

Quantification of Amylase Enzyme from the Calcite Residing Actinobacteria Isolated from Limestone Quarries of Deccan Traps

Syed Raziuddin Quadri^{1,2,3}, Jamal Al Ruwaili¹,
Shaik Mohammed Akbar⁵, Wen-Jun Li^{3,4} and Dayanand Agsar²

¹Department of Medical Laboratory Technology, Faculty of Applied Medical Sciences, Northern Border University, Arar-91431, Northern Borders, Saudi Arabia.

²Department of Microbiology, Gulbarga University, Gulbarga-585106, Karnataka, India.

³Key Laboratory of Microbial Diversity in Southwest China, Ministry of Education and Laboratory for Conservation and Utilization of Bio-resources, Yunnan Institute of Microbiology, Yunnan University, Kunming, Yunnan, 650091, PR China.

⁴State Key Laboratory of Biocontrol and Guangdong Key Laboratory of Plant Resources, College of Ecology and Evolution, Sun Yat-Sen University, Guangzhou 510275, PR China.

⁵Entomology, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Telangana State, India.

(Received: 15 July 2014; accepted: 09 September 2014)

Sixty three actinobacteria were isolated from Limestone quarries of Deccan Traps from South India. The isolates were identified by morphological, biochemical and physiological studies. Among these, six isolates were selected on the basis of their amyolytic activity using submerged cultivation. The potential isolates exhibiting amyolytic activity were DRQ10, DRQ20, DRQ86, DRQ33, DRQ41, and DRQ63 as confirmed by formation of clear zones of starch hydrolysis around the colonies, with maximum activities 2500, 1100, 1200, 1150, 1250, 1400, and 1600 Units, respectively, as determined by DNS (3, 5-dinitrosalicylic acid) assay method. Various physicochemical parameters such as pH, temperature, incubation time were optimized for the production of amylase. Amylase production was maximum on 4th day of incubation. The optimum pH and temperature were found to be 9.0 and 45 °C, and NaCl concentration of 3M for the potential amyolytic isolates. The carbon and nitrogen source for the production of amylase was starch and peptone, respectively.

Key words: Actinobacteria, Amylase, Limestone quarries, Deccan traps.

Actinobacteria are Gram-positive mycelium forming bacteria, capable of producing large variety of different antibiotics and enzymes. Actinobacteria have been isolated from marine¹, as well as terrestrial² sources. Actinobacteria with saprophytic existence occur greatly in natural and manmade environments. Most of them are strictly saprophytes, but some of them form parasitic or mutualistic association with plants and animals. Actinobacteria have become the subject of intensive searches for sources of new, biologically

active compounds, such as, antibiotics, antitumor agents, enzymes, enzyme inhibitors and growth promoting substances. Actinobacteria are filamentous Gram-positive bacteria, characterized by complex life cycle. They belong to the phylum Actinobacteria, which represents one of the largest taxonomic units among the 18 major lineages, currently recognized within the Domain Bacteria³. They play crucial role in recycling refractory biomaterials by decaying complex mixtures of polymers in dead plants, animals and fungal materials. They are also important in soil biodegradation and humus formation as they recycle the nutrients associated with recalcitrant polymers, such as chitin, keratin, and

* To whom all correspondence should be addressed.
Tel +966 594624402, Fax: +966 146644090
E-mail: raziuddinquadri@hotmail.com

lignocelluloses^{4,5}. They produce several volatile substances like geosmin responsible of the characteristic wet earthy odour⁶. The bioactive secondary metabolites produced by microorganisms is reported to be around 23,000, of which 10,000 are produced by actinobacteria, thus representing 45% of all bioactive microbial metabolites discovered⁷.

Large quantities of microorganisms, such as Fungi, Bacteria, Lichen and Algae are seen in a variety of environments in karst areas and they participate in a number of geological and geochemical processes especially the sedimentation and weathering of rocks and minerals and the formation and evolution of soils. Extracellular hydrolytic enzymes secreted by actinobacteria are too important and meriting studies⁸. With the recent advent of biotechnology, there has been a growing interest and demand for enzymes with novel properties. Starch hydrolysis, catalyzed by α -amylase, is one of the most important large scale uses of actinobacteria in terms of enzymatic processes⁹. Many actinobacterial species have been studied for degradation of starch, hemicellulose, lignin and cellulose^{10, 11, 12}. Amylases can breakdown complex sugars such as starch into simple sugars such as glucose, maltose and dextrin. Bacteria, Actinobacteria and Fungi secrete amylases to the outside of the cells to carryout extracellular digestion. α Amylase (endo-1- α -D-glucanohydrolases EC 3.2.1.1) have been widely applied in clinical and medicinal purposes as well as for analytical chemistry. Besides their applications in starch saccharification they are also used in the food, baking, brewing, detergent, textile, paper and distilling industries^{13, 14}. Glucose and maltose-forming α -amylases are employed in alcohol fermentation and sugar syrup formulation, and malto-oligosaccharide-forming α -amylases in food processing^{15, 16}.

The Deccan Volcanic province (DVP) is one of the world's largest igneous provinces and perhaps the best studied continental flood basalt (CFB). It has a total exposed area of about half a million square kilometers, between latitudes 16° - 24° N and longitudes 70° - 77° E. In the northwestern, central and southern Indian peninsula, the approximate volume of the DVP is about 2 x 10⁶ km³ and its estimated age is 64-65 Ma (Million years ago)^{17, 18}. Limestone is naturally

occurring mineral that consists principally of calcium carbonate^{19, 20, 21}. The aim of the present study was to screen amylolytic actinobacteria from the limestone rocks and characterize the enzyme for its stability at harsh conditions like alkalinity and high temperatures to serve the purpose of industrial usage.

MATERIALS AND METHODS

Sample collection from the Deccan traps

Soil samples were collected from limestone quarries around Gulbarga in Karnataka and Betamcherla, Kurnool in Andhra Pradesh, India²².

Isolation of calcite residing Actinobacteria

Samples were screened for actinobacteria by following serial dilution plate culture technique employing starch casein agar and ISP – 2, 3, 6 and 7 media. The inoculated plates were incubated at 35 °C for 3 weeks. Plates were observed at every 24 hours for the growth of actinobacteria. After the completion of incubation period, typical dry powdery colonies were picked up from mixed colonies and sub-cultured on fresh medium to obtain pure cultures. The pure cultures were stored at 4 °C²².

Isolation of amylolytic Actinobacteria

A batch wise bioprocess was carried out in 250 ml Erlenmeyer's flask containing 100 ml starch casein broth. After sterilization of the medium at 121 °C for 15 minutes, 1 ml inoculum with spore count 10⁸ per ml was added. The inoculated flasks were incubated on rotary shaker (180 rpm) at 45 °C for 72 h.

Screening of amylolytic isolates by Starch Iodine plate assay

96 hour cultures were centrifuged at 10,000 rpm, 100 μ l of the supernatants were kept in starch casein agar plates in a well, made in the center with the puncture tube. Following incubation, the plates were flooded with Grams iodine solution and a zone of clearance around culture supernatant of test isolates against a dark blue stained lawn of starch interpreted as a positive result.

Optimization of culture conditions

The various parameters like pH (4.0 to 12.0), temperature (30 °C to 50 °C), incubation time (1 to 6 days), NaCl concentration (0-5 M), carbon (starch, sucrose, maltose, fructose, and xylose) and

nitrogen (casein, oatmeal, peptone, tyrosine, yeast extract) sources were optimized for the production of amylase for the potential actinobacteria isolated from the limestone quarries of Deccan traps.

Determination of extracellular amylase activity

α -Amylase activity was assayed by measuring the release of reducing sugars in a reaction mixture of 1.0 ml of the crude supernatant and 1.0 ml of 1% (w/v) soluble starch (Sigma, MO, USA) solution in 0.05 mol/L phosphate buffer (pH 7.0) incubated at 50 °C for 20 min. Reducing sugars were assayed by the dinitrosalicylic acid (DNS) method at 550 nm spectrophotometrically (Systronics 105)²³. One unit (U) of α -amylase activity corresponded to 1 mmol of glucose equivalent released per minute under the assay conditions²⁴.

RESULTS

Identification of amyolytic isolates

Screening of amyolytic actinobacteria is shown in Fig. 1. Although six isolates were prominent among sixty three actinobacteria that exhibited hydrolysis zones, the actinobacteria DRQ10 strain showed better growth and maximum production of α -amylase.

Influence of incubation time, pH, temperature, NaCl concentration on the production of amylase

Amylase production was maximum on 4th day of incubation. The optimum pH and temperature were found to be 9.0 and 45 °C, respectively, and NaCl concentration of 3 M was required for the potential amyolytic isolates for the production of amylase (Fig. 2, 3, 4, 5).




Fig 1. Synthesis of amylase by six novel isolates of actinobacteria

Table 1. The morphological characters of the selected actinobacterial isolates.

Isolates	Aerial mycelium	Substrate mycelium	Diffusible Pigment	Sporulation pattern
DRQ10	Greyish White	Greyish Brown	Greyish Brown	Flexuous
DRQ20	Brown	Dark Brown	Brownish	Irregular Sporangia and mycelia
DRQ33	Light Pink	Pink	None	Hooked
DRQ41	White	White	None	Straight
DRQ63	White	White	None	Straight
DRQ86	Light Pink	Reddish Pink	None	Hooked /Curled

Effect of carbon and nitrogen source on the production of amylase

The carbon and nitrogen source for the production of amylase was starch and peptone, respectively, for all the potential isolates (Fig. 6, 7).




Fig. 2. Production of amylase by the novel isolates of actinobacteria isolated from the Deccan Traps

DISCUSSION

Actinobacteria have been of major scientific interest in the past decades, with the discovery of large number of metabolites produced by its diverse genera. The present investigation




Fig. 3. Influence of pH on the production of amylase by the novel actinobacterium DRQ10




Fig. 4. Influence of temperature on the production of amylase by the novel actinobacterium DRQ10




Fig. 5. Influence of sodium chloride concentration on the production of amylase by the novel actinobacterium DRQ10




Fig. 6. Influence of carbon source on the production of amylase by the novel actinobacterium DRQ10




Fig. 7. Influence of nitrogen source on the production of amylase by the novel actinobacterium DRQ10

was aimed to isolate amylolytic actinobacteria from the limestone quarries of Deccan traps, from Gulbarga and Betamcherla. The soil scrap of limestone quarries contains a diverse community of organisms differentiated by morphology and biochemical characters. The morphological characters of the selected actinobacterial isolates were shown in Table 1. The enzymatic hydrolysis of starch has been practiced on an industrial scale for many years and is gradually replacing the traditional acid hydrolysis process²⁵. To study the production of α -amylase, the isolates of actinobacteria were inoculated into a suitable medium and morphological characters were studied. The isolates were screened for amylase production on starch agar plates with iodine solution. Amylase producers were confirmed by observing clear zone around the colony (Fig. 1).

High biomass of amylolytic actinobacteria was observed after 96 hour fermentation period for all the potential isolates. The isolate DRQ10 showed highest biomass after 96 hours than the other isolates. The actinobacteria isolates synthesized amylase at pH 9.0, the production decreases slowly beyond pH 9.0, this indicates that the enzyme is optimally produced under alkaline conditions. Alkaline amylases are thought to be important from industrial point of view. Production of amylase was maximum at temperature 45 °C, indicating the thermotolerant nature of the organism. The thermostability of the enzyme can be favourable in industrial operations for traditional brewing and food processing, where temperatures of pasteurization could be used to inactivate the enzymes after fermentation (Stamford et al., 2001). Thermostable amylase has been reported to be active at 57 °C by *Penicillium rugulosum*²⁶. A thermophilic spore-forming strain was isolated from geothermal soil in Antarctica was thermotolerant, and grew well between 45 and 65 °C (optimum at 61 °C)²⁷. *Bacillus subtilis* strain DMO3 grown optimally at 52–55 °C and secret significant amount of alpha-amylase at pH 8²⁸. Amylase production was maximum at 3 M NaCl concentration than in absence of salt, and the production declined with further increase in the salt concentration. Tolerance to high salt concentration designates the halophilic nature of the isolates. Organisms with thermophilic, alkaliphilic and halophilic properties can be explained by the calcite rock

habitat of the isolates and all the isolates showed similar properties. Amylases with thermophilic, alkaliphilic and halophilic characteristics have wide range of industrial applications.

The effect of different carbon and nitrogen sources on amylase production was studied by their addition to nutrient broth. Although the isolates utilized starch, sucrose, maltose, fructose and xylose, starch (0.5%) and sucrose (0.5%) gave significantly higher yield of amylase than other carbon sources added to nutrient broth. Peptone and tyrosine was the most suitable for amylase production than the other nitrogen sources used. Carbon and nitrogen sources are essential for the growth of isolates and optimum production of amylase. The requirement may be explained by the organism's preference for the carbon and nitrogen source for growth rather than for extracellular enzyme formation.

CONCLUSIONS

This study clearly shows that the actinobacteria isolated from the limestone quarries of Deccan Traps possess high potential for producing amylases of thermophilic, alkaliphilic and halophilic nature. Among the sixty three isolates, six isolates were found to possess amylase activity and DRQ10 containing maximum activity.

REFERENCES

1. Selvam K, B. Vishnupriya, VSC Bose., Screening and quantification of marine Actinobacteria producing industrial enzymes amylase, cellulase and lipase from south coast of India, *International Journal of Pharmaceutical & Biological Archives.*, 2011; **2**(5):1481-1487
2. Raghunathan and R Padmadas, 2013. Production, purification and characterization of α -amylase using *Streptomyces* spp. PDS1 and *Rhodococcus* spp. Isolated from Western Ghats. *Int. J Curr Microb. App Sci.*, 2013; **8**: 206-214.
3. Ventura, M.; Canchaya, C.; Tauch, A.; Chandra, G.; Fitzgerald, G.F.; Chater, K.F.; van Sinderen, D. Genomics of Actinobacteria: Tracing the evolutionary history of an ancient phylum. *Microbiol. Mol. Biol. Rev.*, 2007, **71**; 495-548.
4. McCarthy, A.J.; Williams, S.T. Actinobacteria as agents of biodegradation in the environment—a review. *Gene.*, 1992; **115**: 189-192.

5. Stach, J.E.; Bull, A.T. Estimating and comparing the diversity of marine actinobacteria. *Antonie van Leeuwenhoek.*, 2005; **87**: 3-9.
6. Wilkins, K. Volatile metabolites from Actinobacteria. *Chemosphere.*, 1996; **32**: 1427-1434.
7. Berdy, J., Bioactive microbial metabolites. A personal view. *J. Antibiot.*, 2005; **58**: 1-26.
8. a. Pathom-aree W; Y Nogi; IC Sutcliffe; AC Ward; K Horikoshi; AT Bull; Goodfellow M, Williamsia marianensis sp. nov., a novel actinomycete isolated from the Mariana Trench, *Int. J. Syst. Evol. Microbiol.*, 2006; **56**: 1123-1126.
8. b. Pathom-aree W; Y Nogi; AC Ward; K Horikoshi; AT Bull; Goodfellow M, Dermacoccus barathri sp. nov. and Dermacoccus profundus sp. nov., novel actinomycetes isolated from deep-sea mud of the Mariana Trench., *Int. J. Syst. Evol. Microbiol.*, 2006; **56**: 2303-2307.
9. Konsula Z, Liakopoulou-Kyriakides M. Hydrolysis of starches by the action of an α -amylase from Bacillus subtilis., *Proc. Biochem.*, 2004; **39**: 1745-1749.
10. Stamford, T.L.M., Stamford, N.P., Coelho, L.C.B.B., Araujo, J.M., Production and characterization of a thermostable α -amylase from Nocardiosis sp. endophyte of yam bean. *Biores. Technol.*, 2001; **76**: 137-141.
11. Bon EPS, Gottschalk LMF, Nóbrega R. Lignin peroxidase from Streptomyces viridosporus T7A: Enzyme concentration using ultrafiltration. *Appl. Biochem. Biotechnol.*, 2008; **147**: 391-400.
12. Nascimento RP, Alves-Jr. N, Pereira Jr. N, Bon EPS, Coelho RRR. Brewer's spent grain and corn steep liquor as substrate for cellulolytic enzymes production by Streptomyces malaysiensis. *Lett. Appl. Microbiol.*, 2009; **48**: 529-535.
13. Pandey A, Nigam P, Soccol CR, Soccol VT, Singh D and Mohan R. Advances in microbial amylases. *Biotechnol. Appl. Biochem.*, 2000; **31**: 135-152.
14. Saxena RK, Dutt K, Agarwall L, Nayyar P. A highly thermostable and alkaline amylase from a Bacillus sp. PN5. *Biores. Technol.*, 2007; **98**: 260-265.
15. Kirk, O., Borchert, T.V., Fuglsang, C.C., Industrial enzyme application. *Curr. Opin. Biotechnol.*, 2002; **13**: 345-351.
16. Palacios, H.R., Schwarz, P.B., D'Appolonia, L.D., Effect of amylases from different sources on the retrogradation and cyclization of concentrated wheat starch gels: relationship to bread staling. *J. Agr. Food. Chem.*, 2004; **52**, 5978-5986.
17. Glennie, E.A., Gravity Anomalies and the Structure of the Earth's Crust. *Surv. India Prof. Pap.*, 1932; **27**: 10-16.
18. Quershy, M.N., Relation of gravity to elevation and rejuvenation of blocks in India. *J. Geophys. Res.*, 1971; **76**: 545-557.
19. Dunham, R.J., Classification of carbonate rocks according to depositional textures, in Ham W.E. (ed.), Classification of carbonate rocks: Am. Assoc. *Petroleum Geologists Mem*, 1962; **1**: 108-121.
20. Folk, R.L., *Petrology of Sedimentary Rocks*, Hemphill Publishing, Austin, Texas. 1974.
21. Oates, Joseph A. H. Lime and limestone: chemistry and technology, production and uses, Weinheim, Wiley-VCH. 1998.
22. Syed Raziuddin Quadri and Dayanand Agsar., Detection of melanin producing thermo-alkaliphilic Streptomyces from limestone quarries of the Deccan traps., *World Journal of Science and Technology.*, 2012; **2**(2): 08-12
23. Miller L. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal. Chem.*, 1959; **31**: 426-428.
24. Bernfeld P., Amylases, alpha and beta. In: Colowick SP, Kalpan NO (eds.) *Methods in Enzymology*, Academic Press: New York, USA., 1955; **1**: 149-151.
25. Abou-Elela GM, Nermeen, El-Sersy and Wefky SH. Statistical optimization of cold adapted α -amylase production by free and immobilized cells of Nocardiosis aegyptia. *Journal of Applied Science Research.*, 2009; **5**(3): 286-292.
26. Tiwari KL, SK Jadhav, A Fatima., Culture conditions for the production of thermostable amylase by Pencillium rugulosum. *Global journal of Biotechnology and Biochemistry.*, 2007; **2**(1): 21-24.
27. Poli A, E Esposito, L Lama, P Orlando, G Nicolaus, F de Appolonia, A Gambacorta, B Nicolaus, Anoxybacillus amylolyticus sp. nov., a thermophilic amylase producing bacterium isolated from Mount Rittmann (Antarctica), *Systematic and Applied Microbiology.*, 2006; **29**: 300-307.
28. Kishore D and Ashis KM. Characterization of biochemical properties and biological activities of biosurfactants produced by Pseudomonas aeruginosa mucoid and non-mucoid isolates isolated from hydrocarbon-contaminated soil samples. *Appl Microbiol Biotechnol.*, 2005, **69**: 192-199.