

CHAPTER 5

Chickpea and temperature stress: An overview

Viola Devasirvatham¹, Daniel K.Y. Tan¹, Pooran M. Gaur² and Richard M. Trethowan¹

¹Faculty of Agriculture and Environment, Plant Breeding Institute, The University of Sydney, Cobby, NSW, Australia

²International Crops Research Institute for the Semi-Arid Tropics, Patancheru, Telangana, India

5.1 Introduction

Chickpea is an important food grain legume and an essential component of crop rotations throughout the world. However, the adaptation and productivity of chickpea is often limited by low and high temperatures. Cold stress generally occurs in the late vegetative and reproductive stages across the geographical areas of chickpea production. Cold and freezing temperatures (-1.5°C to 15°C) are considered a major problem during the seedling stage of winter-sown chickpea in Mediterranean areas and autumn-sown crops in temperate regions (Singh, 1993). South Australia and parts of north India are most affected by chilling temperatures at flowering (Berger *et al.*, 2011). On the other hand, high day and night temperatures ($>30/16^{\circ}\text{C}$) may cause damage during the reproductive stage on winter-sown chickpea in Mediterranean in-season rainfall areas, south Asia and spring-sown regions (Berger *et al.*, 2011). In chickpea, temperature is a major environmental factor regulating the timing of flowering thus influencing grain yield (Summerfield *et al.*, 1990; Berger *et al.*, 2004). Both low and high temperatures can limit the growth and grain yield of chickpea at all phenological stages.

The FAO climate change technical paper and the Intergovernmental Panel on Climate Change (IPCC) have provided evidence of climate change linked to human activity. Global temperature has been increasing at the rate of 0.74°C per 100 years (IPCC, 2007a). Over the past 50 years, the linear warming trend has been nearly twice the rate of the previous 100 years (FAO, 2009).

Projections to the end of the 21st century estimate a rise in global mean temperature of between 1.8 and 4°C , depending on greenhouse emissions and changes in rainfall patterns (IPCC, 2007a,b). Such changes in climate will impact crop production and some estimates suggest a grain yield decrease of between 8 and 30% (ICRISAT, 2009).

Changes in seasonal temperature and rainfall patterns and their subsequent impact on yield may change the geographic distribution of chickpea production. In Australia, chickpea could expand in new production areas where the frequency of low temperatures ($<15^{\circ}\text{C}$) is higher during early crop growth (Maqbool *et al.*, 2010). However, temperatures lower than 10°C at flowering can reduce grain yield by 15–20% (Chaturvedi *et al.*, 2009). In contrast, the frequency of high temperatures ($>30^{\circ}\text{C}$) during the reproductive stage is often higher in the Australian chickpea production areas of northern New South Wales (NSW) (Devasirvatham *et al.*, 2012a) and any increase in the frequency and duration of these temperatures will limit productivity. A decrease in chickpea yields of 53 kg/ha was observed in north India per 1°C increase in seasonal temperature (Kalra *et al.*, 2008). In south India, the yield loss was estimated to be 10–15% for every 1°C increase beyond the optimum temperature (Upadhyaya *et al.*, 2011). The effect of high and low temperatures on grain quality (grain size and seed coat colour) is also a recognized problem (Wery *et al.*, 1994).

Considerable progress in the improvement of chickpea adaptation to stressful environments has been made. Screening the germplasm in the field and controlled

environments for stress response has increased our knowledge of plant responses to stress in chickpea, and this information has been used in crop improvement. Physiological response (e.g. canopy temperature) and male (pollen) and female (ovary) reproductive function under stress have been investigated to determine their suitability as stress screening techniques (Clarke & Siddique, 2004; Ibrahim, 2011). Sources of tolerance to temperature stress identified using these methods can be used to develop genetic populations to increase our understanding of inheritance. These populations can be used to map quantitative trait loci (QTL) and the resulting linked markers used for marker-assisted selection (MAS). This chapter explores plant responses to high and low temperatures and the implications for stress tolerance breeding in chickpea.

5.2 Impacts on productivity

5.2.1 Temperature stresses during the vegetative period

Cold temperature (<15°C) at emergence reduces crop establishment and results in plants with low vigour. In some sensitive genotypes, cold temperature causes whole plant necrosis and plant death. During chickpea germination, cold temperature also increases susceptibility to soil-borne pathogens thus retarding plant growth and reducing dry matter production (Wery *et al.*, 1994). A plant survival score can be used as an index to describe genotype tolerance to low temperature under field conditions. Chickpea genotypes (Sel 95Th1716 and Sel 96Th11439) were identified as cold tolerant based on plant survival scores in northwestern Iran (Heidarvand *et al.*, 2011). Genotypic differences in chickpea establishment were identified in Australia, and the cold-tolerant cultivars CPI 562896, Semsen and Sombrero showed improved establishment under low temperatures (Wery *et al.*, 1994).

Cold temperatures decrease membrane stability, modify proteins and lipids, and cause changes in respiration and photosynthesis (Croser *et al.*, 2003). Abscisic acid (ABA) content was observed to increase in seedlings at temperatures of 1–7°C compared with the control (23°C) (Nayyar *et al.*, 2005a). The sugar and proline contents also increased under cold stress in the same study. These observations suggest that manipulation of ABA could improve cold tolerance in chickpea.

Field screening for cold tolerance during the late vegetative stage where plants were exposed to –7.4°C for 3 weeks was effective in identifying cold-tolerant genotypes (Malhotra & Singh, 1991). These authors scored materials on a scale of 1 to 9 and concluded that the method was effective in identifying tolerant, moderately tolerant and sensitive genotypes. This method was used to screen cultivated and wild chickpea in the Mediterranean region (Toker, 2005).

Cold temperature generally encourages prolonged vegetative growth in chickpea. Temperature is the main determinant for flower initiation in most environments, although some authors have linked flower initiation in chickpea to a photothermal response (Roberts *et al.*, 1985). In northern NSW, Australia, flower initiation can commence at ≤15°C, although the occurrence of flower abortion will likely be high (Jenkins & Brill, 2011).

The minimum temperature for germination is 10–15°C (Ellis *et al.*, 1986). At high temperatures, greater than 42.5°C, germination decreases significantly (Ibrahim, 2011) and above 45°C no germination is observed due to lack of embryo growth (Singh & Dhaliwal, 1972). Similarly, high temperature affects photosynthesis, transpiration rate and plant growth (Singh & Dhaliwal, 1972) and the length of vegetative period is generally reduced. In other words, phenology can be modified under high temperature (Summerfield *et al.*, 1984). At high temperatures (>35°C), the vegetative period was reduced by 10 to 15 days compared with optimum temperature (28°C) at Kanpur, India (ICRISAT, 2011). High temperature therefore accelerates flowering and reduces the overall crop growth period.

High temperatures can cause cellular abnormalities such as oxidative stress, and denaturation of proteins and enzymes. Oxidative injury occurs as lipid peroxidation, and hydrogen peroxide content tends to increase in heat-sensitive genotypes at day and night temperatures of over 40/30°C compared with heat-tolerant genotypes (Kumar *et al.*, 2012a). ABA remains high at 40/35°C but was observed to decline at 45/40°C (Kumar *et al.*, 2012b). A membrane injury test based on electrolyte leakage from leaves was shown by Ibrahim (2011) to be an effective measure of high temperature sensitivity in chickpea, with sensitive types displaying high degrees of membrane injury. Therefore, heat stress injury can be measured using a combination of oxidative stress assessments, ABA level and membrane injury in chickpea.

Both high and low temperature stresses can affect seed germination, seedling survival, photosynthesis, membrane function, and protein and hormone function. Cold stress encourages a prolonged vegetative period but high temperatures reduce the vegetative period.

5.2.2 Temperature stresses during the reproductive period

Temperature stress at or around flowering is considered a major challenge to yield in many chickpea production areas. Mean daily temperatures at or exceeding 15°C can cause flower abortion (Clarke & Siddique, 1998). Temperatures of less than 10°C during flowering induce flower shedding, low pod set and ultimately poor seed set. Poor pollen viability and germination are the main reasons for low pod set (Savithri *et al.*, 1980). The field and controlled environment screening at ICRISAT identified chickpea genotypic variation for temperature stress during the flowering stage. Plants exposed to mean daily temperatures of 20°C produced more pods than at 15°C. These experiments identified cold-tolerant genotypes such as ICCV 88502 and ICCV 88503 (Srinivasan *et al.*, 1998). Cold temperature also reduces partitioning of assimilates to the vegetative parts, resulting in reduced harvest index (HI). This reduction in harvest index is more common in south Asia and Australia than other production areas (Siddique & Sedgley, 1986; Saxena, 1990).

Poor pod set in chickpea can occur due to the failure of male or female floral parts, or both. Low temperatures at flowering can affect anther dehiscence. Mean daily temperatures of 15°C can also reduce anther dehiscence and pollen load on stigma (Srinivasan *et al.*, 1999). However, at a similar temperature, pollen viability and pollen germination on the stigma were higher in the tolerant lines ICCV 88501, ICCV 88502 and ICCV 88503 than in the sensitive cultivars Chafa and Annigeri (Srinivasan *et al.*, 1999). Pollen function was clearly more sensitive to temperature change than pistil function (esterase activity). Clarke and Siddique (2004) showed that pollen viability and pollen germination on the stigma were the primary reasons why pod set in chickpea was reduced during low-temperature stress. Pollen sensitivity to low temperatures was identified at 5 and 9 days before anthesis (Clarke, 2001). Clarke and Siddique (2004) and Srinivasan *et al.* (1999) also observed that low temperature did not affect the pistil function, i.e. esterase activity. However, pollen tube

growth on the styles of sensitive genotypes was retarded due to cold temperatures. Short pollen tube length at low temperature (15°C) in *in vitro* germination tests was observed at 20°C and 25°C by Savithri *et al.* (1980). Ultimately pod set is reduced by low temperature as observed by Srinivasan *et al.* (1999). They found that pod set was reduced at a low temperature regime of 15/5°C compared with the control (25/15°C).

High temperature during the reproductive stage is a major cause of yield loss due to partial or complete pollen sterility. In chickpea, temperatures at or exceeding 35°C affected male reproductive tissue (anther and pollen), function (pollen germination and tube growth) and pod set. Both anther and pollen showed more structural abnormalities under stress including changes in anther locule number, anther epidermis wall thickening, and pollen sterility rather than functional abnormalities (e.g. *in vivo* pollen germination) (Figure 5.1) (Devasirvatham *et al.*, 2013). Pollen abnormalities can also be found at high temperature, including leakage of pollen protoplast, zigzag pollen tube growth, pseudo-germination and bulbous tip formation in the pollen tube (Devasirvatham *et al.*, 2013). Heat-tolerant chickpea genotypes had clear pollen tube growth on the style following pollen germination and this was confirmed by pod set (Figure 5.2). In heat-sensitive genotypes no pollen germination on the stigma was observed due to complete pollen sterility at temperatures at or exceeding 35°C (Devasirvatham *et al.*, 2012b, 2013) (Figure 5.2). Pollen sterility in the heat-sensitive genotypes is a function of lower sucrose levels, resulting in poor pollen function and pod set (Kaushal *et al.*, 2013).

At very high temperature (45/35°C) both pollen fertility and stigma function can be affected. Observations by Kumar *et al.* (2012b) indicate that oxidative stress in the leaves results in poor fertilization. Devasirvatham *et al.* (2012b, 2013) concluded that the critical temperature affecting pod set was $\geq 37^\circ\text{C}$ for heat-tolerant genotypes (such as ICCV 92944, ICC 1205 and ICC 15614) and $>33^\circ\text{C}$ for heat-sensitive genotypes (ICC 5912, ICC 4567 and ICC 10685).

5.2.3 Temperature stresses during post-anthesis period

Post-anthesis temperature stress, particularly after commencement of pod set, can cause significant pod abortion and decreased grain filling. In chickpea, cold stress decreased the rate and duration of grain filling and




Figure 5.1 Heat-sensitive genotype anther structural abnormalities: anther stained with Alexander's stain. (a) Locule number changed (ICC 4567). (b) Anther epidermis wall is thickened (ICC 4567). (c) Anther shows fertile and sterile pollen grain (ICC 5912). Fertile – red in colour; sterile – green in colour. Scale: 10 μm.

produced smaller seeds (Nayyar *et al.*, 2007; Kaur *et al.*, 2008). At 13/5°C, chickpea average seed weight and size decreased by 41% and 24%, respectively, compared with 28/17°C, largely because seed filling duration reduced from 20 days (non-stressed) to 14 days (cold stressed) (Nayyar *et al.*, 2005b). Similarly, low post-anthesis

temperature reduced yield by 1.3 t/ha in northern NSW, Australia, during 2009 (Moore *et al.*, 2010).

In a controlled environment study, Wang *et al.* (2006) reported a grain yield reduction of 33–39% for post-anthesis heat stress compared with pre-anthesis heat stress. This was possibly due to poor remobilization of




Figure 5.2 Effect of high temperatures on the pollen germination on the stigma. (a) Heat-tolerant: ICC 15614 – pollen germination on the stigma. (b) Heat-sensitive: ICC 10685 – no pollen germination on the stigma. Reproduced from Devasirvatham *et al.* (2013).



Figure 5.3 Comparison of seed size under heat stress. Larger seeds (left side) from non-stressed and smaller seeds (right side) from heat-stressed conditions. Photo courtesy of V. Devasirvatham.

photosynthates to the grain, thus lowering seed weight and seed number per plant (Wang *et al.*, 2006). Both temperature stress extremes influenced seed shape and seed coat colour (Figure 5.3). Generally, temperature stress reduces cotyledon cell number, cell expansion, grain filling rate and ultimately seed weight (Munier-Jolain & Ney, 1998).

Temperature stress can influence grain filling by altering the concentration of hormones, particularly abscisic acid (ABA) and enzymes, in plant tissue. As discussed earlier, ABA plays a significant role in both cold and heat stress tolerance in chickpea (Nayyar *et al.*, 2005a; Kumar *et al.*, 2012a) and is generally downregulated under stress. Exogenous application of ABA increased tolerance to cold stress by improving survival rate through the reproductive stage (Nayyar *et al.*, 2005a). The exogenous application of ABA decreased electrolyte leakage and increased pollen viability,

germination, flower retention, pod set, seed size and grain yield (Kumar *et al.*, 2008). Similarly, Kumar *et al.* (2012b) showed that ABA treatment reduced oxidative injury in chickpea under high temperature. Clearly, exogenous application of ABA will improve grain filling under temperature stress and hence grain yield.

5.3 Impacts on nutritional and processing quality

Environmental stresses during seed development have a negative effect on the quality of chickpea seeds (Behboudian *et al.*, 2001). However, comparatively few studies have dealt with the effect of temperature on seed development and quality in chickpea. Nayyar *et al.* (2005b) reported that under cold stress grain sugar

concentration increased in chickpea but the accumulation of storage proteins, starch and several amino acids decreased. However, the effect was influenced by the stage of seed development. There was a greater reduction of starch, proteins, soluble sugars, fat, crude fibre and storage protein fractions when cold stress occurred in late pod-filling compared to early pod-filling stages (Kaur *et al.*, 2008). However, seed germination was inhibited when plants were stressed at early pod-filling.

The effects of high temperature stress were generally similar to cold stress. High sucrose synthase and low invertase activity were observed in the seeds of heat-tolerant genotypes compared with heat-sensitive types during early pod filling (Chickpea Technical Report, 2011). Generally, high temperature during grain filling reduces dough and baking quality in grain crops (Stone & Nicolas, 1994). However, the available information on grain quality under temperature stresses in chickpea is limited. There is clearly a need to extend our knowledge of grain quality including baking quality under both high and low temperature stresses.

5.4 Breeding for tolerance to temperature stresses

Chickpea improvement has focused on yield potential and regional adaptation through resistance and tolerance to abiotic and biotic stresses, plant type and grain characteristics. At present, the selected bulk method is the most common selection technique used in chickpea breeding (Gaur *et al.*, 2007). The selected bulk method is relatively inexpensive to employ and the response to selection is generally not inferior to more labour-intensive methods such as pedigree selection (Salimath *et al.*, 2007). This section describes some of the breeding strategies used to improve temperature tolerance in chickpea and explores options for future breeding.

5.4.1 High temperature tolerance

A simple but effective field screening technique for heat tolerance at the reproductive stage in chickpea has been developed at ICRISAT (Gaur *et al.*, 2013, 2014). It involves advancing sowing date to synchronize the reproductive phase of the crop with the occurrence of higher temperatures ($\geq 35^{\circ}\text{C}$). This method was effective in identifying heat-tolerant

germplasm at ICRISAT and several other locations in India (Gaur *et al.*, 2013, 2014).

A few heat-tolerant chickpea cultivars (ICCV 88512 and ICCV 513) were identified more than a decade ago (Dua, 2001). However, heat tolerance research in chickpea has only received significant attention in recent years. More recently, Krishnamurthy *et al.* (2011) identified 18 stable heat-tolerant genotypes (e.g. ICC 1205, ICC 637 and ICC 15618) by field screening a reference set of chickpea from southern and central Indian field trials. Short-duration, high-yielding, heat-tolerant genotypes (ICC 5597, ICC 5829, ICC 6121, ICC 7410, ICC 11916, ICC 13124, ICC 14284, ICC 14368 and ICC 14653) were identified by Upadhyaya *et al.* (2011). A heat-tolerant breeding line, ICCV 92944, has been released in Myanmar (as Yezin 6) and in India (as JG 14) and is performing well under late-sown conditions (Gaur *et al.*, 2013). Several breeding lines with higher yields under heat stress than the standard cultivar ICCV 92944 have been identified (Gaur *et al.*, 2013, 2014). Outside India, Kaloki (2010) identified ICCV 92318 as a source of heat tolerance in the semi-arid environments of Kenya through the African Climate Change Breeding Program.

Devasirvatham *et al.* (2012b) confirmed the heat tolerance of ICCV 92944 using a pollen selection method. Devasirvatham *et al.* (2013) also confirmed the heat tolerance of germplasm identified earlier by Krishnamurthy *et al.* (2011) (ICC 1205, ICC 15614) using pollen viability in the field and controlled environment studies, and suggested using this technique to develop heat-tolerant cultivars. These materials have been incorporated into chickpea improvement at ICRISAT and new heat-tolerant progeny are under development as genetic mapping populations (Gaur *et al.*, 2013). Diversity arrays technology (DArT) (Mace *et al.*, 2008) markers with good genome coverage were associated with traits targeted for high temperature tolerance in chickpea, and many genomic regions linked with phenology and grain yield have been identified (Devasirvatham, 2012), thus demonstrating the feasibility of applying genetic association analysis to explore complex traits in future. While there is clearly significant variation for high temperature tolerance in adapted chickpea, there is a compelling need to extend the search for new genetic diversity to provide additional allelic variation for temperature tolerance. The wild annual *Cicer* sp. is a possible source of variation and

could be exploited. This new allelic variation would allow plant breeders to lift the current reproductive temperature limits on chickpea.

5.4.2 Low temperature tolerance

Low temperature stress breeding generally aims to develop materials adapted to the temperature range -1.5 to 15°C at the reproductive stage and less than -1.5°C at the vegetative growth (Croser *et al.*, 2003). Different sources of resistance to cold tolerance are reported by Chaturvedi *et al.* (2009), and several cold-tolerant breeding lines such as ICCVs 88502, 88503, 88506, 88510 and 88516 have been developed that set pods at less than 15°C in India (ICRISAT, 1994). The Indian Agricultural Research Institute (IARI) has also developed a few cold-tolerant genotypes (BGD 112 green, BG 1100, BG 1101, PUSA 1103, BGD 1005, PUSA 1108, DG 5025, DG 5027, DG 5028, DG 5036 and DG 5042) (Gaur *et al.*, 2007). Using pollen as a selection method, Clarke *et al.* (2004) confirmed the cold tolerance of ICCV 88516 and 88510 and the sensitivity of Amethyst, Dooen, Tyson and FLIP84-15C in Western Australia. Accessions of cultivated and wild *Cicer* sp. were screened for cold tolerance at ICARDA (Singh *et al.*, 1995). These authors reported cold tolerance in the lines ILC 8262, ILC 8617 (a mutant) and a FLIP 97-82C from cultivated *Cicer* along with wild annual chickpea such as *C. bijugum* and *C. reticulatum*.

Later, Toker (2005) identified chilling tolerance ($<-1.5^{\circ}\text{C}$) in annual wild *Cicer* sp. of *yamashitae*. Heidarvand *et al.* (2011) identified the genotypes Sel 95Th1716 and Sel 96Th11439 as chilling tolerant based on field screening at the vegetative stage where plants were exposed to -11°C to -25°C at the Dryland Agriculture Research Institute (DARI) of Iran.

Both additive and non-additive gene effects govern cold tolerance in chickpea. Cold tolerance was observed to be dominant over susceptibility for at least five sets of genes (Malhotra & Singh, 1990). Breeding at ICARDA has resulted in the expansion of genetic variability for flowering at low temperatures using cultivated \times wild *Cicer* crosses. The genes responsible for flowering at low temperature have been transferred from wild to cultivated lines (Chaturvedi *et al.*, 2009). These reports suggested that wild relatives of chickpea can be used as a source of tolerance to low temperatures in applied breeding.

5.5 Conclusions

Both high and low temperature stresses cause grain yield loss. Cold stress encourages a prolonged vegetative period while high temperatures reduce the duration of the vegetative period. Reduced pollen viability and pollen germination on the stigma are the primary causes of poor pod set in chickpea following low temperature stress. Similarly, high temperature stress disrupts pollen viability and anther dehiscence. However, stigma receptivity is not affected by either stress. The rate and duration of seed filling are both decreased by cold and high temperature stresses.

Recent chickpea breeding programmes targeting both high and low temperature stresses have been initiated by many countries including India, Australia and Canada, with global centres such as ICARDA and ICRISAT supporting the wider effort through the characterization and exploitation of genetic resources. Screening for tolerance to temperature stresses has identified many promising sources of tolerance to both high and low temperature in chickpea. However, field-based screening is generally based on delayed sowing, and biomass development and the length of the vegetative phase are reduced in such treatments, thus reducing the fitness of plants to survive temperature extremes at flowering. Field-based methods that impose a temperature stress on a normally grown plant should be developed to confirm and validate the response of chickpea lines already identified. The identification of QTLs for temperature stress tolerance and linked molecular markers will undoubtedly improve rates of genetic advance and marker-assisted selection can easily be incorporated into most breeding methods.

Rapid progress has been made in the development of genomic resources for chickpea, and breeders have already started integrating molecular breeding strategies such as marker-assisted backcrossing (MABC) and marker-assisted recurrent selection (MARS) to improving drought tolerance in chickpea (Gaur *et al.*, 2012). Advances in marker systems and genotyping technologies such as DaRT and single nucleotide polymorphisms (SNPs) and genotyping by sequencing (GBS) have made genotyping large numbers of materials cost efficient. The integration of genomic technologies in chickpea breeding will greatly improve efficiency of developing chickpea cultivars that are more resilient to changes in temperature. For example, MARS recombines significant gene effects found among

the progeny of a single population in chickpea (Cobos *et al.*, 2007) and will be useful in enhancing tolerance to temperature extremes. Another potential approach is genomic selection (Nayak *et al.*, 2010), where training populations that are representative of the wider gene pool are assembled, genotyped and phenotyped and all the estimated gene effects used to assemble new chickpea cultivars. However, these breeding-by-design approaches are completely dependent upon the accuracy of the phenotyping data used in estimating the gene effects. The screening methods outlined in this chapter offer scope for rapid and accurate phenotyping for chickpea temperature stress tolerance and when integrated in a molecular breeding scheme, should provide the temperature-tolerant chickpea cultivars required for an increasingly hostile production environment.

References

- Bebhoudian MH, Ma Q, Turner NC, Palta JA (2001) Reactions of chickpea to water stress: yield and seed composition. *J Sci Food Agric* 81: 1288–1291.
- Berger JD, Turner NC, Siddique KHM, *et al.* (2004) Genotype by environment studies across Australia reveal the importance of phenology for chickpea (*Cicer arietinum* L.) improvement. *Crop Pasture Sci* 55: 1071–1084.
- Berger JD, Milroy SP, Turner NC, Siddique KHM, Imtiaz M, Malhotra R (2011) Chickpea evolution has selected for contrasting phenological mechanisms among different habitats. *Euphytica* 180: 1–15.
- Chaturvedi SK, Mishra DK, Vyas P, Mishra N (2009) Breeding for cold tolerance in chickpea. *Trends Biosci* 2: 1–6.
- Chickpea Technical Report (2011) Improving heat tolerance in chickpea for enhancing its productivity in warm growing conditions and mitigating impacts of climate change. ICRIASAT, Patancheru, Andhra Pradesh, India, pp. 1–46
- Clarke HJ (2001) Improving tolerance to low temperature in chickpea. In: *4th European Conference on Grain Legumes. Towards the Sustainable Production of Healthy Food, Feed and Novel Products*. Cracow, Poland, July 8–12, 2001, pp. 34–35.
- Clarke HJ, Siddique KHM (1998) Growth and development. In: Loss S, Brandon N, Siddique KHM (eds), *The chickpea*. Western Australia Agriculture Bulletin 1326. Department of Western Australia Agriculture.
- Clarke HJ, Siddique KHM (2004) Response of chickpea genotypes to low temperature stress during reproductive development. *Field Crops Res* 90: 320–334.
- Clarke HJ, Khan TN, Siddique KHM (2004) Pollen selection for chilling tolerance at hybridization leads to improved chickpea cultivars. *Euphytica* 139: 65–74.
- Cobos MJ, Rubio J, Fernandez-Romero MD, *et al.* (2007) Genetic analysis of seed size, yield and days to flowering in a chickpea recombinant inbred line population derived from a kabuli × desi cross. *Ann App Biol* 151: 33–42.
- Croser JS, Clarke HJ, Siddique KHM, Khan TN (2003) Low-temperature stress: Implications for chickpea (*Cicer arietinum* L.) improvement. *Crit Rev Plant Sci* 22: 185–219.
- Devasirvatham V (2012) The basis of chickpea heat tolerance under semi-arid environments. PhD thesis, The University of Sydney, NSW, Australia.
- Devasirvatham V, Tan DKY, Gaur PM, Raju TN, Trethowan RM (2012a) High temperature tolerance in chickpea and its implications for plant improvement. *Crop Pasture Sci* 63: 419–428.
- Devasirvatham V, Gaur PM, Mallikarjuna N, Raju TN, Trethowan RM, Tan DKY (2012b) Effect of high temperature on the reproductive development of chickpea genotypes under controlled environments. *Funct Plant Bio* 39: 1009–1018.
- Devasirvatham V, Gaur PM, Mallikarjuna N, Raju TN, Trethowan RM, Tan DKY (2013) Reproductive biology of chickpea response to heat stress in the field is associated with the performance in controlled environments. *Field Crops Res* 142: 9–19.
- Dua RP (2001) Genotypic variations for low and high temperature tolerance in gram (*Cicer arietinum*). *Indian J Agri Sci* 71: 561–566.
- Ellis RH, Covell S, Roberts EH, Summerfield RJ (1986) The influence of temperature on seed germination rate in grain legumes. XI. Intraspecific variation in chickpea (*Cicer arietinum* L.) at constant temperatures. *J Exp Bot* 37: 1503–1515.
- FAO (2009) *The State of World Fisheries and Aquaculture 2008*. Also available at: <ftp://ftp.fao.org/docrep/fao/011/i0250e/i0250e01.pdf>. UN Food and Agriculture Organization, Rome.
- Gaur PM, Gowda CLL, Knights EJ, *et al.* (2007) Breeding achievements. In: Yadav SS, Redden RJ, Chen W, Sharma B (eds), *Chickpea Breeding and Management*. pp. 391–416. CAB International, Wallingford, Oxon, UK.
- Gaur PM, Jukanthi AK, Varshney RK (2012) Impact of genomic technologies on chickpea breeding strategies. *Agron* 2: 199–221.
- Gaur PM, Chaturvedi SK, Kumar S, *et al.* (2013) High temperature tolerance in food legumes to mitigate impact of climate change. In: *Proceedings of the National Conference on Crop Improvement and Adaptive Strategies to Meet Challenges of Climate Change*, 22–24 February 2013, University of Agricultural Sciences, Bangalore, India, pp. 29–32.
- Gaur PM, Jukanti AK, Srinivasan S, *et al.* (2014) Climate change and heat stress tolerance in chickpea. In: Tuteja N, Gill SS (eds), *Climate Change and Plant Abiotic Stress Tolerance*. Wiley-VCH, Weinheim, Germany, pp. 839–855.
- Heidarvand L, Amri RM, Naghavi MR, Farayed Y, Sadeghzadeh B, Alizadeh KH (2011) Physiological and morphological characteristics of chickpea accessions under low temperature. *Russian J Plant Phys* 58: 157–163.

- Ibrahim HM (2011) Heat stress in food legumes: evaluation of membrane thermostability methodology and use of infra-red thermometry. *Euphytica* 180: 99–105.
- ICRISAT (1994) Cold tolerant chickpea varieties. Plant material description no. 53. International Crops Research Institute for the Semi-arid Tropics, Patancheru, Hyderabad, Andhra Pradesh, India. pp. 1–4.
- ICRISAT (2009) Climate change in the semi-arid tropics. Available at: www.icrisat.org./rds/Climate_Change_SAT_flyer.pdf.
- ICRISAT (2011) Improving heat tolerance in chickpea for enhancing its productivity in warm growing conditions and mitigating impacts of climate change. Technical report. International Crops Research Institute for the Semi-arid Tropics, Patancheru, Hyderabad, Andhra Pradesh, India.
- IPCC (2007a) *Climate Change 2007: the Physical Science Basis. Summary for Policymakers*. WMO/UNEP, Paris.
- IPCC (2007b) Summary for policymakers. In: Parry ML, Canziani M, Palutikof O, van der Linden J, Hanson C (eds), *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK, pp. 7–22.
- Jenkins L, Brill R (2011) Chickpea time of sowing trial. DAN00156: Northern Pulse Agronomy Project. NSW DPI, Australia.
- Kaloki P (2010) Sustainable climate change adaptation options in agriculture: The case of chickpea in the semi-arid tropics of Kenya. Report of the African Climate Change Fellowship Program. International Crops Research Institute for the Semi-Arid Tropics, Nairobi, Kenya, pp. 1–54.
- Kalra N, Chakraborty D, Sharma A, et al. (2008) Effect of temperature on yield of some winter crops in northwest India. *Curr Sci* 94: 82–88.
- Kaur G, Kumar S, Nayyar H, Upadhyaya HD (2008) Cold stress injury during the pod-filling phase in chickpea (*Cicer arietinum* L.): Effects on quantitative and qualitative components of seeds. *J Agron Crop Sci* 194: 457–464.
- Kaushal N, Awasthi R, Gupta K, Gaur PM, Siddique KHM, Nayyar H (2013) Heat stress induced reproductive failures in chickpea (*Cicer arietinum* L.) are associated with impaired sucrose metabolism in leaves and anthers. *Funct Plant Bio* 40: 1334–1349.
- Krishnamurthy L, Gaur PM, Basu PS, et al. (2011) Large genetic variation for heat tolerance in the reference collection of chickpea (*Cicer arietinum* L.) germplasm. *Plant Gene Res* 9: 59–61.
- Kumar S, Kaur G, Nayyar H (2008) Exogenous application of abscisic acid improves cold tolerance in chickpea (*Cicer arietinum* L.). *J Agron Crop Sci* 194: 449–456.
- Kumar S, Thakur P, Kaushal N, Malik JA, Gaur P, Nayyar H (2012a) Effect of varying high temperatures during reproductive growth on reproductive function, oxidative stress and seed yield in chickpea genotypes differing in heat sensitivity. *Arch Agron Soil Sci* 59: 823–843.
- Kumar S, Kaushal N, Nayyar H, Gaur PM (2012b) Abscisic acid induces heat tolerance in chickpea (*Cicer arietinum* L.) seedlings by facilitated accumulation of osmoprotectants. *Acta Phys Plan* 34: 1651–1658.
- Mace ES, Xia L, Jordan DR, et al. (2008) DarT markers: diversity analyses and mapping in *Sorghum bicolor*. *BMC Genomics* 9: 1–11.
- Malhotra RS, Singh KB (1990) The inheritance of cold tolerance in chickpea. *J Gene Plant Bre* 44: 227–230.
- Malhotra RS, Singh KB (1991) Gene action for cold tolerance in chickpea. *Theor Appl Genet* 82: 598–601.
- Maqbool A, Shafiq S, Lake L (2010) Radiant frost tolerance in pulse crops – a review. *Euphytica* 172: 1–12.
- Moore K, Jenkins L, Hertel K, Callaghan G (2010) Chickpea in 2010: Low temperature effects in 2009. GRDC update paper. Available at: www.grdc.com.au
- Munier-Jolain NG, Ney B (1998) Seed growth rate in grain legumes II. Seed growth rate depends on cotyledon cell number. *J Exp Bot* 49: 1971–1976.
- Nayak SN, Zhu H, Varghese N, et al. (2010) Integration of novel SSR and gene-based SNP marker loci in the chickpea genetic map and establishment of new anchor points with *Medicago truncatula* genome. *Theor Appl Genet* 20: 1415–1441.
- Nayyar H, Bains T, Kumar S (2005a) Low temperature induced flower abortion in chickpea: relationship to abscisic acid and cryoprotectants in reproductive organs. *Environ Exp Bot* 53: 39–47.
- Nayyar H, Bains TS, Kumar S (2005b) Chilling effects during seed filling on accumulation of seed reserves and grain yield. *J Sci Food Agric* 85: 1925–1930.
- Nayyar H, Kaur G, Kumar S, Upadhyaya HD (2007) Low temperature effects during seed filling on chickpea genotypes (*Cicer arietinum* L.): Probing mechanisms affecting seeds reserves and yield. *J Agron Crop Sci* 193: 336–344.
- Roberts EH, Hadley P, Summerfield RJ (1985) Effects of temperature and photoperiod on flowering in chickpeas (*Cicer arietinum* L.). *Ann Bot* 55: 881–892.
- Salimath PM, Toker C, Sandhu JS, et al. (2007) Conventional breeding methods. In: Yadav SS (ed.), *Chickpea Breeding and Management*. CAB International, Wallingford, UK, pp. 369–390.
- Savithri KS, Ganapathy PS, Sinha SK (1980) Sensitivity to low temperature in pollen germination and fruit-set in *Cicer arietinum* L. *J Exp Bot* 31: 475–481.
- Saxena MC (1990) Problems and potentials in chickpea production in the nineties. In: Van Rheenen HA, Saxena MC (eds), *Chickpea in the Nineties. Proceedings of the 2nd International Workshop on Chickpea Improvement*, ICRISAT, Patancheru, India, pp. 13–25.
- Siddique KHM, Sedgley RH (1986) Chickpea (*Cicer arietinum* L.), a potential grain legume for southwestern Australia: seasonal growth and yield. *Crop Pasture Sci* 37: 245–261.
- Singh KB (1993) Problems and prospects of stress resistance breeding in chickpea. In: Singh KB, Saxena MC (eds), *Breeding*

- for Stress Tolerance in Cool-Seasons Food Legumes. John Wiley & Sons, Ltd., Chichester, UK, pp. 17–37.
- Singh KB, Malhotra RS, Saxena MC (1995) Additional sources of tolerance to cold in cultivated and wild *Cicer* species. *Crop Sci* 35: 1491–1497.
- Singh NH, Dhaliwal GS (1972) Effect of soil temperature on seedling emergence in different crops. *Plant Soil* 37: 441–444.
- Srinivasan A, Johansen C, Saxena NP (1998) Cold tolerance during early reproductive growth of chickpea (*Cicer arietinum* L.): Characterization of stress and genetic variation in pod set. *Field Crops Res* 57: 181–193.
- Srinivasan A, Saxena NP, Johansen C (1999) Cold tolerance during early reproductive growth of chickpea (*Cicer arietinum* L.): genetic variation in gamete development and function. *Field Crops Res* 60: 209–222.
- Stone PJ, NicoLas ME (1994) Wheat cultivars vary widely in their responses of grain yield and quality to short periods of post-anthesis heat stress. *Aust J Plant Physiol* 21: 887–900.
- Summerfield RJ, Hadley P, Roberts EH, Minchin FR, Rawsthorne S (1984) Sensitivity of chickpea (*Cicer arietinum* L.) to hot temperatures during the reproductive period. *Exp Agric* 20: 77–93.
- Summerfield RJ, Virmani SM, Roberts EH, Ellis RH (1990) Adaption of chickpea to agroclimatic constraints. In: van Rheenen HA, Saxena MC (eds), *Chickpea in the Nineties. Proceedings of the Second International Workshop on Chickpea Improvement, 4–8 December 1989*. ICRISAT Center, Patancheru, Andhra Pradesh, India, pp. 50–61.
- Toker C (2005) Preliminary screening and selection for cold tolerance in annual wild *Cicer* species. *Genet Resour Crop Evol* 52: 1–5.
- Upadhyaya HD, Dronavalli N, Gowda CLL, Singh S (2011) Identification and evaluation of chickpea germplasm for tolerance to heat stress. *Crop Sci* 51: 2079–2094.
- Wang J, Gan YT, Clarke F, McDonald CL (2006) Response of chickpea yield to high temperature stress during reproductive development. *Crop Sci* 46: 2171–2178.
- Wery J, Silim SN, Knights EJ, Malhotra RS, Cousin R (1994) Screening techniques and sources of tolerance to extremes of moisture and air temperature in cool season food legumes. *Euphytica* 73: 73–83.