

## Entomology

### Host Plant Resistance to Insects in Sorghum: Present Status and Need for Future Research

HC Sharma<sup>1\*</sup>, Belum VS Reddy<sup>1</sup>, MK Dhillon<sup>1</sup>, K Venkateswaran<sup>2</sup>, BU Singh<sup>3</sup>, G Pampapathy<sup>1</sup>, RT Folkertsma<sup>1</sup>, CT Hash<sup>1</sup> and KK Sharma<sup>1</sup>  
(1. ICRISAT, Patancheru 502 324, Andhra Pradesh, India; 2. National Bureau of Plant Genetic Resources, Regional Station, Rajendranagar, Hyderabad 500 030, Andhra Pradesh, India; 3. National Research Center for Sorghum, Rajendranagar, Hyderabad 500 030, Andhra Pradesh, India)

\*Corresponding author: h.sharma@cgiar.org

#### Introduction

Sorghum [*Sorghum bicolor* (L.) Moench] is an important cereal crop in Asia, Africa, the Americas, and Australia. Grain yields on farmers' fields in the semi-arid tropics (SAT) are generally low (500 to 800 kg ha<sup>-1</sup>) mainly due to insect pest damage. Nearly 150 insect species have been reported as pests on sorghum, of which sorghum shoot fly (*Atherigona soccata* Rond.), stem borer (*Chilo partellus* Swin.), armyworm (*Mythimna separata* Walk.), shoot bug (*Peregrinus maidis* Ashmead), aphid (*Melanaphis sacchari* Zehnt.), sorghum midge (*Stenodiplosis sorghicola* Coq.), earhead bug (*Calocoris angustatus* Leth.), and head caterpillars (*Helicoverpa*, *Eublemma*, and *Cryptoblabes*) are the major pests (Sharma 1993). Annual losses due to insect pests in sorghum have been estimated to be over \$1089 million in the SAT.

#### Host plant resistance

**Techniques to screen for resistance to insect pests.** The ability to develop insect-resistant cultivars, use of marker-assisted selection, and development of transgenic plants with insect resistance depends on the precision of resistance screening techniques. Infester row, cage and leaf disc screening techniques have been standardized to evaluate sorghum germplasm, breeding material, and mapping populations for resistance to insect pests under field and greenhouse conditions (Sharma et al. 1992, 2003). However, several of these techniques are being used in the sorghum improvement programs only sparingly because of lack of resources or lack of enthusiasm on the part of the scientists involved. Lack of infrastructure for insect rearing could be an impediment to screening for resistance to stem borer, but infester row and no-choice


cage techniques developed to screen for resistance to shoot fly, midge, and head bugs do not require much investment. There is a need to standardize the techniques to screen for resistance to aphids – an emerging pest problem. One of the problems in selecting for resistance to stem borer is the relative importance of foliar injury, deadhearts, stem tunneling, exit holes, and tiller production (Singh 2002). The effects of different damage parameters on grain yield loss are not fully understood. Another important question is whether the material should be screened in each generation, alternate generations, or only after the material has become homozygous in F<sub>5</sub> – F<sub>6</sub> generations. Extensive studies at ICRISAT have indicated that the material subjected to borer infestation in F<sub>2</sub> to F<sub>5</sub> generations had greater frequency of resistant progenies than the material exposed to borer infestation in the F<sub>5</sub> generation only. Such information needs to be generated for different insect pests.

**Identification and utilization of sources of resistance to insect pests.** Host plant resistance should form the backbone of pest management in sorghum. Over the past five decades, a large proportion of the world sorghum germplasm collection has been evaluated for resistance to insect pests, and a number of lines with resistance to major insect pests have been identified (Sharma et al. 1992, 2003). Large-scale screening of the sorghum germplasm at ICRISAT has resulted in identification of several lines with reasonable levels of resistance to shoot fly, stem borer, midge, and head bugs (Table 1). Sources of resistance to insects in sorghum have been used in the breeding program, and many varieties with resistance to insect pests have been developed (Table 2). However, cultivars with resistance to insect pests are cultivated by farmers only on a limited scale due to an overemphasis by national programs on grain yield as a criterion for release of cultivars. Since sorghum varieties and hybrids with a yield potential of over 10 t ha<sup>-1</sup> are already available in the market, it is important that insect and disease resistance be used as a criterion to identify varieties and hybrids for use by farmers for sustainable crop production.

**Diversification of the cytoplasmic male-sterile systems with resistance to insect pests.** Most of the hybrids grown in India are based on *milo* cytoplasm (A<sub>1</sub> cytoplasm), which is highly susceptible to sorghum shoot fly (Dhillon 2004) (Fig. 1). Extensive use of the A<sub>1</sub> cytoplasm as a source of cytoplasmic male-sterility (CMS) has resulted in narrowing of the genetic base of sorghum hybrids currently being cultivated by farmers, and this might increase the vulnerability of this crop to biotic and abiotic stresses. In general, the CMS lines are more susceptible to sorghum shoot fly (*A. soccata*), sugarcane aphid (*M. sacchari*), shoot bug (*P. maidis*), and midge

(*S. sorghicola*) than the maintainer lines, suggesting that the maintainer lines harbor the factors that influence expression of resistance to insects (Sharma et al. 2004b). Therefore, there is a need to develop a range of CMS, maintainer, and restorer lines with resistance to insect pests, and diversify the CMS systems in sorghum. The A<sub>4</sub>M cytoplasm is slightly less susceptible to shoot fly than the other CMS systems. Recovery from shoot fly damage is better in A<sub>4</sub>M, A<sub>3</sub>, and A<sub>2</sub> cytoplasm than the A<sub>1</sub> cytoplasm. Shoot fly survival and development is also poor on A<sub>4</sub>M and A<sub>4</sub>VzM CMS systems. The A<sub>4</sub>M cytoplasm being less susceptible to shoot fly and having better recovery resistance, can be exploited for developing shoot fly-resistant hybrids in future. However, as a first step, it may be better to transfer the traits associated with resistance to shoot fly into the hybrid parents in A<sub>1</sub> cytoplasm. Another alternative would be to explore opportunities for using male gametocytes and/or temperature and photoperiod-induced male-sterility for sorghum hybrid seed multiplication as these might allow exploitation of the favorable effects of normal maintainer line cytoplasm(s) on expression of resistance to insects in this crop. Of course, the simplest alternative would be to focus on open-pollinated varieties that do not require use of male-sterility for seed multiplication.

**Development of CMS and restorer lines for resistance to insect pests.** Much of the area under high-yielding sorghum cultivars is sown to hybrids in Asia, Australia, and the Americas. Therefore, it is apparent that for host plant resistance to be an important component of pest management in sorghum, we need to transfer the insect resistance genes into male-sterile, maintainer, and restorer lines that can be used by the public institutions and private seed industry to develop insect-resistant hybrids. The CMS, maintainer, and restorer lines with resistance to shoot fly, stem borer, midge, and head bugs



**Figure 1.** Relative susceptibility of different CMS systems to sorghum shoot fly, *Atherigona soccata*.

have been developed at ICRISAT (Table 3). Much of this material has been shared with public institutions and private seed industry over the past decade for use in sorghum improvement, and for developing high-yielding hybrids with resistance to insects. To develop insect-resistant hybrids, the genes conferring resistance to insect pests need to be transferred into both CMS and restorer lines (Fig. 2a, b) (Sharma et al. 2004b). Hybrids based on resistant  $\times$  resistant parents exhibit greater resistance than the hybrids based on other cross combinations (Sharma et al. 1996, Sharma et al. 2004b, Dhillon 2004). The hybrids based on susceptible  $\times$  susceptible parents are highly susceptible. The CMS lines showed a greater influence on the expression of resistance/susceptibility to shoot fly and sorghum midge than the restorer lines. Therefore, efforts should be made to transfer insect resistance genes into a diverse array of both CMS and restorer lines for developing hybrids with resistance to insect pests.

**Wild relatives of sorghum as sources of diverse genes for resistance to insect pests.** Levels of resistance to sorghum shoot fly and stem borer in cultivated germplasm are low to moderate (Sharma et al. 2003). Therefore, it may be important to identify wild relatives of sorghum with high levels of resistance to these pests. Wild relatives of sorghum have been evaluated for resistance to sorghum shoot fly at ICRISAT, and accessions belonging to *Parasorghum* (*S. australiense*, *S. purpureosericeum*, *S. brevicallousum*, *S. timorensis*, *S. versicolor*, *S. matarankense*, and *S. nitidum*) and *Stiposorghum* (*S. angustum*, *S. ecarinatum*, *S. extans*, *S. intrans*, *S. interjectum*, and *S. stipoides*) did not show any shoot fly damage under multi-choice conditions in the field (Venkateswaran 2003) (Table 4). *Heterosorghum* (*S. laxiflorum*) and *Chaetosorghum* (*S. macrospermum*) showed very low damage, while the species belonging to section wild races of *S. bicolor* subsp. *verticilliflorum* (*aethiopicum*, *arundinaceum*, *verticilliflorum*, and *virgatum*) were highly susceptible to shoot fly, as was *S. halepense*. Accessions belonging to *Heterosorghum*, *Parasorghum* and *Stiposorghum* showed little damage by the spotted stem borer under artificial infestation in the field, except for one accession of *Heterosorghum*, which showed 2% deadhearts (Venkateswaran 2003) (Table 5). In contrast, section *Chaetosorghum* (*S. macrospermum*) was highly susceptible to stem borer damage. Within the section *Sorghum*, the four wild races of *S. bicolor* subsp. *verticilliflorum* (races *arundinaceum*, *aethiopicum*, *verticilliflorum*, and *virgatum*) were highly susceptible to stem borer damage. Sorghum midge females did not lay any eggs in the spikelets of *S. angustum*, *S. amplum*, and *S. bulbosum* compared to 30 eggs in spikelets of *S. halepense* under no-choice conditions. Odors from the panicles of *S. halepense* are more attractive to the

females of sorghum midge than the odors from panicles of *S. stipoides*, *S. brachypodum*, *S. angustum*, *S. macropsermum*, *S. nitidum*, *S. laxiflorum*, and *S. amplus* (Sharma and Franzmann 2001). The accessions belonging to the secondary gene pool with diverse mechanisms of resistance to insects can be crossed with cultivated sorghum, while those belonging to the tertiary gene pool may require application of embryo rescue techniques to

transfer resistance genes from the wild relatives into the cultivated sorghums.

**Marker-assisted selection.** It takes five to six generations to transfer a trait within a species into high-yielding locally adapted cultivars through conventional breeding, and one has to evaluate a large number of progenies to be able to select the plants with the appropriate combination

**Table 1. Germplasm accessions identified to be resistant to insect pests in sorghum.**

Insect pest	Germplasm accessions (IS numbers)
Shoot fly	923, 1032, 1034, 1037, 1044, 1054, 1071, 1096, 1104, 1119, 2122, 2123, 2146, 2162, 2168, 2195, 2205, 2265, 2269, 2291, 2309, 2312, 2394, 2681, 3461, 3962, 4224, 4273, 4646, 4663, 4664, 4835, 4881, 4981, 5075, 5076, 5078, 5210, 5429, 5469, 5470, 5480, 5484, 5490, 5511, 5538, 5566, 5571, 5604, 5613, 5619, 5622, 5636, 5648, 8064, 8100, 8320, 8571, 8721, 8811, 8887, 8891, 8918, 8922, 8988, 9009, 9692, 6566, 10711, 10795, 12150, 13674, 14108, 15437, 15896, 16235, 16357, 7726, 17742, 17745, 17747, 17750, 17948, 17966, 18274, 18325, 18366, 18368, 18369, 18371, 18476, 18551, 18580, 18635, 18662, 18700, 18733, 19485, 19569, 19706, 20064, 21871, 21877, 21969, 22039, 22114, 22121, 22144, 22145, 22148, 22149, 22196, 25744, and 26789.
Stem borer	923, 1044, 1051, 1082, 1096, 1104, 2122, 2123, 2146, 2162, 2195, 2263, 2265, 2269, 2290, 2291, 2292, 2312, 2375, 2376, 3962, 4546, 4637, 4646, 4663, 4664, 4756, 4757, 4776, 4835, 4995, 5072, 5210, 5253, 5268, 5469, 5470, 5480, 5484, 5490, 5511, 5566, 5571, 5579, 5585, 5604, 5613, 5619, 5648, 8811, 5658, 6566, 7224, 8165, 8189, 8549, 8671, 12308, 13100, 17745, 17948, 18333, 18371, 18551, 18573, 18577, 18578, 18579, 18581, 18584, 18585, 18662, 18677, 21883, 22039, 22091, 22113, 22114, 22121, 22129, 22144, 22148, 22196, 22778, 23411, 23962, and 24027.
Midge	2290, 2292, 2579, 2687, 2739, 2830, 3461, 6283, 7005, 7134, 7138, 7151, 8100, 8151, 8190, 8196, 8198, 8204, 8577, 8671, 8721, 8729, 8751, 8887, 8891, 8918, 8922, 8988, 10712, 14380, 15107, 8849, 8884, 8946, 9021, 9045, 9107, 9112, 9608, 9807, 18563, 18573, 18695, 18696, 18698, 18733, 19474, 19476, 19512, 19955, 19957, 21006, 21031, 21211, 21871, 21873, 21879, 21881, 21883, 21883-1, 22400, 22464, 22471, 22806, 23748, 26789, 27103, 27466, 31626, 31635, and 31636.
Head bugs	2761, 8064, 14108, 14317, 14334, 14380, 16357, 17610, 17618, 17645, 18579, 19455, 19945, 19948, 19949, 19950, 19951, 19955, 19957, 20024, 20059, 20068, 20638, 20643, 20664, 20740, 21006, 21211, 21443, 21444, 21485, 21525, 21574, 22284, 22507, 23748, 24357, 25069, 25098, 25760, 21512, 25733, 25766, 27329, 27397, 27452, 27466, and 27477.

Sharma et al. (1992, 2003).

**Table 2. Sorghum varieties with resistance to insect pests, developed at ICRISAT.**

Insect pest	Improved lines with resistance to insects (ICSV numbers)
Shoot fly	700, 701, 702, 705, 707, 708, 711 to 714, 717, 726, 89013, 89018, 89025, 93093, and 25001 to 25055.
Stem borer	700, 708, 711, 714, 717, 89008, 89010, 93046, and 25056 to 25162.
Midge	197, 239, 305, 313, 385 to 395, 563, 564, 573, 693, 729 to 758, 804, 830 to 832, 835, 836, 843, 88006, 88013, 88014, 88028, 88032, 88035, 88041, 89001, 89010, 89031, 89034 to 89039, 89042 to 89044, 89049 to 89054, 89057, 89058, 90001 to 90010, 90014, 90016, 90018, 91015, 91025, 92011 to 92013, 92015 to 92018, 92020, 92021, 92023, 92024, 93001 to 93026, 93035, 93046, 93057, 93059, 93065 to 93067, 93069, 95071 to 93093, 95080, 95123 to 95125, 96009 to 96011, 96027, 96031, 96062 to 96082, and 25163 to 25244.
Head bugs	25245 to 25263.

Agarwal BL, Sharma HC, Taneja SL, Reddy BVS, and Stenhouse JH (unpublished).

of traits. The use of DNA markers for indirect selection offers great potential gains for quantitative traits with low heritability, as these are the most difficult characters to work with in the field using direct phenotypic selection. The effectiveness of a marker-assisted selection (MAS) can only be as good as the quality of the phenotypic data on which the development of the marker was based. At ICRISAT, mapping populations have been phenotyped and genotyped for sorghum shoot fly (296B × IS 18551 and BTx 623 × IS 18551), and spotted stem borer, sorghum midge, and aphid (ICSV 745 × PB 15881-3). Genetic linkage maps based on these populations have been constructed to identify quantitative trait loci (QTLs)

associated with resistance to these insects. Polymorphic simple sequence repeat (SSR) loci associated with resistance to shoot fly and the traits associated with resistance to this insect have been identified (Folkertsma et al. 2003). These QTLs are now being transferred into locally adapted hybrid parental lines via SSR based MAS. The QTLs associated with antibiosis and antixenosis mechanisms of resistance to sorghum midge (Tao et al. 2003), and tolerance to green bug (Nagaraj et al. 2005) have also been identified (Table 6). It is hoped that MAS will allow for rapid introgression of the resistance genes, and ultimately gene pyramiding, into the high-yielding varieties and hybrids.

**Table 3. Cytoplasmic maintainer and male-sterile lines with resistance to insect pests, developed at ICRISAT.**

Shoot fly	Stem borer	Midge	Head bugs
ICSB 415, ICSB 416, ICSB 417, ICSB 418, ICSB 419, ICSB 422, ICSB 423, ICSB 425, ICSB 428, ICSB 429, ICSB 432, ICSB 433, ICSB 434, and ICSB 435	ICSB 464, ICSB 467, ICSB 468, ICSB 469, ICSB 472, ICSB 473, and ICSB 474	ICSB 488, ICSB 493, ICSB 494, ICSB 502, ICSB 505, ICSB 508, ICSB 512, ICSB 513, ICSB 516, ICSB 518, ICSB 520, ICSB 524, ICSB 527, and ICSB 541	ICSB 547, ICSB 548, ICSB 550, ICSB 552, ICSB 553, ICSB 555, ICSB 557, and ICSB 563

Reddy BVS, and Sharma HC (unpublished).

**Table 4. Relative susceptibility of wild relatives of sorghum to shoot fly, *Atherigona soccata*.**

Section	Species	Accession	Deadhearts (%)		Adult emergence (%)
			Field conditions	No-choice conditions	
Chaetosorghum	<i>Sorghum macrospermum</i>	TRC 24112	6.7	61.5	-
Heterosorghum	<i>S. laxiflorum</i>	IS 18958	0.0	7.4	6.2
Parasorghum	<i>S. australiense</i>	IS 18954	0.0	10.1	4.2
	<i>S. matarankense</i>	TRC 243576	0.0	0.0	-
	<i>S. purpureosericeum</i>	IS 18943	0.0	50.3	-
	<i>S. nitidum</i>	TRC 243514	0.0	9.7	-
	<i>S. timorensis</i>	TRC 243498	0.0	21.1	-
	<i>S. versicolor</i>	IS 23177	0.0	6.2	-
	<i>S. angustum</i>	TRC 243499	0.0	4.0	0.0
Stiposorghum	<i>S. ecarinatum</i>	TRC 243574	0.0	3.5	-
	<i>S. intrans</i>	TRC 243571	0.0	1.1	-
	<i>S. extans</i>	TRC 243601	0.0	0.0	-
	<i>S. interjectum</i>	TRC 243461	0.0	1.2	-
	<i>S. stipoideum</i>	TRC 243399	0.0	0.0	-
Sorghum	<i>S. aethiopicum</i>	IS 27584	88.9	-	99.5
	<i>S. virgatum</i>	IS 18808	92.2	-	89.0
	<i>S. bicolor</i>	CSH II	96.7	-	-
		S 18551	30.6	70.2	50.8
LSD (P 0.05)			5.8	19.6	18.9

Venkateswaran (2003).


### Development of insect-resistant transgenic sorghums.

Given the wide host range of some of the insect pests, and low levels of resistance in the cultivated germplasm against stem borers, head bugs, and armyworms, it would be highly desirable to combine conventional plant resistance with novel genes from *Bacillus thuringiensis*, protease inhibitors or plant lectins. For plant resistance to be successful in integrated insect pest management, they have to substitute completely or partially for the use of insecticides and/or other methods of pest management, and result in improved economic returns and reduced environmental impact. In addition to the reduction in losses due to insect pests, the deployment of transgenic plants with insecticidal genes will also lead to: i) reduction in insecticide sprays, ii) reduced exposure of farm labour to insecticides, iii) reduction in harmful effects of insecticides on nontarget beneficial organisms, iv) increased activity of natural enemies, v) reduced amounts of pesticide residues in food and food products, and vi) a safer environment to live in. The first transgenic plants were developed in the mid-1980s (Vaeck et al. 1987). Since then, there has been a tremendous progress in development and deployment of transgenic plants for insect resistance (Sharma et al. 2002, James 2003). Toxins from *Bacillus thuringiensis* var *morrisoni* have shown biological activity against the sorghum shoot fly, *A. soccata*. The *B. thuringiensis* toxins Cry1Ac and Cry2A are moderately effective against spotted stem borer, *C. partellus*, while Cry1Ac is effective against *H. armigera* (Sharma et al. 2004a). Sorghum plants having

*cry1Ac* gene have been developed at ICRISAT, and are presently being tested for resistance to spotted stem borer, *C. partellus* (Girijashankar et al. 2005). Combining transgenic resistance to insects with conventional plant resistance will render plant resistance an effective component for pest management in sorghum.

**Genetic engineering of metabolic pathways.** Many secondary plant metabolites such as flavonoids have been implicated in host plant resistance to insects in sorghum. Many compounds of the flavonoid biosynthetic pathway accumulate in response to biotic and abiotic stresses (Heller and Forkman 1993). Genetic engineering offers the opportunity to change the metabolic pathways to increase the amounts of various flavonoids that play an important role in host plant resistance to insect pests. Biotechnology also offers the opportunity to increase the production of secondary metabolites in plants to increase the levels of resistance to insect pests or inhibit the production of toxic metabolites such as HCN in forage sorghum.

**Inducible resistance to insect pests – gene switches.** A wide range of inducible genes have been identified in plants based on endogenous chemical signals such as phytohormones, response to insect attack, or wounding. Chemically induced expression systems or “gene switches” enable the temporal, spatial, and quantitative control of genes introduced into crop plants, or those that are already present in the plants. The best-studied system



**Figure 2.** Insect damage in sorghum hybrids based on resistant (R) and susceptible (S) male-sterile and restorer lines. a: shoot fly, *Atherigona soccata* (FR1 = SPSFR 94006, FR2 = SPSFR 94007, FS1 = ATx623, and FS2 = CK 60A). b: sorghum midge, *Stenodiplosis sorghicola* (MR1 = ICSA 88019, MR2 = ICSA 88020, MS1 = 296A, and MS2 = ICSA 42). FR = Resistant female. FS = Susceptible female. MR = Resistant male. MS = Susceptible male. R = Resistant. S = Susceptible. Sharma et al. (2004b).

utilizes pathogenesis-related protein-1a (*PR 1-a*) gene expression in tobacco (Uknes et al. 1993). *PR 1-a* mRNA levels can also be induced by exogenous application of salicylic acid (Ward et al. 1991). Peptide hormones also induce production of proteinase inhibitors. Systemically induced responses are modified through synthesis and action of jasmonic acid via its lipid precursor, e.g., linoleic acid in tomato. Application of exogenous jasmonate induces

the production of proteinase inhibitors. Enhanced resistance in transgenic rice plants by application of methyl jasmonate and abscisic acid has been reported (Xu et al. 1993).

**Dominant repressible lethal genetic system.** Traditionally, the sterile insect technique has been employed to control several insect pests. However, this system depends on

**Table 5. Relative susceptibility of wild relatives of sorghum to spotted stem borer, *Chilo partellus*.**

Section	Species	Accession	Deadhearts (%)		Larvae recovered (%)
			Field conditions	Greenhouse conditions	
Chaetosorghum	<i>Sorghum macrospermum</i>	TRC 24112	72.9	-	-
Heterosorghum	<i>S. laxiflorum</i>	IS 18958	0.0	82.5	6.0
		TRC 243492	0.0	15.3	0.0
Parasorghum	<i>S. australiense</i>	IS 18955	0.0	10.5	0.0
	<i>S. matarankense</i>	TRC 243576	0.0	5.2	0.0
	<i>S. nitidum</i>	TRC 243514	0.0	0.0	0.0
	<i>S. purpureosericeum</i>	RN 285	0.0	11.1	0.0
	<i>S. timorensis</i>	TRC 243498	0.0	0.0	0.0
	<i>S. versicolor</i>	IS 14262	0.0	0.0	0.0
Stiposorghum	<i>S. angustum</i>	TRC 243499	0.0	0.0	6.0
	<i>S. ecarinatum</i>	TRC 243574	0.0	0.0	0.0
	<i>S. extans</i>	TRC 243601	0.0	0.0	0.0
	<i>S. intrans</i>	TRC 243571	0.0	0.0	0.0
	<i>S. interjectum</i>	TRC 243461	0.0	0.0	0.0
	<i>S. stipoides</i>	TRC 243399	0.0	0.0	0.0
Sorghum	<i>S. aethiopicum</i>	IS 27584	86.7	-	-
	<i>S. virgatum</i>	IS 18808	94.5	98.2	55.0
	<i>S. bicolor</i>	CSH 11	95.5	98.4	90.0
		S 18551	58.0	96.8	40.0
LSD (P 0.05)			10.5	4.4	-

Venkateswaran (2003).

**Table 6. Molecular makers identified to be associated with resistance to insect pests in sorghum.**

Linkage group (LG)	Primers	Linked traits/mechanisms
Sorghum shoot fly, <i>Atherigona soccata</i>		
LG F	<i>Xtxp 258</i> (bp 190/230) <i>Xtxp 289</i> (bp 270/294)	Trichome density
LG G	<i>Xgap 1</i> (bp 180/254) <i>Xtxp 141</i> (bp 154/169)	Deadhearts, leaf glossiness, and trichome density
LG I	<i>IS 328</i> (bp 144/166) <i>IS 264</i> (bp 153/207)	Leaf glossiness
LG J	<i>IS 258</i> (bp 170/193) <i>Xtxp 65</i> (bp 125/134)	Deadhearts and leaf glossiness
Sorghum midge, <i>Stenodiplosis sorghicola</i>		
LG A	<i>RZ 543 ST 698</i>	Antixenosis mechanism of resistance
LG G	<i>ST 1017 SG 14</i>	Antixenosis mechanism of resistance
LG J	<i>TXS 1931 SG 37</i>	Antibiosis mechanism of resistance
Green bug, <i>Schizaphis graminum</i>		
LG 3	<i>Xtxp 12 Xcup 20 Sb1 10</i>	Tolerance mechanism of resistance
LG 5	<i>Xt xp 43 Xtxp 85 Xtxp 335 Xtxp 204</i>	Tolerance mechanism of resistance

Folkertsma et al. (2003), Tao et al. (2003), and Nagaraj et al. (2005).

large-scale production of the target insect, and use of irradiation or chemical sterilization. Release of insects carrying a dominant lethal (RIDL) gene has been proposed as an alternative to the conventional techniques used for insect sterilization (Alphey and Andreasen 2001). This is based on the use of a dominant, repressible, female-specific gene for insect control. A sex-specific promoter or enhancer gene is used to drive the expression of a repressible transcription factor, which in turn controls the production of a toxic gene product. A non-sex specific expression of the repressible transcription factor can also be used to regulate a selectively lethal gene product. Insects produced through genetic transformation using this approach do not require sterilization through irradiation, and could be released in the ecosystem to mate with the wild population to produce sterile insects that will be self-perpetuating.

### Need for future research

- Improvement in precision of screening and selection criteria for resistance to insect pests.
- Gene pyramiding and development of cultivars with multiple resistance to insect pests and diseases.
- Transfer of insect resistance genes into CMS, maintainer, and restorer lines, and exploitation of alternate CMS systems that are less susceptible to insect pests.
- Identification of toxin genes for shoot fly, stem borer, and head bugs, and development of transgenic plants with resistance to insect pests.
- Identification of molecular markers associated with resistance to shoot fly, midge, stem borer, aphids, and head bugs for use in MAS.

**Acknowledgments.** We gratefully acknowledge Dr SL Taneja, Dr BL Agrawal, Dr K Leuschner, Dr KF Nwanze, Dr LR House, Dr SZ Mukuru, Dr JH Stenhouse, Mr CV Abraham, Mr P Vidyasagar, Ms VF Lopez, Mr G Venkateswarulu, Mr MVR Naidu, Dr KV Hariprasad, Mr B Ramaiah, and the collaborating NARS scientists for their contributions in developing insect-resistant varieties.

### References

**Alphey L and Andreasen M.** 2001. Dominant lethality and insect population control. *Molecular and Biochemical Parasitology* 121:173–178.

**Dhillon MK.** 2004. Effects of cytoplasmic male-sterility on expression of resistance to sorghum shoot fly, *Atherigona soccata* (Rondani). Ph.D. Thesis. Hisar, Haryana, India: Department of

Entomology, Chaudhary Charan Singh Haryana Agricultural University, Hisar. 382 pp.

**Folkertsma RT, Sajjanar GM, Reddy BVS, Sharma HC and Hash CT.** 2003. Genetic mapping of QTL associated with sorghum shoot fly (*Atherigona soccata*) resistance in sorghum (*Sorghum bicolor*). Page 42 in Final Abstracts Guide, Plant & Animal Genome XI, Jan 11–15 2003. San Diego, CA, USA: Town & Country Hotel. [http://www.intl-pag.org/11/abstracts/P5d\\_P462\\_XI.html](http://www.intl-pag.org/11/abstracts/P5d_P462_XI.html)

**Girijashankar V, Sharma HC, Sharma KK, Sivarama Prasad L, Royer M, Secundo BS, Lakshmi N and Seetharama N.** 2005. Development of transgenic sorghum for insect resistance against spotted stem borer (*Chilo partellus*). *Transgenic Research* (in press).

**Heller W and Forkman G.** 1993. Biosynthesis of flavonoids. Pages 499–535 in *The Flavonoids, Advances in Research Since 1986* (Harborne JB, ed.). London, United Kingdom: Chapman and Hall.

**James C.** 2003. Preview: Global Status of Commercialized Transgenic Crops: 2003. ISAAA Briefs no. 30. Ithaca, New York, USA: International Service for Acquisition on Agri-Biotech Applications (ISAAA). [http://www.isaaa.org/Publications/briefs\\_26.htm](http://www.isaaa.org/Publications/briefs_26.htm).

**Nagaraj N, Reese JC, Tunistra MR, Smith CM, Amand PT, Kirkham MB, Kofoid KD, Campbell LR and Gerald EW.** 2005. Molecular mapping of sorghum genes expressing tolerance to damage by greenbug (Homoptera: Aphidae). *Journal of Economic Entomology* 98(2):595–602.

**Sharma HC.** 1993. Host plant resistance to insects in sorghum and its role in integrated pest management. *Crop Protection* 12:11–34.

**Sharma HC and Franzmann BA.** 2001. Host plant preference and oviposition responses of the sorghum midge, *Stenodiplosis sorghicola* (Coquillett) (Dipt., *Cecidomyiidae*) towards wild relatives of sorghum. *Journal of Applied Entomology* 125:109–114.

**Sharma HC, Abraham CV, Vidyasagar P and Stenhouse JW.** 1996. Gene action for resistance in sorghum to midge, *Contarinia sorghicola*. *Crop Science* 36:259–265.

**Sharma HC, Ananda Kumar P, Seetharama N, Hariprasad KV and Singh BU.** 2004a. Role of transgenics in pest management in sorghum. Pages 117–130 in *Sorghum Tissue Culture and Transformation* (Seetharama N and Godwin I, eds.). New Delhi, India: Oxford and IBH Publishing Company.

**Sharma HC, Crouch JH, Sharma KK, Seetharama N and Hash CT.** 2002. Applications of biotechnology for crop improvement: prospects and constraints. *Plant Science* 163: 381–395.

**Sharma HC, Dhillon MK, Naresh JS, Ram Singh, Pampapathy G and Reddy BVS.** 2004b. Influence of cytoplasmic male-sterility on the expression of resistance to insects in sorghum. Page 6 in *New Directions for a Diverse*

Planet: Proceedings of the 4<sup>th</sup> International Crop Science Congress, 25 Sept–1 Oct 2004, Brisbane, Queensland, Australia (Fisher T, Turner N, Angus J, McIntyre L, Robertson M, Borrell A and Lloyd D, eds.). Brisbane, Queensland, Australia: <http://www.cropscience.org.au>.

**Sharma HC, Taneja SL, Leuschner K and Nwanze KF.** 1992. Techniques to screen sorghums for resistance to insects. Information Bulletin no. 32. Patancheru 502 324, Andhra Pradesh, India: International Crops Research Institute for the Semi-Arid Tropics. 48 pp.

**Sharma HC, Taneja SL, Kameswara Rao N and Prasada Rao KE.** 2003. Evaluation of sorghum germplasm for resistance to insect pests. Information Bulletin no. 63. Patancheru 502 324, Andhra Pradesh, India: International Crops Research Institute for the Semi-Arid Tropics. 184 pp.

**Singh BU.** 2002. Genetic variability and selection criteria for resistance to spotted stem borer, *Chilo partellus* (Swinhoe) in sorghum [*Sorghum bicolor* (L.) Moench]. Ph.D. Thesis. Hyderabad, Andhra Pradesh, India: Department of Genetics, Osmania University.

**Tao YZ, Hardy A, Drenth J, Henzell RG, Franzmann BA, Jordan DR, Butler DG and McIntyre CL.** 2003. Identifications of two different mechanisms for sorghum midge resistance through QTL mapping. Theoretical and Applied Genetics 107:116–122.

**Uknes S, Dincher S, Friedrich L, Negrotto D, Williams S, Thompson-Taylor H, Potter S, Ward E and Ryals J.** 1993. Regulation of pathogenesis-related Protein-1a gene expression in tobacco. Plant Cell 5:159–169.

**Vaeck M, Reynaerts A, Hofte H, Jansens S, DeBeuckleer M, Dean C, Zabeau M, Van Montagu M and Leemans J.** 1987. Transgenic plants protected from insect attack. Nature 327:33–37.

**Venkateswaran K.** 2003. Diversity analysis and identification of sources of resistance to downy mildew, shoot fly and stem borer in wild sorghums. Ph.D. Thesis. Hyderabad, Andhra Pradesh, India: Department of Genetics, Osmania University.

**Ward ER, Uknes SJ, Williams SC, Dincher SS, Wiederhold DL, Aleander DC, Ahl-Goy P, Mettraux JP and Ryals JA.** 1991. Coordinate gene activity in response to agents that induce systemic acquired resistance. Plant Cell 3:1085–1094.

**Xu D, McElroy D, Thoraburg RW and Wu R.** 1993. Systemic induction of a potato pin 2 promoter by wounding methyl jasmonate and abscisic acid in transgenic rice plants. Plant Molecular Biology 22:573–588.

## **Registration of ICSV 88032: A High Yielding Line Resistant to Sorghum Midge, *Stenodiplosis sorghicola***

**BL Agrawal, HC Sharma\*, CV Abraham and JW Stenhouse** (ICRISAT, Patancheru 502 324, Andhra Pradesh, India)

\*Corresponding author: [h.sharma@cgiar.org](mailto:h.sharma@cgiar.org)

Sorghum [*Sorghum bicolor* (L.) Moench] midge [*Stenodiplosis sorghicola* (Coquillett)] is one of the most important pests of grain sorghum in Asia, Africa, Australia, and the Americas (Harris 1976, Sharma 1993). The larvae of sorghum midge feed on the developing ovary resulting in the production of chaffy spikelets. The damaged panicles present a blasted appearance. Midge damaged spikelets have a pupal case attached to the glumes or have a small exit hole of the midge parasite on the upper glume. Females lay 30–100 eggs singly in the spikelets at flowering during the morning hours, and die by the afternoon. Eggs hatch in 1–4 days. The larvae suck the contents of developing ovaries and complete development in 7–12 days. Larvae pupate inside the glumes, and the pupal period lasts for 3–8 days. Adults live for 2–48 h. Maximum midge abundance occurs during September–October. A small proportion of the larvae enter diapause in the spikelets in each generation, which may last as long as 3 to 4 years.

It is difficult to control sorghum midge with conventional insecticides, as the larvae remain hidden inside the spikelets. Therefore, it is important to develop midge-resistant cultivars to minimize the extent of losses due to this pest (Sharma 1993). Substantial progress has been made in identification and utilization of resistance to sorghum midge (Sharma et al. 1993). The accessions IS 2579C, IS 12666C, TAM 2566, AF 28, DJ 6514, IS 10712, IS 7005, IS 8891, and IS 8721 are diverse sources of resistance to sorghum midge (Sharma 1993), and efforts have been made to transfer resistance into high yielding cultivars in Asia, Australia, and USA. As a result of intensive efforts to breed for resistance to this pest, several midge-resistant varieties with high grain yield have been developed for cultivation by farmers or for use in the national sorghum improvement programs (Johnson et al. 1973, Agrawal et al. 1987, Sharma et al. 1994).

Sorghum line ICSV 88032 is highly resistant to sorghum midge, early, and less susceptible to leaf diseases. It combines high levels of resistance to sorghum midge with yield potential comparable to commercial cultivars. ICSV 88032 (PM 15936-1) is derived by pedigree breeding from a cross between ICSV 197 (midge-resistant line) and ICSV 1 (high-yielding sorghum variety). Its selection number is (ICSV 197 x ICSV 1)-22-1-1-1. The