



Bio-inoculated Nutrient Management Influence on Soil Nutrient Availability Pattern and Growth of Hybrid Pigeonpea (ICPH 2740) under Establishment Methods and Crop Geometry

Bathula Venkatesh¹, M. Malla Reddy¹, Gajanan Sawargaonkar², Ch. Sarada³,
B. Padmaja¹, S. Gopalakrishnan², K. Pavan Chandra Reddy¹, Y.S. Parameswari¹

10.18805/LR-5188

ABSTRACT

Background: A two-year field study was conducted at the ICRISAT research farm during the rainy seasons of 2021 and 2022 to investigate the impact of crop geometry, crop establishment method and sustainable nutrient management practices on nutrient availability pattern and growth of hybrid pigeon pea.

Methods: The experiment followed a split-split plot design. The collected data was analysed using radar graph and heat maps for nutrient availability and dry matter respectively.

Result: Data revealing that transplanted plots registered higher nutrient availability and proportionate root and total dry matter production at various growth stages. Among plant geometry the root and total dry matter production was higher with 100×100 cm. when considering planting methods, transplanting with a square system of 100×100 cm, combined with an integrated nutrient management approach consisting of 150% (or) 100% soil test based NPK, vermicompost at a rate of 5 t ha⁻¹, phosphate-solubilizing bacteria (PSB) and seed treatment with *Rhizobium*, resulted in average of 31.5% higher dry matter production over alone inorganic nutrient management practices. Thus, a square geometry of 100×100 cm, along with sustainable integrated nutrient management (100% soil test based NPK, vermicompost at a rate of 5 t ha⁻¹, PSB and seed treatment with *Rhizobium*), resulted in higher nutrient availability and dry matter production. These findings highlight the importance of careful selection of planting methods, crop geometry and nutrient management practices for maximizing the nutrient mining for production of high dry matter production of hybrid pigeonpea.

Key words: Dry matter production, Hybrid pigeonpea, Phosphorus solubilizing bacteria, *Rhizobium*, Soil nutrient availability.

INTRODUCTION

Now-a-days, farmers face challenges in sowing pigeonpea within the ideal planting window due to irregularities in monsoon patterns (Varatharajan *et al.*, 2019a; Kumar *et al.*, 2021). Moreover, most of the available varieties and hybrids of pigeonpea are photo-sensitive, meaning they enter the reproductive stage when they receive the optimum day length, regardless of the actual sowing time (Varatharajan *et al.*, 2019b). As a result, source development in these plants is constrained (Tigga and Singh, 2019). To address the issue of source-sink imbalances and ensure proper plant development, the concept of transplanting emerged as an alternative to conventional dibbling methods. In this study, the modern transplanting technique was compared with the traditional dibbling method in the main plot treatment. However, it was observed that conventional crop geometry like 150 × 30 cm resulted in significant interplant competition, leading to poor plant expression (Pradeep *et al.*, 2018). Therefore, to avoid these challenges, the plant geometry was considered as a sub-factor in this study.

Higher wider plant geometry especially the hybrid has higher nutrient requirement under clay soils. In order to fulfil demanded nutrient in sustainable manner by integrated approach. Pigeonpea is a deep-rooted crop which can able to extract nutrients from deeper layer of soil at later stages of crop (Singh *et al.*, 2020; 2022a). The initial nutrient

¹Department of Agronomy, College of Agriculture, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad-500 030, Telangana, India.

²International Crops Research Institute for Semi-arid Tropics, Patancheru, Hyderabad-502 324, Telangana, India.

³ICAR-Indian Institute of Oil Seeds Research, Rajendranagar, Hyderabad-500 030, Telangana, India.

Corresponding Author: Bathula Venkatesh, Department of Agronomy, College of Agriculture, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad-500 030, Telangana, India. Email: vihaanvarma179@gmail.com

How to cite this article: Venkatesh, B., Reddy, M.M., Sawargaonkar, G., Sarada, C., Padmaja, B., Gopalakrishnan, S., Reddy, K.P.C. and Parameswari, Y.S. (2023). Bio-inoculated Nutrient Management Influence on Soil Nutrient Availability Pattern and Growth of Hybrid Pigeonpea (ICPH 2740) under Establishment Methods and Crop Geometry. Legume Research. 46(11): 1490-1495. doi: 10.18805/LR-5188.

Submitted: 09-06-2023 **Accepted:** 03-07-2023 **Online:** 27-07-2023

requirement especially nitrogen through biological nitrogen fixation and phosphorus mostly in fixed (or) complex form which may not be available to plant. The unavailable phosphorus makes into available by the activity of phosphorus solubilizing bacteria. With respect to phosphorus at later stages the fixed phosphorus can be

converted to available form by releasing the piscidic acid by roots (Ishikawa *et al.*, 2002). In order to understand the activity of *Rhizobium* and phosphorus solubilizing bacteria (PSB), nutrient availability pattern of soil was examined at various stages of crop phenology. The objective of the present study was to find out the effect of nutrient management practice on nutrient release pattern and proportionate root growth, total dry matter production under planting methods and plant geometry.

MATERIALS AND METHODS

Experimental site

A field study was conducted at the “International Crops Research Institute for Semi-Arid Tropics” (ICRISAT), Patancheru, Hyderabad, during rainy season (*kharif*) of 2021 and 2022. The experimental periods experienced a rainfall of 998.4 mm in 55 rainy days during 2021 and 1000 mm in 53 rainy days during 2022. The experimental plot was characterized by clay soil with a pH of 8.1, an electrical conductivity (EC) of 1.94 dSm⁻¹ and organic carbon content of 0.42%. The soil analysis prior to the commencement of the trial revealed that the available nitrogen (N), phosphorus (P) and potassium (K) were 231.1 kg N ha⁻¹, 29.8 kg P₂O₅ ha⁻¹, 350.5 kg K₂O ha⁻¹, respectively.

Treatment and experimental design

The field study was conducted using a split-split plot design. The main plot treatments consisted of two planting methods of pigeonpea *viz.*, M₁: Dibbling, M₂: Transplanting. The sub-plots treatments were crop geometry *viz.*, S₁: 100 cm × 100 cm (10,000 plants ha⁻¹), S₂: 120 cm × 120 cm (6,944 plants ha⁻¹), S₃: 150 cm × 60 cm (11,111 plants ha⁻¹). The sub-sub plots treatments involve five nutrient management practice *viz.*, N₁: Control, N₂: 100% soil test-based (STB) NPK (25:37.5:8.5 kg ha⁻¹), N₃: 100% STB NPK + vermicompost + Phosphate-solubilizing bacteria (PSB) + Seed treatment (ST) with *Rhizobium*, N₄: 150% STB NPK, N₅: 150% STB NPK + vermicompost + PSB + ST with *Rhizobium*. Dibbling was performed on the same day of transplantation using seedlings that were 23 days and 25 days old during the *kharif* season of 2021 and 2022, respectively. The sowing was done on 20th June and 23th June during *kharif* 2021 and 2022 respectively. The hybrid used in this investigation was ICPH 2740 which was released in 2015 in Telangana on name of mannemkonda kandi.

Data collection

The weight of dry matter is an index of productive capacity of the plant. The plants harvested from each plot boarder rows for estimating dry matter production. The roots were clipped off from each selected plant, transferred to properly labeled cloth bags separately and then partially dried in the shade. Later on, they were subjected to oven drying at 65±2°C until constant weights were recorded and expressed as total dry matter production (kg ha⁻¹) and root dry weight (kg ha⁻¹) at 45, 90, 135 DAS and harvest (Rana *et al.*, 2014).

Available nitrogen (kg ha⁻¹)

Available nitrogen in soil was determined by 2 M KCl extraction with continuous flow analyser at 560 nm method as described by Keeney and Nelson (1983).

Available phosphorus (kg ha⁻¹)

Available phosphorus was extracted from soil by Olsen's reagent (0.5 N NaHCO₃, pH 8.5). The colour development (blue colour) was done by ascorbic acid method given by Watanabe and Olsen (1965) and the intensity of the blue colour was measured at 8800 nm wavelength by using continuous flow analyser.

Statistical analysis

The heat map was generated using GraphPad Prism Version 9.5.1(733) to visualize the root dry matter and total dry matter production at different stages of crop growth under different agronomic practices during the *kharif* season of 2021 and 2022. To ensure proper comparison, the data for root and dry matter production were normalized column-wise. The normalization process involved setting the minimum value as zero per cent and the maximum value as one hundred percent. This allowed for a standardized representation of the data on the heat map.

RESULTS AND DISCUSSION

Soil available nitrogen and phosphorus

For easy and better understanding of soil available nitrogen and phosphorus at different stages of crop depicted in the form of radar graph in Fig 1 and 2, respectively during *kharif* 2021. In this radar graph, the treatments are distributed around the circle, the nutrient availability of pattern at different stages represented with different colour, from centre to periphery there is scale which represents increasing order of nutrient availability from centre to outside of circle. The outer circle which represented with green colour represents the higher and inner circle with red colour denotes lower nutrient availability, respectively.

From this graph, we knew that the soil nitrogen and phosphorus availability was increased from 45 to 90 DAS, but, there onwards gradually increase in trend was shown in all the treatment combinations. Higher soil nutrient availability was recorded in combination of integrated nutrient approach *i.e.* 100% soil test based NPK + vermicompost @ 5 t ha⁻¹ + vermicompost enriched with PSB + seed treatment with *Rhizobium* which was on par with 150% soil test based NPK + vermicompost @ 5 t ha⁻¹ + vermicompost enriched with PSB + seed treatment with *Rhizobium*. Compared to control and inorganic plots the average nutrient (nitrogen and phosphorus) availability was increased 28.1 and 22.2%; 11.6 and 14.8% higher with integrated nutrient management practices during 90 DAS (where highest nutrient availability was registered). With respect to the planting methods at 45 and 90 DAS, higher nutrient availability was recorded with transplanting over dibbling in both the years. Whereas, in plant geometry the

nutrient availability was found non-significant at 45 and 90 DAS. Due to addition of vermicompost and bio-inoculants, the activity of soil micro-fauna was improved so nutrient availability increased by mineralization process (Kumar *et al.*, 2014, 2022), in this way the integrated approach treatments recorded higher nutrient availability compared to the sole inorganic fertilizer application. These results are in tune with Choudhary *et al.* (2013) and Yadav *et al.* (2015).

Dry matter production

The data on root and total dry matter production at 45, 90, 135 DAS and harvest were represented in the form of heat maps (Fig 3, 4) during *kharif* 2021 and 2022, respectively. Analysis of the heat map reveals that the combination of transplanting with wider spacing (100×100 cm plant geometry) in conjunction with the application of 150% soil test based NPK + vermicompost @ 5 t ha⁻¹ + vermicompost enriched with PSB + seed treatment with *Rhizobium*, resulted in higher root dry matter and total dry matter production at all the stages as depicted by the intensive red colour. However, when considering, root dry matter plant⁻¹, it was

observed that transplanting with wider spacing (120×120 cm plant geometry) in combination with 150% soil test based NPK + vermicompost @ 5 t ha⁻¹ + vermicompost enriched with PSB + seed treatment with *Rhizobium*, resulted in higher values compared to the wider square geometry (100×100 cm) as mentioned earlier.

Dry matter production was not proportionately increased under wider plant geometry despite same level of soil nutrient availability. This might be because of higher nutrient availability due to lack of optimum plant population in wider plant geometry (Kumar *et al.*, 2020, 2021). Whereas, with planting methods slow initial growth and lower root development at later stages in case of dibbling over transplanting led to low nutrient uptake and use-efficiency (Rajpoot *et al.*, 2016, 2021).

At harvest, total dry-matter production was 26.1 and 30.8% higher in case of transplanting over dibbling during *kharif* 2021 and 2022, respectively. In case of transplanting, due to vigorous growth during initial stages led to effective uptake and utilization in initial period as a result of nutrient availability. Not only by increased availability, but also required optimum plant population to utilize the available nutrients in case of 100×100 cm




Fig 1: Soil available nitrogen pattern of hybrid pigeonpea under agronomic practices during *kharif* 2021.




Fig 2: Soil available phosphorus pattern of hybrid pigeonpea under agronomic practices during *kharif* 2021.

plant geometry registering 18.6 and 16.9% higher total dry matter production (kg ha^{-1}) over 120×120 cm plant geometry during *kharif* 2021 and 2022 despite the per plant dry matter was higher with later plant geometry. When considering all the three factors, transplanting with (100×100 cm plant geometry) in conjunction with 100% soil test based NPK + vermicompost @ 5 t ha^{-1} + vermicompost enriched with PSB + seed treatment with *Rhizobium* recorded 28.7 and 23.7% higher than sole

100% soil test based NPK. Same treatment combination in comparison to control obtained 56.7 and 57.2% higher during *kharif* 2021 and 2022, respectively at harvest.

Effective utilization of available nutrients is known to increase nutrient-use efficiency, thereby, dry matter production (Rajpoot *et al.*, 2018, 2019). The bio-inoculated integrated nutrient management also plays a crucial role in making nutrients available forms at initial stages of crop




Fig 3: Heat maps showing root and stem dry matter production influenced by agronomic practices during *kharif* 2021.




Fig 4: Heat maps showing root and stem dry matter production influenced by agronomic practices during *kharif* 2022.

growth (Suri *et al.*, 2006; Choudhary *et al.*, 2013; Singh *et al.*, 2022b). Additionally, crop geometry provides an opportunity for better nutrient uptake, thereby, optimum plant expression in terms of growth and development was obtained (Rajpoot *et al.*, 2018, 2021; Gupta *et al.*, 2022). Therefore, the combination of transplanting with square geometry and integrated nutrient management practices can enhance soil nutrient availability and dry matter production by effective use of available nutrients.

CONCLUSION

The hybrid pigeonpea exhibited a highly favourable environment for effective utilisation of available nutrients with maximum root development. This was achieved through the implementation of the transplanting method of establishment along with a square geometry of 100×100 cm. Furthermore, the provision of the required nutrient levels through integrated 100% soil test based NPK + vermicompost 5 t ha⁻¹ + vermicompost enriched with PSB + *Rhizobium* seed treatment, contributed to higher total dry matter production pays way for nutrient use efficiency and higher yield in pigeonpea.

Conflict of interest: None.

REFERENCES

- Choudhary, A.K., Thakur, S.K. and Suri, V.K. (2013). Technology transfer model on integrated nutrient management technology for sustainable crop production in high value cash crops and vegetables in North-Western Himalayas. *Communications in Soil Science and Plant Analysis*. 44(11): 1684-1699.
- Gupta, G., Dhar S., Kumar, A., Choudhary, A.K., Dass, A., Sharma, V.K., Shukla, L., Upadhyay, P.K., Das, A., *et al.* (2022). Microbes-mediated integrated nutrient management for improved rhizo-modulation, pigeonpea productivity and soil bio-fertility in a semi-arid agro-ecology. *Frontiers in Microbiology*. 13: 924407. DOI: 10.3389/fmicb.2022.924407.
- Ishikawa, S., Adu-Gyamfi, J.J., Nakamura, T., Yoshihara, T., Watanabe, T. and Wagatsuma, T. (2002). Genotypic variability in phosphorus solubilizing activity of root exudates by pigeonpea grown in low-nutrient environments. *Food Security in Nutrient-Stressed Environments: Exploiting Plants' Genetic Capabilities*. 111-121.
- Keeney, D.R. and Nelson, D.W. (1983). Nitrogen-inorganic forms. *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties*. 9: 643-698.
- Kumar, R., Sahoo, P.K, Choudhary, A.K. and Mani, I. (2020). Design, development and performance evaluation of tractor-drawn raised-bed pulse-planter for precision sowing of pigeonpea. *Indian Journal of Agricultural Sciences*. 90(9): 1800-1809.
- Kumar, A., Rana, K.S., Choudhary, A.K., Bana, R.S., Sharma, V.K., Prasad, S., Gupta, G., Choudhary, M., Pradhan, A., Rajpoot, S.K., Kumar, A., Kumar, A. and Tyagi, V. (2021). Energy budgeting and carbon footprints of zero-tilled pigeonpea-wheat cropping system under sole or dual crop basis residue mulching and Zn-fertilization in a semi-arid agro-ecology. *Energy*. 231: 120862. <https://doi.org/10.1016/j.energy.2021.120862>.
- Kumar, A., Rana, K.S., Choudhary, A.K., Bana, R.S., Sharma, V.K., Gupta, G., Rajpoot, S.K. *et al.* (2022). Sole- or dual-crop basis residue-mulching and Zn-fertilization lead to improved productivity, rhizo-modulation and soil health in zero-tilled pigeonpea-wheat cropping system. *Journal of Soil Science and Plant Nutrition*. 22(2): 1193-1214. <http://doi.org/10.1007/s42729-021-00723-6>.
- Kumar, A., Suri, V.K. and Choudhary, A.K. (2014). Influence of inorganic phosphorus, VAM fungi and irrigation regimes on crop productivity and phosphorus transformations in okra (*Abelmoschus esculentus* L.)-pea (*Pisum sativum* L.) cropping system in an Acid Alfisol. *Communications in Soil Science and Plant Analysis*. 45(7): 953-967.
- Rajpoot, S.K, Rana, D.S., Choudhary, A.K. (2018). Bt-cotton-vegetable-based intercropping systems as influenced by crop establishment method and planting geometry of Bt-cotton in Indo-gangetic plains region. *Current Science*. 115(3): 516-522.
- Rajpoot, S.K., Rana, D.S., Choudhary, A.K. (2016). Influence of diverse crop management practices on weed suppression, crop and water productivity and nutrient dynamics in Bt-cotton (*Gossypium hirsutum*) based intercropping systems in a semi-arid indo-gangetic plains region. *Indian Journal of Agricultural Sciences*. 86(12): 1637-1641. <https://epubs.icar.org.in/index.php/IJAgS/article/view/65688>.
- Rajpoot, S.K., Rana, D.S., Choudhary, A.K. (2021). Crop and water productivity, energy auditing, carbon footprints and soil health indicators of Bt-cotton transplanting led system intensification. *Journal of Environmental Management*. 300: 113732. <https://doi.org/10.1016/j.jenvman.2021.113732>.
- Rajpoot, S.K., Rana, D.S., Choudhary, A.K., Pande, P. (2019). Cotton establishment methods based system-intensification: Effects on Bt-cotton growth, weed suppression, system crop and water productivity, system-profitability and land-use efficiency in indo-gangetic plains region. *Indian Journal of Agricultural Sciences*. 89(2): 253-260. <https://doi.org/10.56093/ijas.v89i2.87016>.
- Rana, K.S., Choudhary, A.K., Sepat, S., Bana, R.S. and Dass, A. (2014). *Methodological and Analytical Agronomy*. [ISBN: 978-93-83168-07-1]. Post Graduate School, IARI, New Delhi-110 012, India. 276+xii.
- Singh, U., Choudhary, A.K. and Sharma, S. (2020). Comparative performance of conservation agriculture vis-a-vis organic and conventional farming in enhancing plant attributes and rhizospheric bacterial diversity in *Cajanus cajan*: A field study. *European Journal of Soil Biology*. 99: 103197. <https://doi.org/10.1016/j.ejsobi.2020.103197>.
- Singh, U., Choudhary, A.K. and Sharma, S. (2022a). A 3-year field study reveals that agri-management practices drive the dynamics of dominant bacterial taxa in the rhizosphere of (*Cajanus cajan*). *Symbiosis*. 86(2): 215-227. <https://doi.org/10.1007/s13199-022-00834-3>.
- Singh, U., Choudhary, A.K., Varatharajan, T., Sharma, S. (2022b). Agri-management practices affect the abundance of markers of phosphorus cycle in soil: Case study with pigeonpea and soybean. *Journal of Soil Science and Plant Nutrition*. 22: 3012-3020. <https://doi.org/10.1007/s42729-022-00863-3>.

- Suri, V.K., Chander, G., Choudhary, A.K. and Verma, T.S. (2006). Co-inoculation of VA-mycorrhizae (VAM) and phosphate solubilizing bacteria (PSB) in enhancing phosphorus supply to wheat in typic hapludalf. *Crop Research*. 31(3): 357-361.
- Tigga, R. and Singh, S. (2019). Technology transfer through field trials for increasing productivity and profitability of pigeon pea. *Journal of Plant Development Sciences*. 11(2): 97-100.
- Varatharajan, T., Choudhary, A.K., Pooniya, V., Dass, A. and Harish, M.N. (2019a). Integrated crop management practices for enhancing productivity, profitability, production-efficiency and monetary-efficiency of pigeonpea (*Cajanus cajan*) in indo-gangetic plains region. *Indian Journal of Agricultural Sciences*. 89(3): 559-563. <http://epubs.icar.org.in/.../87606>.
- Varatharajan, T., Choudhary, A.K., Pooniya, V., Dass, A., Meena, M.C., Gurung, B., Harish, M.N. (2019b). Influence of integrated crop management practices on yield, photosynthetically active radiation interception, resource-use-efficiency and energetics in pigeonpea (*Cajanus cajan*) in north Indian plains. *Journal of Environmental Biology*. 40(6): 1204-1210.
- Watanabe, F.S. and Olsen, S.R. (1965). Test of an ascorbic acid method for determining phosphorus in water and NaHCO₃ extracts from soil. *Soil Science Society of America Journal*. 29(6): 677-678.
- Yadav, A., Suri, V.K., Kumar, A. Choudhary, A.K. and Meena, A.L. (2015). Enhancing plant water relations, quality and productivity of pea (*Pisum sativum* L.) through AM fungi, inorganic phosphorus and irrigation regimes in a Himalayan acid Alfisol. *Communications in Soil Science and Plant Analysis*. 46(1): 80-93.
- Pradeep, S., Ullasa, M.Y., Kumar Naik, A.H., Ganapathi and Divya, M. (2018). Effect of different organic nutrient management practices on growth, yield and economics of pigeonpea, [*Cajanus cajan* (L.) Millsp.] and soil properties. *International Journal of Farm Sciences*. 7(2): 10-14.