



Genome-wide screening and characterization of *phospholipase A (PLA)*-like genes in sorghum (*Sorghum bicolor* L.)

Vidhi J Sapara^{1,2} · Aishwarya R Shankhapal^{1,3,4} · Palakolanu Sudhakar Reddy¹

Received: 24 January 2024 / Accepted: 13 June 2024

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2024

Abstract

Main conclusion The characterisation of *PLA* genes in the sorghum genome using in-silico methods revealed their essential roles in cellular processes, providing a foundation for further detailed studies.

Abstract *Sorghum bicolor* (L.) Moench is the fifth most cultivated crop worldwide, and it is used in many ways, but it has always gained less popularity due to the yield, pest, and environmental constraints. Improving genetic background and developing better varieties is crucial for better sorghum production in semi-arid tropical regions. This study focuses on the phospholipase A (*PLA*) family within sorghum, comprehensively characterising *PLA* genes and their expression across different tissues. The investigation identified 32 *PLA* genes in the sorghum genome, offering insights into their chromosomal localization, molecular weight, isoelectric point, and subcellular distribution through bioinformatics tools. *PLA*-like family genes are classified into three groups, namely patatin-related phospholipase A (pPLA), phospholipase A1 (*PLA*₁), and phospholipase A2 (*PLA*₂). In-silico chromosome localization studies revealed that these genes are unevenly distributed in the sorghum genome. *Cis*-motif analysis revealed the presence of several developmental, tissue and hormone-specific elements in the promoter regions of the *PLA* genes. Expression studies in different tissues such as leaf, root, seedling, mature seed, immature seed, anther, and pollen showed differential expression patterns. Taken together, genome-wide analysis studies of *PLA* genes provide a better understanding and critical role of this gene family considering the metabolic processes involved in plant growth, defence and stress response.

Keywords Bioinformatics tools · *Cis*-motif · Chromosomal localization · Gene expression studies · In-silico · Metabolic process · Sorghum genome

Communicated by Dorothea Bartels.

Vidhi Sapara and Aishwarya R Shankhapal have contributed equally to this work.

✉ Palakolanu Sudhakar Reddy
Sudhakarreddy.Palakolanu@icrisat.org;
palakolanusreddy@gmail.com

¹ Cell Molecular Biology and Trait Engineering, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru, Hyderabad, Telangana 502324, India

² Department of Genetics, Osmania University, Hyderabad, Telangana, India

³ Present Address: Division of Plant and Crop Sciences, School of Biosciences, University of Nottingham, Nottingham, UK

⁴ Present Address: Plant Sciences for the Bio-Economy, Rothamsted Research, Harpenden, Hertfordshire, UK

Abbreviations

PLA Phospholipase A
pPLA Patatin-related phospholipase A
PLA₁ Phospholipase A1
PLA₂ Phospholipase A2

Introduction

Sorghum, a widely cultivated crop belonging to the grass family Poaceae, serves multiple purposes, particularly as a vital resource for livestock. As a cereal crop and a source of nourishment for humans and domesticated animals and poultry, sorghum holds a crucial position in India's agricultural landscape (Hariprasanna and Rakshit 2016). Ranking as the country's fifth most important cereal crop and third most significant overall, it is a primary food grain for over 750 million people in semi-arid regions (Tsusaka et al. 2015).

Various essential cellular processes, such as carbon partitioning, cell elongation, defense response, seedling establishment, and plant growth, are associated with liberating fatty acids from membranes and storing lipids in plants. The enzymatic activities of phospholipases, a complex and vital class of enzymes, play a significant role in hydrolyzing phospholipids into free fatty acids (FFAs), phosphatidic acid (PA), diacylglycerol (DAG), lysophospholipids, and soluble head groups. Phospholipases also perform numerous physiological functions within plant (Ali et al. 2022).

Phospholipases are crucial in various biological processes, including intracellular signal transmission, phospholipid metabolism, and cell membrane maintenance. These enzymes are divided into three main groups: PLA, which hydrolyze phospholipids to produce lysophospholipids and free fatty acids; phospholipase C (PLC), facilitating hydrolysis of phospholipids to produce diacylglycerol (DAG) and a phosphorylated head group, and phospholipase D (PLD) helps in the hydrolysis of glycerophospholipids into phosphatidic acid (PA) and a free head group, such as choline. Within the PLA group, three subtypes exist pPLA, PLA₁, and PLA₂, which differ in their specific phospholipid cleavage positions. PLA₁ acts on the sn-1 position, PLA₂ on the sn-2 position, and pPLA exhibits activity in both positions (Chen et al. 2013).

The first pPLA subtype in the PLA gene family is the largest class of phospholipases facilitating hydrolysis of phospholipids and other glycerolipids at both the sn-1 and sn-2 positions. It plays a crucial role in lipid hydrolysis, with Ca²⁺ catalytic sites and a patatin domain (Chen et al. 2013). This gene family, essential in cellular processes like cell growth, signal transduction, lipid metabolism, and stress response, is classified into four groups: pPLAI, pPLAII, pPLAIII, and a fourth class including SDP1, SDP1-like, and ATGL-like (Scherer 2002; Eastmond 2006). pPLAs influence signaling molecules, contributing to diverse cellular activities such as defense signals, anther dehiscence, cell elongation, and gravitropism (Chen et al. 2013). The pPLA-I class of phospholipases regulates basal but not pathogen or wound-induced jasmonic acid production (Yang et al. 2007). Notably, *pPLA-IIa* overexpression in Arabidopsis increases plant cell death and enhances resistance to pathogen attacks (Ackermann et al. 1994). Knockout studies of *pPLA-IIc* in Arabidopsis reveal its essential role in the root response to phosphate deficiency (Rietz et al. 2010). Furthermore, *pPLA-III d* overexpression results in a *STURDY* mutant phenotype characterized by a rigid inflorescence stem, thick leaves, short siliques, large seeds, round flowers, and delayed growth (Huang et al. 2001). Pollen-specific pPLAs induce haploidy, and this phenomenon is functionally conserved in monocots, including foxtail millet, maize, and wheat (Gilles et al. 2017; Kelliher et al. 2017; Liu et al. 2017, 2020; Yao et al. 2018; Cheng et al. 2021).

The second, PLA₁ subtype within the PLA gene family is characterized by a highly conserved GX SXG motif and is calcium-independent, with molecular masses ranging from 45 to 50 kDa (Chen et al. 2013). PLA₁ is further classified into three classes, I, II, and III, based on specific sequences at the N-terminal and similarities in the catalytic region. In Arabidopsis, these are distributed among chloroplasts, plastids, cytosols, and mitochondria (Chen et al. 2013). Arabidopsis contains group I, II, and III PLA₁, along with phosphatidic acid-specific PLA₁ (PA-PLA1) and lecithin: cholesterol acyltransferase-like PLA₁ (LCAT-PLA1) (Chen et al. 2011). In rice, there are group I, II PLA₁, and PA-PLA1 (Singh et al. 2012). PLA₁ transcripts are detected in almost all organs of plants, but the expression levels of the individual isoform have been changed according to the tissue specificity (Seo et al. 2008). PLA₁ genes play crucial roles in various biological processes, including the production of jasmonic acid, plant growth and development, senescence, ultraviolet B (UV-B) defense signaling, and shoot gravitropism (Ishiguro et al. 2001; Kato et al. 2002; Lo et al. 2004; Seo et al. 2008, 2011; Hyun et al. 2008; Ellinger et al. 2010).

The third subtype of the PLA gene family, the PLA₂ superfamily in animals, is broadly classified into five main families: secretory PLA₂s (sPLA₂), cytosolic PLA₂s (cPLA₂), Ca²⁺-independent PLA₂s (iPLA₂s), platelet-activating factor acetyl hydrolases (PAF-AHs), and lysosomal PLA₂s (Schaloske and Dennis 2006). Apart from Arabidopsis, PLA₂ have also been characterized in tobacco (Fujikawa et al. 2012), soybean (Mariani et al. 2012) and wheat (Verlotta et al. 2013). In plants, only low molecular weight secretory PLA₂s have been identified, characterized by molecular masses ranging from 13 to 18 kDa and a PA2c domain (Lee et al. 2005). The genomes of rice and Arabidopsis each have three and four secretory PLA₂ paralogs, respectively, linked to responses to both biotic and abiotic stress. These enzymes play crucial roles in various physiological processes, including auxin control, cell elongation, gravitropism, plant growth and development, and cellular signalling (Mariani and Fidelelio 2019; Takáč et al. 2019).

A comprehensive genome-wide analysis of the PLA gene family has been conducted for Arabidopsis (Chen et al. 2011) and rice (Singh et al. 2012) model crops. Comprehensive identification of this important family of genes has been very useful in exploring a broad range of catalytic properties and biological functions. In this study, we have identified the PLA-like gene family in the sorghum genome, which is mainly classified into three main classes: PLA, PLA₁, and PLA₂. Genome-wide analysis of these genes will help establish a solid foundation for characterizing these genes in sorghum. This comprehensive approach will facilitate the thorough analysis of individual functions across different classes.

Materials and methods

Plant material

The sorghum cultivar ICSR 14001 (Parbhani Shakti) was used for expression studies. Different developmental tissues representing the leaf, seedling, root, anther, pollen, immature seed and mature seed were collected from the life cycle of sorghum. The plants for all experiments were grown in glasshouse conditions under natural day-light oscillations, with day/night average temperatures of approximately 28/22 °C and relative humidity of 70/90%.

Identification of *PLA*-like genes in *Sorghum bicolor*

The *PLA*-like protein sequences of *Arabidopsis* and rice were compiled from the Phytozome database (www.phytozome.net/). Every sequence was individually compared with functional annotations by browsing the sorghum genome database (<https://phytozome.jgi.doe.gov/pz/portal.html>) and NCBI BLASTp, resulting in the identification of *PLA* genes in sorghum. The unclassified *PLA* genes were classified into different isoforms by comparing the phylogenetic relationship of their putative protein sequences with clearly classified *PLA* genes from the phytozome server (<http://www.phytozome.net/search.php?show=blast>). The protein sequence length, molecular weight (MW), and theoretical isoelectric point (pI), three significant physicochemical characteristics of the discovered proteins, were examined using the Expert Protein Analysis System (EXPASY) (https://web.expasy.org/compute_pi/) (Gasteiger et al. 2005). The protein subcellular localization was predicted using the online tool WoLF PSORT, available at <https://wolfpsort.hgc.jp/> (Horton et al. 2007).

Multiple sequence alignment and phylogenetic tree analysis of *PLA*-like genes

ClustalW software was used to perform multiple sequence alignment of all *PLA*-like proteins. The neighbor-end joining method (NJ) created the phylogenetic tree with 1000 bootstrap replications using MEGA11 software (Tamura et al. 2021). The tree was modified further by iTOL (<https://itol.embl.de>) (Letunic and Borp 2021). The consensus of protein sequences was visualized by Geneious Prime®2023.2.1 software.

Gene structure, domain and motif identification of *SbPLA* protein

Gene structures of the sorghum *PLAs* were mapped using TBtools software (Chen et al. 2020), and the domains were identified using NCBI CDD (<https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi>) (Wang et al. 2023). The conserved motifs in the sorghum *PLA* genes family were identified using MEME (<http://meme-suite.org/>) (Bailey et al. 2015), with the maximum number of motifs to be discovered set at 5.

Chromosomal localization, syntenic analysis and protein–protein interactions

The position of the putative *PLAs* in sorghum was procured from the phytozome database, and genes were mapped with their relative distance on the chromosome using TBtools software. The Multiple Collinearity Scan toolkit (MCS-canX) in TBtools with default parameters enabled the gene duplication study in *Sorghum bicolor*, *Oryza sativa*, and *Zea mays*. Protein–protein interaction and pathway analysis were done using the online tool STRING (<https://string-db.org/>) (Szklarczyk et al. 2015).

Cis-acting elements analysis of *SbPLA*-like genes

To elucidate the possible *cis*-acting elements that are responsible for gene expression in response to various factors. 1500-bp of genomic DNA sequences upstream from the initiation codon (ATG) of the *SbPLA* genes were extracted and analyzed. The PlantCARE (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>) (Lescot 2002) database was adopted to identify the *cis*-acting elements in the promoter regions.

RNA isolation and quantitative real time-PCR

100 mg of sorghum tissue was used for RNA isolation using Qiagen RNeasy Mini Kit (Germantown, MD, USA). The RNA quality and quantity were analyzed using 1.4% formaldehyde agarose gel electrophoresis and a Qubit RNA BR assay kit (Thermo Fisher Scientific, Waltham, MA, USA). 1500 ng RNA was used for cDNA synthesis using Superscript III (Invitrogen, Carlsbad, CA, USA) and as a template after diluting it with nuclease-free water (1:10). qPCR was carried out in 96-well optical reaction plates, and PCR reaction was performed in a total volume of 10 µL containing 0.5 µM of each primer (1.5 µL), cDNA (1.0 µL), and Sensi Master Mix (2X) and dH₂O were added up to 2.7 µL. The PCR primers were designed using Primer3 and had a GC content of 40–60%, a T_m > 50 °C, a primer length of 20–25 nucleotides, and an expected product size of 90–180 bp. The

qPCR reactions were carried out by following the standard thermal profile: 95 °C for 10 s and then 40 cycles of 15 s at 95 °C, 15 s at 61 °C with fluorescent signal recording, and 15 s at 72 °C. After the 40th cycle, amplicon dissociation curves were measured by heating at 58 to 95 °C with fluorescence measured within 20 min. All qPCR data were obtained from three biological replicates with three technical replicates. Normalized expression was calculated with qBase⁺ software (Schmidt and Delaney 2010) with reference genes *eukaryotic initiation factor4α* (*SbEIF4α*) and *protein phosphatase2A* (*SbPP2A*) (Sudhakar Reddy et al. 2016).

Results

Identification of PLA-like genes in *Sorghum bicolor*

Based on sequence homology, several phospholipase-like genes were identified from the sorghum genome. Consequently, 32 full-length protein-coding PLA-like genes were identified in the sorghum genome. In the sorghum genome, these PLA-like genes, categorized into pPLAs, PLA₁, and sPLA₂ subtypes, are represented by 21, 7, and 4 members. Identified 32 sorghum PLA genes were characterised based on the presence of the number of introns, pI, chromosome localization, number of amino acids, molecular weight, domain localization, cellular localization, and number of transmembrane domains. These identified PLA proteins showed the presence of amino acids ranging from 154 to 1338. Molecular weights ranged from 16.1 to 147.4 kDa, and the pI ranged from 5.55 to 10.07. Subcellular localization prediction results showed that *SbPLA*-like genes are mostly distributed in organelles such as chloroplast, nucleus, and cytoplasm (Table 1).

Multiple sequence alignment and phylogenetic tree analysis of PLA-like genes

Multiple sequence alignments across various sorghum PLA classes have verified the presence of highly conserved regulatory and catalytically crucial motifs. In all PLA₁ members, a notably conserved GX SXG motif was identified. Corresponding to earlier findings, pPLA members exhibited the distinctive esterase box GTSTG and the anion-binding DGGGXRG motif. Similarly, sPLA₂s displayed a well-defined PA2c domain with a highly conserved Ca²⁺ binding loop marked by the YGKYCGxxxxGC motif and the catalytic site LDACCxxHDxCV (Fig. S1).

The study of PLA gene evolution in sorghum, Arabidopsis, and rice identified distinct subgroups in each species. Phylogenetic analysis divided these genes into three subtypes: pPLAs, PLA₁, and sPLA₂, based on their evolutionary relationships between Arabidopsis and rice (Fig. 1).

Additionally, sorghum PLA-like genes were further classified into subgroups according to their functional domains.

Gene structure, domain and motif identification of *SbPLA* protein

Coding and genomic sequences were extracted to analyse the intron–exon structure. The position of introns and exons was analysed in all 32 candidate *SbPLA* genes. All genes showed the presence of none to 8 introns in their structure except *SbpPLA-I* (Fig. 2c). The *SbpPLA-I* gene showed the presence of 18 introns in the gene organization. The position and presence of introns were different in all subtypes. While the length of each exon was similar for most members in each subfamily, some deviations were also noted. Intron exon length varied widely within all 32 *SbPLA* genes. In a nutshell, *SbPLA* showed a complex structural organization with varied lengths of introns and exons.

To study the functions of different *SbPLA*-like proteins, their protein domains were analyzed according to the subtype. The *SbpPLA* subtype contains the patatin domain. The patatin domain was found in all *SbpPLAs* except *SbpPLA-IIIc*, *SbpPLA-III f*, *SbpPLA-III e* and *SbpPLA-III g*. In the *SbsPLA2* subtype, a phospholipase_A2_1 domain was identified in all except *SbsPLA2b*. The *SbPLA1* subtype showed the presence of lipase_3 domain (Fig. 2b). To understand the structural features of *SbPLA* proteins, we also examined the motif of the PLA proteins according to their subtypes. The genes contained in the evolutionary tree subtype *SbpPLA* subdivided into four categories based on the number of motifs like *SbpPLA-I*, *SbpPLA-II*, *SbpPLA-III*, and *SbSDP1* contained one, five, two and none, respectively (Fig. S2a). The second type, the *SbPLA1* class, contained five conserved motifs in most of the sequences, except *SbPLA1-IIa* and *SbPLA1-Ia* which showed four motifs in their protein structure (Fig. S2b). The third subtype, *SbsPLA2*, contained motifs ranging from three to five (Fig. S2c).

Chromosomal localization and syntenic analysis of *SbPLA*-like genes

In silico chromosome localization prediction showed that the *SbPLA* genes were distributed throughout the sorghum genome except for chromosomes 6 and 9 (Fig. 3). All chromosomes carried at least 1 (chromosome 4 and 8) and a maximum of 8 (chromosome 1) of *SbPLA* genes. *SbpPLA* genes are distributed in all chromosomes except chromosomes 4, 6 and 9. The *SbPLA1* gene subtype was found to be distributed in chromosomes 2, 3, 7 and 10. Meanwhile, *SbsPLA2* genes were distributed in chromosomes 1, 4, and 5. The MCscanX results also contributed to detecting syntenic and collinear relationships of sorghum with other crop species. We constructed collinearity syntenic maps of sorghum

Table 1 In-silico characterization of the candidate genes

Annotated name	Gene ID	Introns	Number of amino acid	MW (kDa)	pI	Instability index	Aliphatic index	Gravity	Cellular localization
pPLA									
SbpPLA-I	Sobic.002G320900	32	1338	147.4	5.82	54.93	95.55	- 0.097	Nucleus
SbpPLA-IIa	Sobic.002G228700	5	482	52.4	8.86	31.27	93.79	- 0.095	Cytoplasm
SbpPLA-IIb	Sobic.007G158600	3	435	46.8	6.27	30.26	90.14	- 0.199	Chloroplast
SbpPLA-IIc	Sobic.007G158700	4	487	53.3	5.58	41.73	86.08	- 0.144	Chloroplast
SbpPLA-IId	Sobic.007G158800	8	401	44	6.11	29.73	88.1	- 0.215	Chloroplast
SbpPLA-IIe	Sobic.005G186100	2	410	44.5	6.27	26.75	88.1	- 0.146	Chloroplast
SbpPLA-IIf	Sobic.005G186200	2	405	43.9	6.08	25.11	89.43	- 0.136	Chloroplast
SbpPLA-IIg	Sobic.005G186400	3	413	44.8	5.88	33.1	88.18	- 0.151	Cytoplasm
SbpPLA-IIh	Sobic.003G389400	4	333	45.4	9.34	34.21	77.36	- 0.326	Chloroplast
SbpPLA-IIQ	Sobic.001G348600	3	437	47.4	9.34	42.62	85.79	- 0.136	Chloroplast
SbpPLA-IIk	Sobic.002G377600	4	403	43.7	5.76	32.06	84.07	- 0.218	Cytoplasm
SbpPLA-IIm	Sobic.007G158500	4	437	46.6	6.47	35.35	90.85	- 0.129	Chloroplast
SbpPLA-IIa	Sobic.001G433900	1	460	48.9	6.11	38.78	74.46	- 0.245	Nucleus
SbpPLA-IIb	Sobic.001G067500	1	449	46.6	9.37	38.89	81.92	- 0.080	Chloroplast
SbpPLA-IIc	Sobic.002G031800	2	492	50.3	10.07	51.47	77.97	- 0.112	Cytoplasm
SbpPLA-IIId	Sobic.010G228100	1	471	48.2	9.02	46.32	80.42	- 0.013	Chloroplast
SbpPLA-IIe	Sobic.001G157700	0	461	47	9.06	44.69	80.87	0.036	Chloroplast
SbpPLA-IIIf	Sobic.008G166050	2	484	49.2	9.33	46.55	76.26	- 0.071	Chloroplast
SbpPLA-IIIg	Sobic.001G157900	0	445	45.6	8.76	42.42	78.74	0.015	Chloroplast
SbSDP1	Sobic.003G304200	3	907	100	6.12	53.25	86.96	- 0.306	Nucleus
SbSDP1-L	Sobic.001G041900	7	841	92.9	6.3	52.78	87.24	- 0.230	plastid
PLA1									
SbPLA1-Ia	Sobic.007G036900	0	509	53.9	9.58	54.6	87.31	- 0.12	Chloroplast
SbPLA1-Ib	Sobic.010G047800	0	547	60.5	6.11	42.68	74.9	- 0.4	Chloroplast
SbPLA1-Ic1	Sobic.003G432600	1	526	58.1	6.54	39.35	78.97	- 0.327	Chloroplast
SbPLA1-IIa	Sobic.002G177200	4	277	30	7.06	36.07	101.12	0.16	Nucleus
SbPLA1-IIb	Sobic.003G239500	1	440	46.9	6.27	40.35	82.8	- 0.188	Chloroplast
SbPLA1-IIc	Sobic.003G239200	2	408	45.5	5.98	41.27	79.85	- 0.445	Cytoplasm
SbPLA1-IIId	Sobic.003G274700	0	473	51	5.55	38.64	86.83	- 0.149	Chloroplast
sPLA2s									
SbsPLA2a	Sobic.001G120400	3	155	16.5	5.76	64.79	85.61	0.098	golgi
SbsPLA2b	Sobic.005G145100	3	169	18	5.75	64.57	84.97	- 0.026	ER
SbsPLA2c	Sobic.004G357800	2	157	16.7	8.72	53.86	85.1	0.221	ER
SbsPLA2d	Sobic.001G429900	3	154	16.1	6.03	46.59	90.58	0.144	Extra

The selected candidate genes are characterized by the total number of introns, number of amino acids, molecular weight (MW), isoelectric point (pI), instability index, aliphatic index, gravity, and cellular localization


and the two other crops, *Oryza sativa* and *Zea mays*. The *PLA* gene pairs within the significant collinear blocks (zero e-value) were identified in sorghum-rice (22 pairs) and sorghum-maize (23 pairs) (Fig. 4).

In silico analysis of *SbPLA* promoter regions

To identify putative *cis*-acting elements in the promoter region of *SbPLA*-like genes, genomic sequences approximately 1500 bp upstream from the translational start site were extracted and analysed using the PlantCARE database. The analysis mainly focussed on abiotic stress, phytohormone responsiveness and development-related *cis*-elements, as they are key players in signalling different

stresses and biological processes. The abiotic stress-related *cis*-acting elements identified in promoters of the *PLA* genes included LTR, *cis*-acting element involved in low-temperature responsiveness, *STRE*, stress-responsive elements and *WUN*-motif, wounding-related. The phytohormone-responsive *cis*-acting elements included *P*-box, *TCA*, *CGTCA* motif: *cis*-acting element involved in the *MeJA*-responsiveness, *TGACGA* motif, *ABRE*: *cis*-acting element involved in the abscisic acid responsiveness, *TATC* box and *TGA* element: auxin-responsive element. The development-related *cis*-acting elements included *GCN4* motif, *RY* element, *MBS* (*MYB* binding site involved in drought-inducibility), circadian (*cis*-acting regulatory element involved in circadian control),

Fig. 1 Phylogenetic tree of PLAs from *Sorghum bicolor*, *Arabidopsis thaliana* and *Oryza sativa*. Multiple sequence alignments were performed using amino acid sequences from the PLA class of genes from Arabidopsis, rice and sorghum. Default parameters for ClustalW in Mega 11.0 were used to perform the alignment. The alignment was then used to construct a phylogenetic tree in MEGA 11.0 by the neighbour-joining method with 1000 bootstrap replications. The tree was classified into three different subtypes (pPLAs, sPLA₂ and PLA₁) based on the clades and is indicated by different colour



CCGTCC box and CCGTCC motif. Almost all promoter regions showed the presence of CGTCA motif, TGCGA motif, STRE, and ABRE motif. From this, it can be concluded that all genes mostly contain developmental-specific and hormone-responsive *cis*-acting elements (Fig. 5).

In silico protein–protein interaction analysis

A STRING database was used to analyse protein–protein interactions. This prediction analysis showed that the *SbPLA* genes are involved in various pathways such as linoleic acid metabolism, arachidonic acid metabolism, Ether lipid metabolism, alpha-linolenic acid metabolism, and glycerophospholipid metabolism. The reported functions of this class of candidate genes are lipid degradation, lipid metabolism, and hydrolases. Figure 6 suggests that CM000760.3, CM000766.3, CM000767.3, CM000763.3, and LOC8065835 putative lipoxygenase 5 regulate the genes' major number of the *SbPLA* class. If considering only the *SbPLA* family, *SbSDP1-L* and *SbsPLA2a* are related to a maximum number of proteins in the *PLA* family (Fig. 6).

Expression profile of *SbpPLA* genes in various tissues of sorghum

The expression levels of *SbpPLA* candidate genes in various tissues of sorghum were analyzed by qPCR. Seedling, root, post anther, pollen, mature seed, leaf, and immature seed were considered to examine the expression level of the *SbpPLA* candidate genes. The qPCR results showed that the expression pattern of the *pPLA* gene family is quite diverse. *SbpPLA-I*, *SbpPLA-IIb*, *SbpPLA-IIc*, *SbpPLA-IId*, *SbpPLA-IIe*, *SbpPLA-IIf*, *SbpPLA-IIg*, *SbpPLA-IIh*, and *SbpPLA-IIi* showed low to moderate expression levels in immature seed, leaf, mature seed and pollen whereas downregulation occurred in post anther, root and seedling stages. However, *SbpPLA-IIa* showed a moderate level of expression in pollen and post anther stages, while *SbpPLA-IIQ* and *SbpPLA-IIk* showed a moderate level expression in pollen. *SbpPLA-IIIlb* and *SbpPLA-IIIc* were moderately expressed at the mature seed stage. On the other hand, expression levels of the *SbpPLAIII* class of genes such as *SbpPLA-IIIlb*, *SbpPLA-IIId*, *SbpPLA-IIIf*, *SbpPLA-IIIg* showed an upregulation in roots and seedling stage of the plant (Fig. 7). This




Fig. 2 Phylogenetic relationships, conserved protein domain and gene structure of *PLA* genes in *Sorghum bicolor*. **a** Phylogenetic tree of 32 amino acid sequences *SbPLA* gene family. **b** Protein conserved domain in *PLA*-like genes of sorghum. Blue boxes represent the presence of patatin domain, yellow boxes represent phospholipase_A2_1 domain, pink boxes represent lipase_3 domains while the grey boxes

represents the rest of the protein sequences except the conserved domain in the respective *SbPLA* protein sequences. **c** The exon–intron structure of *SbPLA*-like genes. The purple boxes represent untranslated region (UTR), the orange boxes indicate exons and black lines indicate introns

signifies the role of the *SbPLA* gene family in vital tissue functioning. Overall, these qPCR results depict diverse transcript abundance at different stages of the tissue, proving a solid backbone in regulating growth, defence and stress regulation mechanisms.

Discussion


Proteins of the phospholipase family are categorized into four types: PLAs, PLBs, PLCs, and PLDs, distinguished by their diverse enzymatic functions (Wang 2001). PLAs and PLBs remove hydrophobic fatty acid tails from the glycerol of phospholipids, whereas PLCs and PLDs eliminate hydrophilic phosphate bands or side chains from protein structures.

The exhaustive exploration of the sorghum genome through various methodologies has resulted in the identification of 32 *SbPLA* genes. Similar to Arabidopsis and rice, three distinct groups of PLAs were discerned. Rice exhibited 31 *PLAs* (Singh et al. 2012), while Arabidopsis

had 29 (Chen et al. 2011). According to the evolutionary relation between *SbPLA* with rice and Arabidopsis, *PLA* genes are distributed based on homologous protein similarity and into clusters. Phylogenetic analysis suggests the division of *SbPLA* proteins into three groups, namely pPLA, PLA₁, and sPLA₂, consistent with the classification proposed in Arabidopsis, which showed 13, 12, and 4 members, respectively (Chen et al. 2011), and rice, which exhibited 16, 11, and 3 members, respectively (Singh et al. 2012). Similarly, these proteins are distributed across these classes in sorghum, comprising 21, 7, and 4 members, respectively (Fig. 1).

Results from *in silico* tools for subcellular localization revealed the variable distribution of *SbPLA* genes in chloroplasts, cytoplasm, nucleus, and other organelles. Notably, the *SbPLA* genes followed the localization pattern observed in Arabidopsis (Chen et al. 2011) and rice (Singh et al. 2012). Mostly all the *SbPLA* genes were found in chloroplasts (Table 1). The plant PLA₁ class exhibits a molecular weight range of approximately 45–60 kDa (Matos and Pham-Thi 2009). Similarly, the *SbPLA1* class consists of higher

Fig. 5 The distribution of *cis*-acting elements in the promoter sequences of *SbPLA*-like genes in *Sorghum bicolor*. The upper bar graph uses different colours to recognize various *cis*-acting elements representing the total number of *cis*-acting elements, such as abiotic responsive, phytohormone responsive, and plant development related. The bubble graph is plotted based on the number of *cis*-acting elements in the promoter region. The bubble size is proportional to the abundance of each *cis*-acting element (1.5 kb upstream of the translation start site)



consistent with studies in Arabidopsis, while subtype 2 members contained 1–2 introns in sorghum, similar to rice. However, pPLAs exhibited differences in intron–exon structure compared to Arabidopsis (Scherer 2002) (Fig. 2c). Chromosomal mapping of *SbPLA* genes showed variable distribution on 10 chromosomes, excluding chromosomes 6 and 9. Similarly, in rice, *PLA* genes are distributed on 11 chromosomes except chromosome 4 (Singh et al. 2012) (Fig. 3).

Previous findings have indicated the significance of the *phospholipase* gene family in early developmental stages, as many candidates exhibit developmental-specific elements in their promoter regions. These elements include MBS, circadian, GCN4 motif, RY element, and CCGTCC-box, as well as hormone-responsive elements such as CGTCA

motif, TGCGA motif, STRE, P-box, TCA element, TATC-box, TGA box, and ABRE motif, reflecting diverse functions in plants (Fig. 5). Phospholipases are crucial in developmental and hormone-specific gene functions, contributing to abiotic stress-triggered signalling pathways (Takáč et al. 2019). Moreover, diverse PLA members are present in various plant tissues and overall plant development (Saddhe and Potocky 2023).

The application of STRING analysis on the genes facilitated the construction of a network, offering insights into the potential utility of the candidates. It provided an overview of diverse pathways where the candidates may play a role. The analysis highlighted the predominant involvement of these candidates in lipid degradation, lipid metabolism, and hydrolases. Notably, *SbSDP1-L* and *SbsPLA2a* emerged as key

controllers, participating in various processes and regulating the network of the *PLA* gene family. These proteins belong to the lipolytic acyl hydrolase family, indicating their crucial role in governing membrane function regulation.

The *PLA* gene family is actively involved in various developmental processes of plants, including seed development, root development, and cell elongation (Ali et al. 2022). *PLA₂* plays a role in auxin-induced cell elongation in plants (Scherer 2002). Based on previous studies, the pPLA class is involved in different expression studies in developmental tissue in rice (Singh et al. 2012), Arabidopsis (Matos et al. 2008), and cotton (Wei et al. 2023). This study extends this analysis to examine the pPLA class's expression profiling in various sorghum tissues. Our findings reveal the upregulation of *SbpPLA-III*, particularly in the root and seedling stages, suggesting its potential role in seed development and root elongation processes. In Arabidopsis, certain members of pPLA have been reported to be up-regulated under drought stress conditions, influencing root development via auxin and phosphate deficiency pathways (Matos et al. 2008). Conversely, the expression patterns of *SbpPLA-IIa*, *SbpPLA-IIQ*, and *SbpPLA-IIk* exhibit moderate levels of pollen-specific expression (Fig. 7), hinting at their potential involvement in the fertilization process and suggesting further investigation into their role in haploid development. Phospholipase genes specific to pollen are pivotal in haploid induction across various crops. For instance, the maternal haploid induction system in monocots, based on pPLA, is known as *MATRILINEAL/NOT LIKE DAD/ZmPHOSPHOLIPASE-A1* (Kelliher et al. 2017; Liu et al. 2017; Gilles et al. 2017), and has been successfully implemented in maize (Kelliher et al. 2017), rice (Yao et al. 2018), wheat (Liu et al. 2020), and foxtail millet (Cheng et al. 2021). Recent studies have also demonstrated haploid induction in japonica rice and Arabidopsis through the loss of function in pollen-expressed phospholipase *OsMATL2* (*OspPLA-IIα*) gene and gynoeceium-expressed phospholipase *AII* (*pPLAIIγ*) genes, respectively (Jang et al. 2023a, b).

Conclusion

In this investigation, we successfully identified and characterized 32 *PLA* genes in *Sorghum bicolor*. Utilizing phylogenetic analysis, the 32 *SbPLAs* were classified into three distinct subtypes: pPLA, *PLA₁*, and *sPLA₂*. In silico sequence analysis gained comprehensive insights into the *SbPLA* gene family, encompassing gene structure, conserved motifs, chromosomal localization, promoter elements, and protein–protein interactions. Expression analysis revealed the up-regulation of the pPLA subtype in pollen tissue, while most other genes exhibited a moderate expression level in leaves. Identifying pollen-specific genes in sorghum

suggests their potential role in haploid induction, shortening the breeding cycle process. The phospholipase genes identified in sorghum present promising avenues for future studies exploring signalling pathways related to biotic-abiotic stress, growth hormones, lipid degradation, lipid metabolism, and hydrolases. These investigations pave the way for detailed molecular characterization and an exploration of the in-planta roles of *PLA* genes in sorghum.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00425-024-04467-2>.

Acknowledgements All authors sincerely thank ICRISAT for providing financial assistance and support for conducting research. VS acknowledges the DST-INSPIRE fellowship (DST/INSPIRE Fellowship/IF190079) from the Department of Science and Technology, Government of India for her doctoral research work.

Author contributions Conceptualization: PSR; Methodology: VS, AS, PSR; software: VS and AS; data analysis: VS and AS; writing—original draft preparation, VS and AS; writing—review and editing, PSR; supervision and funding acquisition: PSR. All authors have read and agreed to the published version of the manuscript.

Data availability Data are contained within the article or supplementary material.

Declarations

Conflict of interest The authors declare no conflict of interest.

References

- Ackermann EJ, Kempner ES, Dennis EA (1994) Ca²⁺-independent cytosolic phospholipase A2 from macrophage-like P388D1 cells Isolation and Characterization. *J Biol Chem* 269:9227–9233. [https://doi.org/10.1016/S0021-9258\(17\)37098-9](https://doi.org/10.1016/S0021-9258(17)37098-9)
- Ali U, Lu S, Fadlalla T et al (2022) The functions of phospholipases and their hydrolysis products in plant growth, development and stress responses. *Progr Lipid Res* 86:101158. <https://doi.org/10.1016/j.plipres.2022.101158>
- Bailey TL, Johnson J, Grant CE, Noble WS (2015) The MEME suite. *Nucleic Acids Res* 43:W39–W49. <https://doi.org/10.1093/nar/gkv416>
- Chen G, Snyder CL, Greer MS, Weselake RJ (2011) Biology and biochemistry of plant phospholipases. *Crit Rev Plant Sci* 30:239–258. <https://doi.org/10.1080/07352689.2011.572033>
- Chen G, Greer MS, Weselake RJ (2013) Plant phospholipase A: advances in molecular biology, biochemistry, and cellular function. *Biomol Concepts* 4:527–532. <https://doi.org/10.1515/bmc-2013-0011>
- Chen C, Chen H, Zhang Y et al (2020) TBtools: an integrative toolkit developed for interactive analyses of big biological data. *Mol Plant* 13:1194–1202. <https://doi.org/10.1016/j.molp.2020.06.009>
- Cheng Z, Sun Y, Yang S et al (2021) Establishing in planta haploid inducer line by edited *SiMTL* in foxtail millet (*Setaria italica*). *Plant Biotechnol J* 19:1089–1091. <https://doi.org/10.1111/pbi.13584>
- Eastmond PJ (2006) *SUGAR-DEPENDENT1* encodes a patatin domain triacylglycerol lipase that initiates storage oil breakdown in germinating Arabidopsis seeds. *Plant Cell* 18:665–675. <https://doi.org/10.1105/tpc.105.040543>

- Ellinger D, Stingl N, Kubigsteltig II et al (2010) DONGLE and DEFECTIVE IN ANOTHER DEHISCENCE1 lipases are not essential for wound- and pathogen-induced jasmonate biosynthesis: redundant lipases contribute to jasmonate formation. *Plant Physiol* 153:114–127. <https://doi.org/10.1104/pp.110.155093>
- Fujikawa Y, Fujikawa R, Iijima N, Esaka M (2012) Characterization of secretory phospholipase a 2 with phospholipase a 1 activity in tobacco, *Nicotiana tabacum* (L.). *Lipids* 47:303–312. <https://doi.org/10.1007/s11745-011-3632-3>
- Gasteiger E, Hoogland C, Gattiker A et al (2005) Protein identification and analysis tools on the ExpASY server. In: Walker JM (ed) *The proteomics protocols handbook*. Humana Press, Totowa, pp 571–607
- Gilles LM, Khaled A, Laffaire J et al (2017) Loss of pollen-specific phospholipase NOT LIKE DAD triggers gynogenesis in maize. *EMBO J* 36:707–717. <https://doi.org/10.15252/embj.201796603>
- Hariprasanna K, Rakshit S (2016) Economic importance of sorghum. In: Rakshit S, Wang Y-H (eds) *The sorghum genome*. Springer International Publishing, Cham, pp 1–25
- Holk A, Rietz S, Zahn M et al (2002) Molecular identification of cytosolic, patatin-related phospholipases a from *Arabidopsis* with potential functions in plant signal transduction. *Plant Physiol* 130:90–101. <https://doi.org/10.1104/pp.006288>
- Horton P, Park K-J, Obayashi T et al (2007) WoLF PSORT: protein localization predictor. *Nucleic Acids Res* 35:W585–W587. <https://doi.org/10.1093/nar/gkm259>
- Huang S, Cerny RE, Bhat DS, Brown SM (2001) Cloning of an *Arabidopsis* patatin-like gene, *STURDY*, by activation T-DNA tagging. *Plant Physiol* 125:573–584. <https://doi.org/10.1104/pp.125.2.573>
- Hyun Y, Choi S, Hwang H-J et al (2008) Cooperation and functional diversification of two closely related galactolipase genes for jasmonate biosynthesis. *Dev Cell* 14:183–192. <https://doi.org/10.1016/j.devcel.2007.11.010>
- Ishiguro S, Kawai-Oda A, Ueda J et al (2001) The *DEFECTIVE IN ANOTHER DEHISCENCE1* gene encodes a novel phospholipase a1 catalyzing the initial step of jasmonic acid biosynthesis, which synchronizes pollen maturation, anther dehiscence, and flower opening in *Arabidopsis*. *Plant Cell* 13:2191–2209. <https://doi.org/10.1105/tpc.010192>
- Jang JH, Noh G, Seo HS et al (2023a) Loss of function of pollen-expressed phospholipase *OsMATL2* triggers haploid induction in *japonica* rice. *Plant Physiol* 193:1749–1752. <https://doi.org/10.1093/plphys/kiad422>
- Jang JH, Seo HS, Widiez T, Lee OR (2023b) Loss-of-function of gynoecium-expressed phospholipase *pPLA1y* triggers maternal haploid induction in *Arabidopsis*. *New Phytol* 238:1813–1824. <https://doi.org/10.1111/nph.18898>
- Kato T, Morita MT, Fukaki H et al (2002) SGR2, a phospholipase-like protein, and ZIG/SGR4, a SNARE, are involved in the shoot gravitropism of *Arabidopsis*. *Plant Cell* 14:33–46. <https://doi.org/10.1105/tpc.010215>
- Kelliher T, Starr D, Richbourg L et al (2017) *MATRILINEAL*, a sperm-specific phospholipase, triggers maize haploid induction. *Nature* 542:105–109. <https://doi.org/10.1038/nature20827>
- Lee HY, Bahn SC, Shin JS et al (2005) Multiple forms of secretory phospholipase A2 in plants. *Progr Lipid Res* 44:52–67. <https://doi.org/10.1016/j.plipres.2004.10.002>
- Lescot M (2002) PlantCARE, a database of plant cis-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. *Nucleic Acids Res* 30:325–327. <https://doi.org/10.1093/nar/30.1.325>
- Letunic I, Bork P (2021) Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Res* 49(W1):W293–W296. <https://doi.org/10.1093/nar/gkab301>
- Liu C, Li X, Meng D et al (2017) A 4-bp insertion at *ZmPLA1* encoding a putative phospholipase a generates haploid induction in maize. *Mol Plant* 10:520–522. <https://doi.org/10.1016/j.molp.2017.01.011>
- Liu H, Wang K, Jia Z et al (2020) Efficient induction of haploid plants in wheat by editing of *TaMTL* using an optimized *Agrobacterium*-mediated CRISPR system. *J Exp Bot* 71:1337–1349. <https://doi.org/10.1093/jxb/erz529>
- Lo M, Taylor C, Wang L et al (2004) Characterization of an ultraviolet B-induced lipase in *Arabidopsis*. *Plant Physiol* 135:947–958. <https://doi.org/10.1104/pp.103.036376>
- Mariani ME, Fidelio GD (2019) Secretory phospholipases A2 in plants. *Front Plant Sci* 10:451927
- Mariani ME, Villarreal MA, Cheung F et al (2012) In silico and in vitro characterization of phospholipase A2 isoforms from soybean (*Glycine max*). *Biochimie* 94:2608–2619. <https://doi.org/10.1016/j.biochi.2012.07.021>
- Matos AR, Pham-Thi AT (2009) Lipid deacylating enzymes in plants: old activities, new genes. *Plant Physiol Biochem* 47(6):491–503. <https://doi.org/10.1016/j.plaphy.2009.02.011>
- Matos AR, Gigon A, Laffray D, Pêtres S, Zuily-Fodil Y, Pham-Thi AT et al (2008) Effects of progressive drought stress on the expression of patatin-like lipid acyl hydrolase genes in *Arabidopsis* leaves. *Physiol Plant* 134(1):110–120. <https://doi.org/10.1111/j.1399-3054.2008.01123.x>
- Rietz S, Dermendjiev G, Oppermann E et al (2010) Roles of *Arabidopsis* patatin-related phospholipases a in root development are related to auxin responses and phosphate deficiency. *Mol Plant* 3:524–538. <https://doi.org/10.1093/mp/ssp109>
- Saddhe AA, Potocký M (2023) Comparative phylogenomic and structural analysis of canonical secretory PLA2 and novel PLA2-like family in plants. *Front Plant Sci* 14:1118670. <https://doi.org/10.3389/fpls.2023.1118670>
- Schaloske RH, Dennis EA (2006) The phospholipase A2 superfamily and its group numbering system. *Biochim Biophys Acta (BBA)* 1761:1246–1259. <https://doi.org/10.1016/j.bbalip.2006.07.011>
- Scherer GFE (2002) Secondary messengers and phospholipase A2 in auxin signal transduction. In: Perrot-Rechenmann C, Hagen G (eds) *Auxin molecular biology*. Springer Netherlands, Dordrecht, pp 357–372
- Schmidt GW, Delaney SK (2010) Stable internal reference genes for normalization of real-time RT-PCR in tobacco (*Nicotiana tabacum*) during development and abiotic stress. *Mol Genet Genomics* 283(3):233–41. <https://doi.org/10.1007/s00438-010-0511-1>
- Seo YS, Kim EY, Mang HG, Kim WT (2008) Heterologous expression, and biochemical and cellular characterization of *CaPLA1* encoding a hot pepper phospholipase A1 homolog. *Plant J* 53:895–908. <https://doi.org/10.1111/j.1365-3113X.2007.03380.x>
- Seo YS, Kim EY, Kim WT (2011) The *Arabidopsis* sn-1-specific mitochondrial acylhydrolase AtDLAH is positively correlated with seed viability. *J Exp Bot* 62:5683–5698. <https://doi.org/10.1093/jxb/err250>
- Singh A, Baranwal V, Shankar A et al (2012) Rice phospholipase A superfamily: organization, phylogenetic and expression analysis during abiotic stresses and development. *PLoS ONE* 7:e30947. <https://doi.org/10.1371/journal.pone.0030947>
- Sudhakar Reddy P, Srinivas Reddy D, Sivasakthi K et al (2016) Evaluation of sorghum [*Sorghum bicolor* (L.)] reference genes in various tissues and under abiotic stress conditions for quantitative real-time PCR data normalization. *Front Plant Sci* 7:529. <https://doi.org/10.3389/fpls.2016.00529>
- Szklarczyk D, Franceschini A, Wyder S et al (2015) STRING v10: protein–protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* 43:D447–D452. <https://doi.org/10.1093/nar/gku1003>
- Takáč T, Novák D, Šamaj J (2019) Recent advances in the cellular and developmental biology of phospholipases in plants. *Front Plant Sci* 10:362. <https://doi.org/10.3389/fpls.2019.00362>

- Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular evolutionary genetics analysis version 11. *Mol Biol Evol* 38(7):3022–3027. <https://doi.org/10.1093/molbev/msab120>
- Tsusaka TW, Msere HW, Homann Kee-Tui S et al (2015) Sorghum in semi-arid subsistence agriculture: the case of central Mozambique. *ICRISAT SocioEconomics Discuss Pap Ser* (33):58. <https://doi.org/10.13140/RG.2.1.4061.5920>
- Verlotta A, Liberatore MT, Cattivelli L, Trono D (2013) Secretory phospholipases A2 in durum wheat (*Triticum durum* Desf.): gene expression, enzymatic activity, and relation to drought stress adaptation. *Int J Mol Sci* 14:5146–5169. <https://doi.org/10.3390/ijms14035146>
- Wang X (2001) Plant phospholipases. *Annu Rev Plant Physiol Plant Mol Biol* 52:211–231. <https://doi.org/10.1146/annurev.arplant.52.1.211>
- Wang J, Chitsaz F, Derbyshire MK et al (2023) The conserved domain database in 2023. *Nucleic Acids Res* 51:D384–D388. <https://doi.org/10.1093/nar/gkac1096>
- Wei Y, Chong Z, Lu C, et al (2023) Genome-wide identification and expression analysis of the cotton patatin-related phospholipase a genes and response to stress tolerance. *Planta* 257(3):49. <https://doi.org/10.1007/s00425-023-04081-8>
- Yang W, Devaiah SP, Pan X et al (2007) *AtPLAI* is an acyl hydrolase involved in basal jasmonic acid production and *Arabidopsis* resistance to *Botrytis cinerea*. *J Biol Chem* 282:18116–18128. <https://doi.org/10.1074/jbc.M700405200>
- Yao L, Zhang Y, Liu C et al (2018) *OsMATL* mutation induces haploid seed formation in indica rice. *Nature Plants* 4:530–533. <https://doi.org/10.1038/s41477-018-0193-y>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.