

## Registration of Pearl Millet Parental Lines ICMA 92666 and ICMB 92666 with Multiple Disease Resistance

ICMB 92666 (Reg. no. PL-31, PI 596509) pearl millet [*Pennisetum glaucum* (L.) R. Br.] is the maintainer line of a male-sterile line ICMA 92666 (Reg. no. PL-32, PI 596510). ICMA 92666 and ICMB 92666 were released in 1996 by ICRISAT Asia Center, Patancheru, AP, India, for use as seed parents in hybrid breeding programs. These lines are resistant to ergot (caused by *Claviceps fusiformis* Loveless), smut [caused by *Moesziomyces penicillariae* (Bref.) K. Vánky; syn. *Tolyposporium penicillariae* Bref.], and downy mildew [caused by *Sclerospora graminicola* (Sacc.) J. Schröt.].

The male-sterile line, ICMA 92666, derives its A<sub>1</sub> cytoplasm from ICMA-1 (81A) (1). ICMB 92666 was developed from a BC<sub>1</sub> population [ICMPES 34 × (843B × ICMPES 34)]. 843B (= BKM 2068) is a *d*<sub>2</sub> dwarf, early-maturing, large-seeded maintainer line of male-sterile line 843A (= AKM 2068). 843A and 843B were developed at the Fort Hays Experiment Station, Kansas State University, and introduced into India in 1982 by ICRISAT. 843B is a good general combiner, but is highly susceptible to ergot, smut, and downy mildew. ICMPES 34, developed at ICRISAT Asia Center (IAC) as an ergot resistance source, is a tall, late-maturing line with high levels of resistance to ergot, smut, and downy mildew (2).

During the 1985 cool-dry season, 432 BC<sub>1</sub>F<sub>1</sub> plants selected from among >1000 plants on the basis of early flowering, shorter height, and good tillering, were screened for ergot. Sixty-four of these BC<sub>1</sub>F<sub>1</sub> plants that had <10% ergot and medium to large seed size were selected and evaluated as F<sub>2</sub> progeny rows during the 1986 rainy season. One of the progenies (F<sub>2</sub>-155) flowered in 45 d and had 2% ergot, whereas 843B (control) flowered in 39 d and had 85% ergot. Further pedigree selection within this progeny led to the identification of an F<sub>4</sub> progeny that had <1% ergot (95% in 843B) during the 1987 rainy season. When crossed onto male-sterile line 81A, this F<sub>4</sub> progeny produced a male-sterile hybrid. Eleven additional generations of bulk pedigree selection for agronomic traits in this F<sub>4</sub> progeny and 10 generations of concurrent backcrossing onto male-sterile backcross progenies led to the development of ICMA 92666 and ICMB 92666 during the 1992 warm-dry season. Intermittent evaluation of resistance to downy mildew and ergot during the generations of selfing was done for the F<sub>4</sub>-derived lines. Ergot severity in these lines ranged from 1 to 14%, compared with 75 to 95% in 843B.

Under high disease pressure in four greenhouse seedling inoculation tests and in a disease nursery at Mysore in southern India, ICMB 92666 registered 3 to 13% downy mildew incidence, compared with 58 to 77% in 843B. In an artificial inoculation test at IAC and under natural conditions at two locations with high disease incidence in northern India (Hisar and Gwalior), ICMB 92666 was free of smut, whereas 841A, a commercial male-sterile line of similar height and maturity, had 22 to 41% smut.

In yield trials conducted in four environments, ICMB 92666 had a mean height of 1.4 m, required 51 d from seeding to 50% flowering, and had a mean grain yield of 1.66 t ha<sup>-1</sup>, which was 15% less than ICMB 88004, a commercial male-sterile line of similar maturity and 14 cm shorter height. ICMB 92666 has good tillering, small panicles (17.5 cm) that taper towards the tip, and large seed size (10.5 g 1000 seed<sup>-1</sup>). ICMA 92666 and ICMB 92666 set 70 to 90% seed under open pollination in the main panicle. These lines have creamy anthers and gray-brown seed of globular shape. Male sterility of ICMA 92666 has been stable across seasons and sites.

Seed of ICMA 92666 and ICMB 92666 will be maintained by the Genetic Enhancement Division, ICRISAT Asia Center, and

have been distributed to several hybrid breeding programs in India. Small quantities of seed of these two lines will be provided upon request.

K. N. RAI,\* R. P. THAKUR, AND A. S. RAO (3)

### References and Notes

1. Anand Kumar, D.J. Andrews, R.P. Jain, and S.D. Singh. 1984. ICMA-1 and ICMB-1 pearl millet parental lines with A<sub>1</sub> cytoplasmic-genic male sterility system. *Crop Sci.* 24:832.
2. Thakur, R.P., K.N. Rai, S.B. King, and V.P. Rao. 1993. Identification and utilization of ergot resistance in pearl millet. *Res. Bull.* 17. ICRISAT, Patancheru, India.
3. K.N. Rai and A.S. Rao, Genetic Enhancement Division; R.P. Thakur, Crop Protection Division, Patancheru 502 324, Andhra Pradesh, India. Approved as Journal Article no. 1992 by ICRISAT. Accepted 30 June 1997. \*Corresponding author (k.raai@cgnnet.com) (for 1998, kraai@unlvm.unl.edu).

Published in *Crop Sci.* 38:575 (1998).

## Registration of Smut-Resistant Pearl Millet Parental Lines ICMA 88006 and ICMB 88006

ICMB 88006 (Reg. no. PL-29, PI 596507) pearl millet [*Pennisetum glaucum* (L.) R. Br.] is the maintainer line of ICMA 88006 (Reg. no. PL-30, PI 596508). ICMA 88006 and ICMB 88006 were developed and released in 1996 by ICRISAT Asia Center, Patancheru, AP, India, for use in hybrid breeding programs. ICMA 88006 is resistant to smut [caused by *Moesziomyces penicillariae* (Bref.) K. Vánky; syn. *Tolyposporium penicillariae* Bref.] and downy mildew [caused by *Sclerospora graminicola* (Sacc.) J. Schröt.].

The male-sterile line, ICMA 88006, derives its A<sub>1</sub> cytoplasm from ICMA-1 (81A) (1). ICMB 88006 was developed by crossing 843B (= BKM 2068) with a smut- and downy mildew-resistant line derived from a cross between 81B and a smut resistant source line (SRL 53-1). 843A (= AKM 2068) and 843B were introduced into India in 1982 by ICRISAT from Fort Hays Experimental Station, Kansas State University. SRL 53-1 was developed at ICRISAT Asia Center (IAC) as a source of smut resistance. This line is also resistant to downy mildew. The cross involving 843B was made in the 1982 dry season and the F<sub>1</sub> was planted in the 1982 rainy season. Pedigree selection up to F<sub>4</sub> generation, followed by three generations of bulk pedigree selection for high grain yield, large seed size, earliness, and high levels of resistance to smut and downy mildew produced a maintainer F<sub>7</sub> progeny in 1986. Its male-sterile hybrid on 81A and the F<sub>8</sub> progeny were established as an A-B pair during the 1986 rainy season. Individual plants from the F<sub>8</sub> progeny were further selfed and backcrossed onto individual plants of the sterile hybrid. This process of selfing in the maintainer progeny with concurrent backcrossing of individual plants onto individual plants of the sterile backcross progeny led to a male-sterile BC<sub>4</sub> progeny and the counterpart B-line (F<sub>11</sub> progeny). Two additional backcrosses were made using bulk pollen from the B-line to develop ICMA 88006 and the selfed bulk of the B-line as ICMB 88006 in 1988.

In yield trials conducted in 11 year-location environments in India, ICMA 88006 had 1.62 t ha<sup>-1</sup> mean grain yield, 1.1 m plant height, and 2.2 panicles plant<sup>-1</sup>, which was similar to a male-sterile line 81A (2). ICMA 88006 required 50 d to flowering (4 d earlier than 81A) and had a 1000-seed mass of 12.7 g, 84% higher than that of 81A. Smut severity in smut nurseries at IAC ranged from 0 to 1% in ICMA 88006 and from 17 to 68% in 81A. Downy mildew incidence in downy mildew nurseries and