

QTL and QTL × Environment Effects on Agronomic and Nitrogen Acquisition Traits in Rice

Senapathy Senthilvel^{1†}, Kunnummal Kurungara Vinod^{1‡}, Palaniappan Malarvizhi^{2§}
and Marappa Maheswaran^{3*§}

¹Centre for Plant Breeding and Genetics, Tamil Nadu Agricultural University, Coimbatore 641003, India;

²Department of Soil Science and Agricultural Chemistry, Tamil Nadu Agricultural University, Coimbatore 641003, India;

³Centre for Plant Molecular Biology, Tamil Nadu Agricultural University, Coimbatore 641003, India)

Abstract

Agricultural environments deteriorate due to excess nitrogen application. Breeding for low nitrogen responsive genotypes can reduce soil nitrogen input. Rice genotypes respond variably to soil available nitrogen. The present study attempted quantification of genotype × nitrogen level interaction and mapping of quantitative trait loci (QTLs) associated with nitrogen use efficiency (NUE) and other associated agronomic traits. Twelve parameters were observed across a set of 82 double haploid (DH) lines derived from IR64/Azucena. Three nitrogen regimes namely, native (0 kg/ha; no nitrogen applied), optimum (100 kg/ha) and high (200 kg/ha) replicated thrice were the environments. The parents and DH lines were significantly varying for all traits under different nitrogen regimes. All traits except plant height recorded significant genotype × environment interaction. Individual plant yield was positively correlated with nitrogen use efficiency and nitrogen uptake. Sixteen QTLs were detected by composite interval mapping. Eleven QTLs showed significant QTL × environment interactions. On chromosome 3, seven QTLs were detected associated with nitrogen use, plant yield and associated traits. A QTL region between markers RZ678, RZ574 and RZ284 was associated with nitrogen use and yield. This chromosomal region was enriched with expressed gene sequences of known key nitrogen assimilation genes.

Key words: doubled haploid population; nitrogen acquisition; quantitative trait loci; quantitative trait loci × environment interaction; rice.

Senthilvel S, Vinod KK, Malarvizhi P, Maheswaran M (2008). QTL and QTL × environment effects on agronomic and nitrogen acquisition traits in rice. *J. Integr. Plant Biol.* 50(9), 1108–1117.

Available online at www.jipb.net

Rice (*Oryza sativa* L.) is the staple food crop for more than 50% of the world's population. To feed the fast increasing global population in the coming years, a considerable quantity of rice

will be required (FAO 2002) and this increase in production will have to come from the receding extent of land, soil and water. This warrants the situation in which other production needs to be prudently used. Among the production inputs, nitrogen remains as the most naturally scarce primary input in irrigated rice ecosystems. For over a century now, rice farmers have become increasingly dependent on chemical nitrogenous fertilizers. However, the efficiency of fertilizer nitrogen use by rice is low due to substantial nitrogen losses from flooded soil (De Datta and Buresh 1989). Further, the excessive use of nitrogen fertilizers causes serious adverse ecological consequences (Conway and Pretty 1988). Socolow (1999) cautioned that eutrophication and loss of biodiversity in the ecosystem would be a very serious danger due to the over-application of nitrogen in agriculture.

The efficiency of the rice plant to use soil available nitrogen effectively is the prime factor to its overall yield potential. Efforts to improve this must be guided by a thorough understanding of the genetic processes that govern nitrogen use efficiency (NUE). Significant genotypic differences in NUE have been shown in

Received 10 Nov. 2007 Accepted 27 Mar. 2008

Supported by the International Rice Biotechnology Program Schemes RF 98001 #642 and RF 98001 #700 from the Rockefeller Foundation, New York to M. Maheswaran.

[†]Present address: International Crops Research Institute for the Semi-Arid Tropics, Patancheru 502324, India.

[‡]Present address: Rubber Research Institute of India, Kadaba 574221, Karnataka, India.

[§]Present address: Agricultural College and Research Institute, Tamil Nadu Agricultural University, Killikulam, Vallanadu 628252, Tamil Nadu, India.

*Author for correspondence.

Tel: +91 4630 246 1226;

Fax: +91 4630 246 1268;

E-mail: <mahes@tnau.ac.in>.

rice, which can be better exploited in the rice improvement (Tirol-Padre et al. 1996). The limited understanding on the genetics of this complex trait remains as an impediment to breeders in genetic enhancement of rice for better NUE. Nevertheless, the genotypes are having varying responses to different nitrogen sources and levels, indicating the presence of strong genotype × environment (G × E) interaction.

Quantitative genetic analyses in recent times rely extensively on mapping of loci on molecular linkage maps to be used in marker assisted selection (MAS) programs. Hirel et al. (2001) mapped quantitative trait loci (QTLs) for several agronomic and physiological traits such as nitrate content, nitrate reductase (NR) and glutamine synthetase (GS) in maize. Rauh et al. (2002) mapped specific QTLs to varying nitrogen sources such as nitrate, ammonium, and ammonium nitrate in *Arabidopsis*. Similarly, Loudet et al. (2003) mapped 48 QTLs for traits such as shoot growth, total nitrogen, nitrate and free amino acid contents in *Arabidopsis*. In rice, Obara et al. (2004) found that the QTL region of chromosome 2 was linked to panicle number and panicle weight contained a regulatory gene (*GS1*) for glutamine synthetase activity, and selections based on this QTL region proved to be superior in tillering ability, panicle number and total panicle weight, particularly under low nitrogen. Lian et al. (2005) mapped QTLs for low nitrogen tolerance at the seedling stage involving a recombinant inbred lines (RIL) population in rice.

Based on the success of the above works, we attempted to locate the QTLs associated with nine traits of agronomic importance presumably related to nitrogen nutrition and three parameters that are directly associated with nitrogen uptake and nitrogen use efficiency in dihaploid (DH) lines of IR 64 × Azucena grown under three nitrogen regimes.

Results

Phenotypic variation and heritability

The mean performance of parents and DH lines under three nitrogen regimes is given in Table 1. IR64 recorded higher grain yield per plant in g (SPY) together with higher values of number of tillers (TLN), number of productive tillers (PTN) and spikelet fertility percentage (SFP) than Azucena under all three nitrogen levels. IR64 was dwarfed and early flowering compared to Azucena. It recorded higher values for percentage nitrogen content (NCP) in plant and nitrogen uptake in g (NUP). IR64 was more nitrogen responsive compared with Azucena. However, Azucena recorded higher NUE than IR64 at the native N level. The limitation of nitrogen availability had a systematic decreasing effect on all of the agronomic traits substantially in IR64 except for spikelets per primary panicle (SPP) and SFP. The mean values of DH lines were in between the parental values for all of the traits. However, a large number of trans-

gressive segregants were observed for every trait. The broad sense heritability (h^2) values for all agronomic traits except plant height in cm (PHT) recorded moderate to low values (Table 2). All of the nitrogen parameters recorded a very low heritability. For NCP, the h^2 value ranged between 27% and 42% across three nitrogen levels, whereas it was 12%–56% for NUP and 49%–53% for NUE.

Correlation analysis

The correlation analysis carried out across the twelve parameters, indicated that the traits NUP and NUE were positively associated with SPP, filled grains per panicle (FGP), SFP and SPY (Table 3). NCP showed positive correlations with NUP; while it had significant negative associations with days-to-flowering (DTF), PHT and panicle length in cm (PLG). NUP had positive association with PHT, TLN and PTN. Among agronomic traits, SPY showed significant positive correlation with PTN, SPP, FGP and SFP.

Genotype-by-environment interaction

The stratification of variance among genotypes, environments and G × E sources of variation varied appreciably from trait to trait (Table 2). The genotypic effect was large and significant for all traits except for the PTN and NUP, for which the environment effect was predominant. The environmental effect was significant for TLN and SPY also. For all other traits, the proportion of environment effect was negligible. Highly significant ($P < 0.01$) G × E effects were noticed for DTF, SPP, FGP, SFP, SPY, NCP, NUP and NUE. The proportion of G × E effect was the highest in NCP followed by genotype and environment effects. The G × E effect was significant ($P < 0.05$) for TLN, PTN and PLG. PHT was the only trait having non-significant G × E effect.

Principal component analyses

Principal component analyses carried out independently for three nitrogen regimes, extracted four components cumulatively describing total variance of 75.20% (native nitrogen) to 81.09% (high nitrogen). In all cases, principal component 1 (PC1) and principal component 2 (PC2) together described more than 50% of the total variance. Component weights of individual factors, varied considerably between the nitrogen regimes, even shifting the direction of influence across regimes. For PC1, TLN and PTN under native levels showed high negative component weights, which shifted to low positive values in the optimum level and reverted back to negative values under high nitrogen regimes. Similar shifting of vector directions can be seen with NCP and NUP. NUE had positive component weights under native and normal nitrogen regimes and negative influence under high nitrogen (Table 4). The factor orientations also show

Table 1. Phenotypic mean values of parents and range for double haploid (DH) lines for various agronomic traits and nitrogen uptake parameters under different levels of nitrogen

Trait	Mean						Range		
	IR64			Azucena			DH lines		
	Native	Normal	High	Native	Normal	High	Native	Normal	High
Days-to-flowering (DTF)	105.33	118.33	96.33	118.33	128.00	126.00	88.70–129.70	89.00–126.00	95.00–134.70
Plant height (PHT) (cm)	58.67	61.00	62.67	99.83	105.67	107.00	45.70–127.70	46.30–124.70	47.00–131.30
Number of tillers (TLN)	6.33	7.00	10.67	2.00	3.00	4.00	2.00–9.30	3.30–9.30	3.30–13.30
Number of productive tillers (PTN)	6.33	7.00	10.67	1.67	2.67	4.00	2.00–7.00	2.70–9.30	3.30–12.70
Panicle length (PLG) (cm)	18.70	20.50	21.00	23.30	26.00	26.17	15.30–31.80	16.50–30.70	16.00–30.50
Spikelets per primary panicle (SPP)	80.44	173.40	149.84	175.76	186.87	143.69	22.00–185.00	20.30–188.31	18.12–198.00
Filled grains per panicle (FGP)	61.00	131.00	125.00	67.67	117.00	88.67	10.30–113.70	9.70–118.30	7.70–149.70
Spikelet fertility percentage (SFP)	75.83	73.55	83.42	38.50	62.61	61.71	8.00–83.20	7.10–78.00	8.30–90.30
Grain yield per plant (SPY) (g)	7.52	10.30	15.08	5.93	6.31	3.47	0.59–7.19	0.43–15.77	0.57–15.64
Percent nitrogen content (NCP)	1.14	1.09	1.16	1.07	0.82	0.80	0.58–1.25	0.67–1.25	0.63–1.19
Nitrogen uptake (NUP) (g)	0.17	0.19	0.34	0.14	0.16	0.18	0.05–0.21	0.08–0.30	0.12–0.38
Nitrogen use efficiency (NUE)	39.69	55.32	44.23	45.66	36.32	21.06	4.94–93.62	2.89–73.67	2.68–73.91

Table 2. Heritability and proportion of variances due to genotype (G), environment (E) and genotype × environment (G × E) for various agronomic traits and nitrogen uptake parameters under different levels of nitrogen

Traits	Heritability (h^2)			Variance		
	Native	Normal	High	Genotype (G)	Environment (E)	G × E
Days-to-flowering (DTF)	0.65	0.74	0.57	76.32**	0.00	23.68**
Plant height (PHT) (cm)	0.85	0.92	0.87	93.30**	1.22	5.48
Number of tillers (TLN)	0.24	0.32	0.39	44.10**	36.24**	19.66*
Number of productive tillers (PTN)	0.23	0.30	0.42	38.82**	45.19*	15.99*
Panicle length (PLG) (cm)	0.62	0.66	0.63	83.27**	4.30	12.43*
Spikelets per primary panicle (SPP)	0.34	0.28	0.43	63.84**	4.10	32.06**
Filled grains per panicle (FGP)	0.48	0.45	0.51	70.01**	2.73	27.26**
Spikelet fertility percentage (SFP)	0.53	0.48	0.60	75.83**	0.15	24.02**
Grain yield per plant (SPY) (g)	0.36	0.49	0.61	54.43**	20.46**	25.11**
Percent nitrogen content (NCP)	0.42	0.27	0.27	41.29**	3.44	55.25**
Nitrogen uptake (NUP) (g)	0.56	0.12	0.29	25.09**	47.31**	27.60**
Nitrogen use efficiency (NUE)	0.49	0.53	0.53	66.13**	1.00	32.87**

* $P = 0.05$; ** $P = 0.01$.

that the position of IR64 was determined greatly by the direction encompassed by SPY, SFP, SPP, NUE and NUP, while position of Azucena was bracketed by PHT and PLG except in native nitrogen regime (Figure 1).

Main-effect QTLs

Quantitative trait loci analysis revealed 16 main effect QTLs associated with the traits except NUP, distributed on chromosomes 1, 3, 5, 7, 8, 9 and 11 (Table 5). There were three QTLs identified each for PHT and PTN, two for NCP, and one each for DTF, TLN, PLG, FGP, SPP, SFP, SPY and NUE. The likelihood

of odds (LOD) values for identified QTLs ranged from 2.37 to 12.43.

Genetic effects of QTL and QTL-by-environment interaction

The QTL identified for PLG, FGP and SPP had non-significant additive effects within individual environments. The average additive genetic effect of putative QTL for DTF ($qDTF-3$) was 3.15 explaining 16% of the total phenotypic variation (Table 5). Of the three QTLs identified for PHT, $qPHT-1$ and $qPHT-3$ were contributed by IR64, while $qPHT-8$ came from Azucena. These

Table 3. Pearson's correlation coefficients showing interrelations of agronomic traits and nitrogen uptake parameters across different levels of nitrogen

Trait	DTF	PHT	TLN	PTN	PLG	SPP	FGP	SFP	SPY	NCP	NUP
PHT	0.316**										
TLN	-0.211	-0.498**									
PTN	-0.218*	-0.476**	0.965**								
PLG	0.146	0.762**	-0.474**	-0.438**							
SPP	0.019	0.094	-0.076	0.004	0.206						
FGP	-0.102	0.202	-0.319**	-0.272*	0.374**	0.664**					
SFP	0.110	0.007	0.132	0.208	0.004	0.767**	0.062				
SPY	0.062	0.030	0.192	0.297**	0.112	0.822**	0.334**	0.826**			
NCP	-0.336**	-0.452**	0.211	0.221*	-0.273*	0.167	0.054	0.159	0.205		
NUP	0.016	0.251*	0.242*	0.274*	0.178	0.427**	0.253*	0.348**	0.541**	0.340**	
NUE	0.041	-0.072	0.050	0.151	0.081	0.718**	0.296**	0.744**	0.851**	0.040	0.099

* $P=0.05$; ** $P=0.01$. DTF, days-to-flowering; FGP, filledgrains per panicle; NCP, percent nitrogen content; NUE, nitrogen use efficiency; NUP, nitrogen uptake (g); PHT, plant height (cm); PLG, panicle length (cm); PTN, number of productive tillers; SFP, spikelet fertility percentage; SPP, spikelets per primary panicle; SPY, grain yield per plant (g); TLN, number of tillers.

Table 4. Component weights of the extracted four principal components (PC1 to PC4) for twelve traits under three regimes of nitrogen

Trait	Native N				Normal N				High N			
	PC1	PC2	PC3	PC4	PC1	PC2	PC3	PC4	PC1	PC2	PC3	PC4
DTF	0.164	0.001	0.357	-0.405	0.095	-0.217	-0.192	0.022	0.026	0.103	0.455	0.351
PHT	0.378	0.118	0.477	0.061	0.126	-0.430	0.238	0.303	0.098	0.446	0.124	0.419
TLN	-0.449	-0.179	0.154	0.060	0.021	0.491	0.201	0.315	-0.293	-0.386	0.047	0.331
PTN	-0.424	-0.238	0.131	-0.036	0.079	0.486	0.230	0.293	-0.313	-0.382	0.050	0.282
PLG	0.332	0.023	0.446	0.289	0.177	-0.402	0.248	0.146	0.112	0.463	0.044	0.280
SPP	0.241	-0.429	-0.086	0.159	0.461	-0.059	-0.066	-0.177	-0.370	0.322	-0.016	-0.178
FGP	0.130	-0.181	-0.186	0.716	0.283	-0.233	0.219	0.033	-0.148	0.364	-0.267	-0.240
SFP	0.178	-0.358	-0.020	-0.385	0.425	0.138	-0.238	-0.211	-0.404	0.088	0.200	-0.042
SPY	0.160	-0.505	0.027	-0.115	0.490	0.134	-0.030	0.065	-0.479	0.112	0.072	-0.019
NCP	-0.284	-0.249	0.076	0.047	0.026	0.131	0.296	-0.773	-0.126	-0.018	-0.647	0.046
NUP	-0.211	-0.325	0.509	0.149	0.253	0.066	0.628	-0.022	-0.275	0.127	-0.382	0.497
NUE	0.284	-0.359	-0.313	-0.142	0.399	0.115	-0.400	0.149	-0.387	0.094	0.300	-0.305
Eigenvalue	3.335	2.982	1.513	1.195	3.456	3.026	1.572	1.209	3.825	2.804	1.777	1.325
Cumulative variance (%)	27.79	52.64	65.25	75.20	28.80	54.01	67.11	77.18	31.88	55.25	70.05	81.09

DTF, days-to-flowering; FGP, filledgrains per panicle; NCP, percent nitrogen content; NUE, nitrogen use efficiency; NUP, nitrogen uptake (g); PHT, plant height (cm); PLG, panicle length (cm); PTN, number of productive tillers; SFP, spikelet fertility percentage; SPP, spikelets per primary panicle; SPY, grain yield per plant (g); TLN, number of tillers.

QTLs together explained 13.1% of the total phenotypic variation. There were three QTLs associated with PTN, of which two were located on chromosome 3 (*qPTN-3-1* and *qPTN-3-2*) and one on chromosome 9 (*qPTN-9*). Together, they accounted for 13.7% of the total phenotypic variation and were contributed by Azucena. For FGP, a QTL was located on chromosome 7, flanked by RG477 and PGMS0.7 (*qFGP-7*) having a contribution of 3.2%. The QTLs for TLN (*qTLN-1*) and SPP (*qSPP-1*) were found located on the same marker interval of chromosome 1 flanked by RG246 and K5. *qTLN-1* had 12.8% contribution with

an average additive effect of -0.444, and had allelic influence from Azucena, while *qSPP-1* had an average additive effect of 2.81, with allelic favor of IR64. Two QTLs, *qPLG-11* and *qSFP-11* shared the same marker interval (RG247-RG103) on chromosome 11, with the QTL peaks located at 100 cM and 80 cM, respectively. The QTLs for SPY, NCP and NUE, *qSPY-3*, *qNCP-3-1* and *qNUE-3*, shared contiguous marker intervals on chromosome 3, involving markers, RG191, RZ678, RZ574 and RZ 284, all having considerable influence on their respective traits. Similarly the QTLs, *qDTF3*, *qPTN-3-2* and

Table 5. Effect of detected main effect quantitative trait loci (QTL) and QTL × environment interaction for three N regimes by composite interval mapping

Trait	Chr.	Marker interval	QTL	Position of QTL peak (cM)	LOD	R^2	QTL additive effect (a)				MS (QTL × E)	General variance explained
							Native	Normal	High	Average		
DTF	3	RZ448 – RZ519	<i>qDTF-3</i>	300	2.37	15.70	2.82*	4.06**	2.58*	3.15**	44.96*	16.60
PHT	1	RG690–RZ730	<i>qPHT-1</i>	198	12.43	50.30	5.06**	6.26**	6.18**	5.84**	14.69*	13.10
	5	RZ390–RG313	<i>qPHT-5</i>	16	4.64	24.00		4.38*			8.05*	
	8	RZ66–AC5	<i>qPHT-8</i>	142	5.76	27.70		–4.32*	–4.67*	–4.07*	26.19*	
TLN	1	RG246–K5	<i>qTLN-1</i>	28	4.65	24.30	–0.32*	–0.44**	–0.58**	–0.44**	1.54*	12.80
PTN	3	RG348–RZ329	<i>qPTN-3-1</i>	28	4.37	22.00						13.70
	3	RZ519–Pgi1	<i>qPTN-3-2</i>	320	4.57	22.70						
	9	RZ12–RG667	<i>qPTN-9</i>	112	9.06	40.20	–0.28*	–0.41*	–0.50*	–0.40**		
PLG	11	RG247–RG103	<i>qPLG-11</i>	100	4.61	22.80					1.71*	1.50
FGP	7	RG477–PGMS0.7	<i>qFGP-7</i>	80	3.99	20.30					169.17*	3.200
SPP	1	RG246–K5	<i>qSPP-1</i>	46	4.19	22.20					809.92*	0.000
SFP	11	RG247–RG103	<i>qSFP-11</i>	80	4.71	23.30	–9.82**		–7.47*	–7.58**	110.75*	8.50
SPY	3	RZ678–RZ574	<i>qSPY-3</i>	122	5.05	24.70	0.43*	0.82*	0.95*	0.73*	4.79*	10.30
NCP	3	RG191–RZ678	<i>qNCP-3-1</i>	58	6.12	29.100		–0.05*	–0.03*	–0.03*		48.20
	3	Pgi1–CDO87	<i>qNCP-3-2</i>	372	4.67	23.600	–0.04*			–0.03**		
NUE	3	RZ574 – RZ284	<i>qNUE-3</i>	124	5.46	26.40		5.66*	3.64*	4.11**	19.12*	11.00

* $P = 0.05$; ** $P = 0.01$. DTF, days-to-flowering; FGP, filled grains per panicle; LOD, likelihood of odds; NCP, percent nitrogen content; NUE, nitrogen use efficiency; NUP, nitrogen uptake (g); PHT, plant height (cm); PLG, panicle length (cm); PTN, number of productive tillers; SFP, spikelet fertility percentage; SPP, spikelets per primary panicle; SPY, grain yield per plant (g); TLN, number of tillers.

qNCP-3-2 shared contiguous marker intervals on chromosome 3 bracketed by markers RZ448 and CDO 87.

Quantitative trait loci-by-nitrogen level interaction was generally low among the main effect QTLs identified barring a QTL for SPP. High level of interaction was shown by this QTL (*qSPP-1*) having high MS(QTL × E) value (809.92) and very low independent contribution toward the general variance. The additive effects of this QTL were insignificant among the nitrogen regimes. There were also differences among the additive effects of the QTLs under different nitrogen regimes. The QTLs for PTN and NCP did not show significant QTL × environment interaction.

Discussion

Breeding crop plants for better NUE is one of the primary objectives in modern agriculture, considering the importance of imparting low nitrogen tolerance to mitigate the adverse effects of nitrogen saturation and also to decrease fertilizer usage, which bloats the cost of cultivation. This depends on the identification of heritable physiological or biochemical traits, causally related to grain yield. In the present study, a total of nine agronomic parameters associated with nitrogen nutrition and three parameters quantifying NUE were studied. The genetic

loci associated with each of the parameters were mapped on a molecular linkage map of IR64/Azucena.

Considerable levels of genotype and genotype × environment variation were observed among the DH lines and parents for all of the traits studied under different nitrogen regimes. Responses of genotypes to the soil as well as applied nitrogen not only depend on the applied form (Rauh et al. 2002) but also on the levels (Gallais and Hirel 2004). The SPP and FGP of IR64 observed in the native regime are smaller than that of normal and high nitrogen regimes. This had possibly occurred due to low nitrogen susceptibility of the IR64 at low (native) levels, which might have considerably reduced the number of filled grains in panicles other than primary. Primary panicle had more filled spikelets because the development of which might have used more of mobilizing nitrogen within the plant system. In Azucena, FGP was lesser in low N conditions, but it produced more SPP, probably as an adaptive mechanism for low nitrogen. The mean performance of DH lines indicated that there were not many differences in DTF, PHT and PLG across the nitrogen levels, whereas there was an increasing trend in TLN, PTN, FGP, SPY and NUP with increasing nitrogen level. The genetic control of these traits is governed by quantitative genes, which may be dominant, additive and/or epistatic. They may have an array of suppressors and modifiers that are under disparate regulatory influences and modify the cumulative expression of the ultimate trait. They are thus differentially expressed under




Figure 2. Biplots showing scatter of double haploid (DH) lines and components under three nitrogen regimes by principal component analyses.

The positions of both the parents IR 64 and Azucena are shown. DTF, days-to-flowering; FGP, filledgrains per panicle; NCP, percent nitrogen content; NUE, nitrogen use efficiency; NUP, nitrogen uptake (g); PHT, plant height (cm); PLG, panicle length (cm); PTN, number of productive tillers; SFP, spikelet fertility percentage; SPP, spikelets per primary panicle; SPY, grain yield per plant (g); TLN, number of tillers.

form. On the other hand, a limited number of plants are capable of absorbing nitrogen in the ammoniacal form and rice is one among them (Wang et al. 1993; Kronzucker et al. 1999). However, absorption of nitrate and ammonia by roots depends on many transport systems, namely, high affinity transport systems (HATS) and low affinity transport systems (LATS). Members of nitrate transporters and ammonium transporters belonging to both these systems are reported (Glass et al. 2002). Among these genes, NRT 1 and NRT 2 families are very crucial in nitrate uptake in plants (Crawford and Glass 1998). Furthermore, the nitrate pathway is primarily regulated by enzymes such as nitrate reductase (NR), nitrite reductase (NiR), glutamine synthetase (GS) and glutamate synthase (GOGAT) (Meyer and Stitt 2001). In addition, many genes encoding for enzymes like glutamate dehydrogenase, asparagine synthase, urease may also be involved (Gerendas et al. 1998; Gievarec et al. 2004; Lillo 2004). Microarray analyses had revealed, as many as 115 genes upregulated in tomato roots (Wang et al. 2001) and 1 176 genes either being up or downregulated in *Arabidopsis* roots after nitrate induction, based on the mRNA hybridization (Wang et al. 2003).

The prime upshot of this study is the establishment of main effect QTLs, and their interactions to the varying nitrogen regimes, for describing the phenotypic performance of the DH population. One prominent QTL was identified with 12.43 LOD and an R^2 value of 50.3 for PHT in the present study. This QTL, *qPHT1* was located on chromosome 1 (RG690–RZ730) and found expressed in all of the nitrogen levels. This can be considered as a stable QTL. Fang and Wu (2001) found a stable QTL flanked between RZ730–RZ801, corresponding to the locus of the semi-dwarfing gene, *sd-1*, which was tightly linked to the marker RZ730 on chromosome 1 in IR64/Azucena DH line population (Yu 1991; Cho et al. 1994; Huang et al. 1996). The other two QTLs for PHT were not detected under native nitrogen, but in normal and high nitrogen regimes. Similar detection of QTLs for plant height that interacted with nitrogen levels has been reported (Fang and Wu 2001). It was seen here that chromosome 3 harbored seven QTLs out of 16 identified. Among this, the QTL for SPY was detected in all of the nitrogen regimes, while its nearest interval addressing NUE was expressed only under nitrogen-fertilized regimes. This suggested that this location might be sensitive to added nitrogen. Fang and Wu (2001) identified a plant height affecting QTL associated closely with marker RZ284, sensitive to low nitrogen levels under nutrient solution culture and soil culture. Abundance of QTLs affecting nitrogen-related traits on chromosome 3 has also been reported recently by Laza et al. (2006). These QTLs were found to be associated with nitrogen uptake, shoot nitrogen content, plant height, filled grains per panicle, panicle number, and above ground dry weight. In this study, no QTL was identified for NUP, which recorded the highest proportion of environment effect followed by genotype \times environment effect.

It is striking to note that 11 out of 17 QTLs detected showed significant ($P < 0.05$) interaction with nitrogen levels. This indicates that significant influences are on for agronomic and nitrogen assimilation rates, at varying levels of nitrogen. Studies on quantitative genetic control of nitrogen assimilation across species show complex patterns of regulatory mechanisms. Evidence suggests that HATS and LATS genes in higher plants are more root-specific, and nitrate inducible and are repressed by nitrogen metabolites (Daniel-Vedele et al. 1998). Also, it is likely that both the transporter systems are active during the early stages of crop growth, but HATS continue to function even in the later stages. Thus HATS probably plays a major role in improving grain protein content and the number of fully developed grains (Abrol et al. 1999). Besides, key enzymes in the nitrogen assimilation pathway show complex regulatory mechanisms. Particularly, nitrate reductase is very sensitive to stress and shows low affinity to nitrate, and is unstable *in vitro*. Higher plants also harbor multiple conserved sequences of the *Nia* gene encoding for nitrate reductase, and evidence suggests that plants do not benefit from excess activity of this enzyme. However, it may be worth mentioning here that overexpression of the enzymes of ammonia assimilation increases the total protein accumulation under nitrogen deprivation and showed higher carbon dioxide assimilation (Chichkova et al. 2001). This may further support the studies of Hirel et al. (2001), wherein the QTLs were detected for physiological traits such as nitrate content, nitrate reductase, and glutamine synthetase activities in maize. Yamaya et al. (2002) have reported putative QTLs associated with the contents of glutamine synthetase, which is probably involved in the export of nitrogen from senescing organs.

It is known that nitrogen content in plants is predominantly affected by the Rubisco content, which strongly affects photosynthesis. About 50% of the total soluble protein and 25% of the total nitrogen are in Rubisco protein in rice leaves (Makino 2003). Ishimaru et al. (2001) detected QTL for Rubisco and soluble protein on chromosome 8 in rice. An additional QTL for Rubisco was also identified on chromosome 9. To have the better results, the physiological parameters should be measured by rapid nitrate tissue tests (Diphenylamine-, Merckoquant-, Reflectoquant-method) and chlorophyll meter-tests to evaluate the nitrogen-nutritional status of individual genotype across environments. The rapid nitrate tests were already used in wheat, barley, rape, sugar beet and maize in field experiments at sites differing in total nitrogen content of the soil and in mineralization rate of nitrogen.

The QTLs mapped can be regarded as intermediaries in the genetic analyses, between statistically defined loci and Mendelian genes (Paran and Zamir 2003). This multiple trait QTL mapping method will certainly improve the efficiency of QTL mapping as indicated by (Jiang and Zeng 1995). Of a total of 26 expressed candidate gene sequences associated with enzymes involved in the nitrogen metabolism in rice distributed over some

of the identified QTL positions in the present study can be compared to the loci of few expressed candidate genes involved in nitrogen metabolism in rice. We attempted to place these candidate genes by *in silico* methods on rice chromosomes 1, 3, 5, 7, 8, 9 and 11 (Figure 1). It was interesting to find that co-localization of genes putatively expressing glutamine synthetase, glutamate synthase, nitrate transporter, ammonium transporter and 14-3-3-like protein along with the detected QTLs for nitrogen use efficiency and plant yield. Furthermore, high level of QTL effects and QTL × environment interactions observed in this study implies manifold regulation of nitrogen response genes in rice. This could be facilitated additionally by QTLs distributed outside the vicinity of candidate gene loci. This corroborates the need to generate precise information about the QTL mapped for nitrogen response in rice using explicit phenotypic parameters and accurate phenotyping, which will positively lead to the establishment of functionality to the genes of nitrogen metabolism by combining various approaches such as QTL mapping, candidate genes mapping and *in silico* biology.

Materials and Methods

Plant materials and environments

A subset of 82 double haploid rice (*Oryza sativa* L.) lines drawn out of 135 lines obtained from the International Rice Research Institute, the Philippines was used in this study. These lines were generated through *in vitro* anther culture of F₁ between IR64, an *indica* variety adapted to irrigated conditions and Azucena, a traditional upland *japonica* variety from the Philippines (Guiderdoni et al. 1992).

Sowing of the selected lines along with their parents was carried out in seedling boxes. Twenty-day-old seedlings of each line were planted in pots measuring 25 × 30 cm filled with approximately 4.5 kg of soil (12.74% moisture). The soil was a nitrogen-deficient clay loam falling under the group Typic Haplustalf with an ammoniacal N (KCl extractable) concentration of 37.07 µg/g of soil and available N (KMnO₄-N) concentration of 277 kg/ha. A single plant was maintained in each pot from seedling to maturity. A total of nine pots were maintained for each line split into three sets corresponding to different nitrogen levels (native, optimum and high) with three replications per level.

Three levels of nitrogen, namely, native (low), optimum and high were maintained. Native level was not supplied with any nitrogen fertilization (0 kg/ha); optimum level was applied with 1 630 mg of nitrogen fertilizer (750 mg N) per pot in the form of urea, which is the normal recommended level for rice (100 kg/ha) and high level was added with 3 260 mg of urea per pot (1 500 mg N) that was two times more than the normal dose (200 kg/ha). Nitrogen was applied in four split doses: 25% as

basal and 25% each at tillering, panicle initiation and heading stages. Other major nutrients, phosphorus and potassium were applied to all of the pots in the form of superphosphate and muriate of potash at the rate of 2 345 mg (375 mg P₂O₅) and 625 mg (375 mg K₂O) per pot, respectively. All fertilizers were applied in the form of aqueous solution.

Agronomic traits

Nine traits of agronomic importance that are presumably affected by nitrogen nutrition, namely, days-to-flowering (DTF), plant height (PHT), total number of tillers including non-productive ones (TLN), number of productive tillers alone (PTN), panicle length (PLG), spikelets per primary panicle (SPP), filled grains per primary panicle (FGP), spikelet fertility in percentage (SFP) and grain yield (SPY) were measured on a single plant basis from all three replications across different nitrogen levels. At maturity, the plants were cut just above ground level. The grains were threshed manually and sun-dried. Filled and unfilled grains were separated manually. Grain yield at 14% moisture level was recorded.

Nitrogen analysis

The dry weight of filled grains, unfilled grains, straw and rachis were determined after oven drying to a constant weight at 70 °C. The oven-dried unfilled grains, straw, and rachis were pooled and ground in a cyclone mill and the grains were ground separately to obtain fine powdered samples. Three grams of composite sample was prepared by taking a portion of the straw sample (straw + unfilled grain + rachis) and the grain sample proportionate to their weight and stored in tightly sealed vials. A sub-sample of 0.5 g was used to determine total nitrogen content by the Microkjeldhal method (Humphries 1956). The biomass of filled grains, unfilled grains, rachis and straw were added and expressed as plant dry weight (g/plant) on oven-dry weight basis (70 °C). Three parameters, namely, percent nitrogen content in plant (percentage of nitrogen in the pooled sample of straw and grain, NCP), total nitrogen uptake (plant dry weight (g/plant) × percent nitrogen content in plant, NUP) and nitrogen use efficiency (grain yield (g/plant)/total nitrogen uptake, NUE) were computed.

Data analysis

The data on individual traits under varying nitrogen regimes were subjected to analysis of variance, association analysis, principal component analysis and QTL analyses. The phenotypic and genotypic variance and heritability of the traits were computed. The frequency distribution of the traits in the mapping population was also determined.

The molecular marker data developed by Huang et al. (1997) for IR64/Azucena mapping population with 175 markers including 135 restriction fragment length polymorphism (RFLP) markers and 40 isozyme and/or Random Amplified Polymorphic DNA (RAPD) markers was used to identify putative QTL for various traits involved in the study. The mean phenotypic values from three replications for all 12 parameters under three different nitrogen regimes were subjected to composite interval mapping using multi-environment analysis option of PLABQTL v1.2 (Utz and Melchinger 2003). To get an unbiased estimate of LOD threshold for each trait, 1 000 permutations were run using a permutation test (Churchill and Doerge 1994). The detected QTLs were named as per McCouch et al. (1997), with a “q” prefixed to the trait name followed by the chromosome number separated by a hyphen (-). If more than one QTL was located on the same chromosome for the same trait, the QTLs were serially numbered and prefixed by another hyphen (-).

References

- Abdi H** (2003). Factor rotation. In: Lewis-Beck M, Bryman A, Futing T, eds. *Encyclopedia for Research Methods for the Social Sciences*. Sage, Thousand Oaks, CA, USA. pp. 792–795.
- Abrol YP, Chatterjee SR, Kumar PA, Jain V** (1999). Improvement in nitrogen use efficiency: physiological and molecular approaches. *Curr. Sci.* **76**, 1357–1364.
- Chichkova S, Arellano J, Vance CP, Hernández G** (2001). Transgenic tobacco plants that overexpress alfalfa NADH-glutamate synthase have higher carbon and nitrogen content. *J. Exp. Bot.* **52**, 2079–2087.
- Cho YG, Eun MY, McCouch SR, Chea YA** (1994). The semi dwarf gene, *sd-1*, of rice (*Oryza sativa* L.). II. Molecular mapping and marker assisted selection. *Theor. Appl. Genet.* **89**, 54–59.
- Churchill GA, Doerge RW** (1994). Empirical threshold values for quantitative trait mapping. *Genetics* **138**, 963–971.
- Conway GR, Pretty JN** (1988). Fertilizer risks in developing countries. *Nature* **334**, 207–208.
- Crawford NM, Glass ADM** (1998). Molecular and physiological aspects of nitrate uptake in plants. *Trends Plant Sci.* **3**, 389–395.
- De Datta SK, Buresh RJ** (1989). Integrated nitrogen management in irrigated rice. *Adv. Soil Sci.* **10**, 143–169.
- Daniel-Vedele F, Filleur S, Caboche M** (1998). Nitrate transport: a key step in nitrate assimilation. *Curr. Opin. Plant Biol.* **1**, 235–239.
- FAO** (2002). Concern about rice production practices. Retrieved 9 November 2007, from Food and Agriculture Organization of the United Nations' website: <http://www.fao.org/english/newsroom/news/2002/7538-en.html>
- Fang P, Wu P** (2001). QTL × N-level interaction for plant height in rice (*Oryza sativa* L.). *Plant Soil* **236**, 237–242.
- Gallais A, Hirel B** (2004). An approach to the genetics of nitrogen use efficiency in maize. *J. Exp. Bot.* **55**, 295–306.

- Glass ADM, Britto DT, Kaiser BN, Kinghorn JR, Kronzucker HJ, Kumar A et al.** (2002). The regulation of nitrate and ammonium transport systems in plants. *J. Exp. Bot.* **53**, 855–864.
- Gerendas JG, Zhu Z, Sattelmacher B** (1998). Influence of N and Ni supply on nitrogen metabolism and urease activity in rice (*Oryza sativa* L.). *J. Exp. Bot.* **49**, 1545–1554.
- Glevarec G, Bouton S, Jaspard E, Riou MT, Cliquet JB, Suzuki A et al.** (2004). Respective roles of the glutamine synthetase/glutamate synthase cycle and glutamate dehydrogenase in ammonium and amino acid metabolism during germination and post-germinative growth in the model legume *Medicago truncatula*. *Planta* **219**, 286–297.
- Lillo C** (2004). Light regulation of nitrate uptake, assimilation and metabolism. In: Amancio S, Stule I, eds. *Plant Ecophysiology, Vol 3: Nitrogen Acquisition and Assimilation in Higher Plants*. Kluwer Academic Publishers, Dordrecht. pp. 149–184.
- Guiderdoni E, Galinato E, Luistro J, Vergava G** (1992). Anther culture of tropical japonica × indica hybrids of rice (*Oryza sativa* L.). *Euphytica* **62**, 219–224.
- Hirel B, Bertin P, Quillere I, Bourdoncle W, Attagnant C, Dellay C et al.** (2001). Towards better understanding of the genetic and physiological basis for nitrogen use efficiency in maize. *Plant Physiol.* **125**, 1258–1270.
- Huang N, Courtois B, Khush GS, Lin H, Wang G, Wu P et al.** (1996). Association of quantitative trait loci for plant height with major dwarfing genes in rice. *Heredity* **77**, 130–137.
- Huang N, Parco A, Mew T, Magapantay G, McCouch SR, Guiderdoni E et al.** (1997). RFLP mapping of isozymes, RAPD, and QTLs for grain shape, brown planthopper resistance in a doubled haploid rice population. *Mol. Breeding* **3**, 105–113.
- Humphries EC** (1956). Mineral components and ash analysis. *Soil Sci.* **100**, 112–117.
- Ishimaru K, Kobayashi N, Ono K, Yano M, Ohsugi R** (2001). Are contents of rubisco, soluble protein and nitrogen in flag leaves of rice controlled by the same genetics? *J. Exp. Bot.* **52**, 1827–1833.
- Jiang C, Zeng ZB** (1995). Multiple trait analysis of genetic mapping for quantitative trait loci. *Genetics* **140**, 1111–1127.
- Kronzucker HJ, Siddiqi MY, Glass ADM, Kirk GJD** (1999). Nitrate-ammonium synergism in rice. A subcellular flux analysis. *Plant Physiol.* **119**, 1041–1045.
- Laza MR, Kondo M, Ideta O, Barlaan E, Imbe T** (2006). Identification of quantitative trait loci for $\delta^{13}\text{C}$ and productivity in irrigated lowland rice. *Crop Sci.* **46**, 763–773.
- Lian X, Xing Y, Yan H, Xu C, Li X, Zhang Q** (2005). QTLs for low nitrogen tolerance at seedling stage identified using a recombinant inbred line population derived from an elite rice hybrid. *Theor. Appl. Genet.* **112**, 85–96.
- Loudet O, Chaillou S, Merigout P, Talbotec J, Daniel-Vedele F** (2003). Quantitative trait loci analysis of nitrogen use efficiency in *Arabidopsis*. *Plant Physiol.* **131**, 345–358.
- McCouch SR, Cho YG, Yano M, Paul E, Blinstrub M, Morishima H et al.** (1997). Suggestions for QTL nomenclature for rice. *Rice Genet. Newsl.* **14**, 11–13.
- Mattson WJ, Haack RA, Lawrence RK, Slocum SS** (1991). Considering the nutritional ecology of the spruce budworm in its management. *For. Ecol. Manage.* **39**, 183–210.
- Makino A** (2003). Rubisco and nitrogen relationships in rice: leaf photosynthesis and plant growth. *Soil Sci. Plant Nutr.* **49**, 319–327.
- Meyer C, Stitt M** (2001). Nitrate reduction and signalling. In: Lea PJ, Morot-Gaudry JF, eds. *Plant Nitrogen*. Springer-Verlag, Berlin. pp. 37–59.
- Obara M, Sato T, Sasaki S, Kashiba K, Nagano A, Nakamura I et al.** (2004). Identification and characterization of a QTL on chromosome 2 for cytosolic glutamine synthetase content and panicle number in rice. *Theor. Appl. Genet.* **110**, 1–11.
- Paran I, Zamir D** (2003). Quantitative traits in plants: beyond the QTL. *Trends Genet.* **19**, 303–306.
- Rauh BL, Basten C, Buckler ES** (2002). Quantitative trait loci analysis of growth response to varying nitrogen sources in *Arabidopsis thaliana*. *Theor. Appl. Genet.* **104**, 743–750.
- Rieseberg LH, Archer MA, Wayne RK** (1999). Transgressive segregation, adaptation and speciation. *Heredity* **83**, 363–372.
- Socolow RH** (1999). Nitrogen management and the future of food: lessons from the management of energy and carbon. *Proc. Natl. Acad. Sci. USA* **96**, 6001–6008.
- Tirol-Padre A, Ladha JK, Singh U, Laurels E, Punzalan G, Akita S** (1996). Grain yield performance of rice genotypes at sub optimal levels of soil N as affected by uptake and utilization efficiency. *Field Crops Res.* **46**, 127–142.
- Wang MY, Siddiqi MY, Ruth TJ, Glass ADM** (1993). Ammonium uptake by rice roots. I. Fluxes and subcellular distribution of $^{13}\text{NH}_4^+$. *Plant Physiol.* **103**, 1249–1258.
- Wang YH, Garvin DF, Kochian LV** (2001). Nitrate-induced genes in tomato roots. Array analysis reveals novel genes that may play a role in nitrogen nutrition. *Plant Physiol.* **127**, 345–359.
- Wang R, Okamoto M, Xing X, Crawford NM** (2003). Microarray analysis of the nitrate response in *Arabidopsis* roots and shoots reveals over 1,000 rapidly responding genes and new linkages to Glucose, Trehalose-6-Phosphate, iron, and sulfate metabolism. *Plant Physiol.* **132**, 556–567.
- Yu Z** (1991). *Molecular Mapping of Rice (Oryza sativa L.) Genes via Linkage to Restriction Fragment Length Polymorphism (RFLP) Markers*. Ph.D Dissertation. Graduate School, Cornell University, Ithaca.
- Yamaya T, Obara M, Nakajima H, Sasaki S, Hayakawa T, Sato T** (2002). Genetic manipulation and quantitative trait loci mapping for nitrogen recycling in rice. *J. Exp. Bot.* **53**, 917–925.
- Utz HF, Melchinger AE** (2003). *PLABQTL: A Computer Program to Map QTL, Version 1.2*. Institute of Plant Breeding, Seed Science, and Population Genetics, University of Hohenheim, Stuttgart, Germany.