

Evaluation of Wild *Cicer* Species for Resistance to Ascochyta Blight and Botrytis Gray Mold in Controlled Environment at ICRISAT, Patancheru, India

S Pande^{1,*}, D Ramgopal², GK Kishore¹, N Mallikarjuna¹, M Sharma¹, M Pathak¹ and J Narayana Rao¹
(1. ICRISAT, Patancheru 502 324, Andhra Pradesh, India; 2. Acharya NG Ranga Agricultural University, College of Agriculture, Rajendranagar, Hyderabad 500 030, Andhra Pradesh, India)

*Corresponding author: s.pande@cgiar.org

Chickpea (*Cicer arietinum* L.) is the third most important food legume crop grown over 45 countries across five continents. It maintains soil fertility through biological nitrogen fixation and contributes to the sustainability of cropping systems in cereal-legumes rotation.

Ascochyta blight (AB, caused by *Ascochyta rabiei*) and Botrytis gray mould (BGM, caused by *Botrytis cinerea*) are destructive fungal foliar diseases of chickpea (Davidson et al. 2004; Pande et al. 2004 and Pande et al. 2005) that can cause up to 100% yield losses. Cool and wet weather favour these diseases and their epidemic development. Management of AB and BGM rely on fungicides, but these are not effective when the disease pressure is high. Deployment of resistant genotypes could be an effective way to minimize yield losses due to AB and BGM. Since adequate levels of disease resistance are not available in the cultivated chickpea germplasm, wild *Cicer* spp. have been identified

as good sources of resistance to these diseases and there is a potential to transfer resistance genes from these species into cultivated *C. arietinum* species (Singh et al. 1992 and Haware et al. 1992). Therefore, in our quest to identify durable levels of resistance to AB and BGM, we initiated a large-scale screening of wild *Cicer* accessions under optimal disease development conditions at ICRISAT.

Ascochyta blight. Following the controlled environment screening technique (CEST), 148 wild accessions belonging to seven *Cicer* spp. viz., *C. bijugum*, *C. cuneatum*, *C. echinospermum*, *C. judaicum*, *C. pinnatifidum*, *C. reticulatum* and *C. yamashitae* were evaluated for AB resistance. Eight seedlings each of the test entry and a susceptible genotype were raised in rows in plastic trays filled with sand-vermiculite mixture (4:1) in a greenhouse. Nine test entries and a susceptible check Pb7 were sown in each tray. These trays with 12-day-old seedlings were transferred to controlled environment facility (CEF) maintained at $20\pm1^\circ\text{C}$ and ~ 1500 Lux light intensity for 12 h a day, allowed to acclimatize for 24 h and inoculated with the conidial suspension (5×10^4 conidia ml⁻¹) till runoff. The *A. rabiei* conidia were produced on the autoclaved seeds of chickpea and harvested into sterile distilled water to prepare the conidial suspension for inoculation. After inoculation, the seedlings were allowed to dry partially for 30 min; thereafter 100% relative humidity (RH) was maintained till the end of the experiment. Disease severity was recorded on a 1–9 rating scale 10 days after inoculation (Pande et al. 2005). The experiment was repeated once. Based on the mean disease score of two repetitions (16 seedlings), individual chickpea lines were categorized as asymptomatic

Table 1. Evaluation of wild *Cicer* accessions for resistance to Ascochyta blight in controlled environment.

<i>Cicer</i> species	No. of lines tested	Reaction to Ascochyta blight infection ^a				
		A	R	MR	S	HS
<i>C. bijugum</i>	30	—	—	7	20	3
<i>C. cuneatum</i>	3	—	—	1	2	—
<i>C. echinospermum</i>	4	—	—	—	3	1
<i>C. judaicum</i>	47	—	5	34	8	—
<i>C. pinnatifidum</i>	27	—	—	13	13	1
<i>C. reticulatum</i>	31	—	—	—	15	16
<i>C. yamashitae</i>	6	—	—	—	—	6
Total	148	—	5	55	61	27

a. Based on the disease score the wild accessions were categorized for their reaction to Ascochyta blight infection as follows: 1 = asymptomatic (A); 1.1–3.0 = resistant (R); 3.1–5.0 = moderately resistant (MR); 5.1–7.0 = susceptible (S); 7.1–9.0 = highly susceptible (HS).

Table 2. Evaluation of wild *Cicer* accessions for resistance to Botrytis gray mold in controlled environment.

<i>Cicer</i> species	No. of lines tested ^b	Reaction to Botrytis gray mold infection ^a				
		A	R	MR	S	HS
<i>C. bijugum</i>	28	—	3	18	7	—
<i>C. cuneatum</i>	3	—	—	3	—	—
<i>C. echinospermum</i>	2	—	—	1	—	1
<i>C. judaicum</i>	45	—	23	18	4	—
<i>C. pinnatifidum</i>	26	—	—	4	20	2
<i>C. reticulatum</i>	27	—	3	6	18	—
<i>C. yamashitae</i>	5	—	—	—	2	3
Total	136	—	29	50	51	6

a. Based on the disease score the wild accessions were categorized for their reaction to Botrytis gray mold infection as follows: 1 = asymptomatic (A); 1.1–3.0 = resistant (R); 3.1–5.0 = moderately resistant (MR); 5.1–7.0 = susceptible (S); 7.1–9.0 = highly susceptible (HS).
b. 12 lines did not germinate.

Table 3. Identification of combined resistance to Ascochyta blight and Botrytis gray mold diseases in controlled environment.

Accession No ¹	Disease reaction (on 1–9 rating scale)					
	Ascochyta blight			Botrytis gray mold		
	Test 1	Test 2	Mean	Test 1	Test 2	Mean
ICC 17211	2.7	2.0	2.3	4.0	3.0	3.5
IG 69986	2.5	3.5	3.0	4.5	2.5	3.5
IG 70030	3.5	2.5	3.0	4.5	2.5	3.5
IG 70037	2.0	4.0	3.0	4.0	2.0	3.0
IG 70038	2.7	3.0	2.8	3.5	2.0	2.8

1. All accessions belong to *Cicer judaicum*.

(disease score 1.0), resistant (disease score 1.1–3.0), moderately resistant (disease score 3.1–5.0), susceptible (disease score 5.1–7.0) and highly susceptible (disease score 7.1–9.0).

Out of 148 accessions evaluated, five accessions of *C. judaicum* (ICC 17211, IG 69986, IG 70030, IG 70037 and IG 70038) were resistant. Of the remaining lines, 55 accessions were moderately resistant, 61 were susceptible and 27 were found to be highly susceptible to AB infection (Table 1).

Botrytis gray mold. One hundred and forty-eight wild *Cicer* accessions belonging to seven *Cicer* spp. viz., *C. bijugum*, *C. cuneatum*, *C. echinospermum*, *C. judaicum*, *C. pinnatifidum*, *C. reticulatum* and *C. yamashitae* were raised similar to AB resistance screening procedures in

the greenhouse and tested for BGM resistance in CEF. There were eight seedlings of each of the nine test genotypes and a BGM susceptible line (JG 62 as indicator) in each tray. Trays with 12-day-old seedlings were transferred to CEF adjusted at $15\pm2^\circ\text{C}$ and ~ 1500 Lux light intensity for 12 h a day, allowed to acclimatize for 24 h and inoculated with the conidial suspension (3×10^5 conidia ml^{-1}) till runoff. After inoculation, the seedlings were allowed to dry for 30 min; thereafter 100% RH was maintained till the end of experiment. The *B. cinerea* inoculum was multiplied on autoclaved petals of marigold (*Tagetes erecta*) flowers for 8 days at 25°C and 12 h photoperiod. Conidia from the profusely sporulating culture were harvested into sterile distilled water and used for inoculations. The experiment was repeated once. Disease severity was recorded on a 1–9 rating scale as

done for AB at 20 DAI, and based on the mean disease score of two repetitions (16 seedlings) individual chickpea lines were categorized as asymptomatic, resistant, moderately resistant and susceptible or highly susceptible.

Of the 148 wild accessions evaluated, 29 accessions were found to be resistant. Out of 29 resistant accessions 23 were from *C. judaicum* (ICC 17194, ICC 17205, ICC 17149, ICC 17148, ICC 17204, IG 69977, IG 70033, IG 72931, IG 72932, IG 17150, IG 69959, IG 69969, IG 70032, IG 70038, ICC 17151, ICC 17190, ICC 17192, ICC 17195, IG 69943, IG 69997, IG 69998, IG 70034 and IG 70037); three from *C. bijugum* (IG 69981, IG 70023 and IG 70006) and three from *C. reticulatum* (IG 72959, IG 72933 and IG 72941). The remaining 107 were categorized as moderately resistant (50), susceptible (51) and highly susceptible (6) to BGM (Table 2). Twelve lines did not germinate.

Ascochyta blight and Botrytis gray mold. Five AB resistant accessions belonging to *C. judaicum* (ICC 17211, IG 69986, IG 70030, IG 70037 and IG 70038) were separately evaluated for AB and BGM twice in the CEF to identify combined resistance to both the diseases. Procedures for raising the seedlings, inoculum preparation, inoculations, and disease scoring were similar to AB and BGM evaluations explained earlier. Two accessions (IG 70037 and IG 70038) were found to be resistant (≤ 3.0 , on 1–9 scale) to both the diseases and the remaining three (ICC 17211, IG 69986 and IG 70030) were moderately resistant (Table 3). These wild *Cicer*

accessions, found resistant to AB, BGM and/or to both the diseases, can be used in the chickpea foliar disease resistance breeding programs as resistant donor parents.

References

- Davidson JA, Pande S, Bretag TW, Lindbeck KD and Kishore GK.** 2004. Biology and management of *Botrytis* spp. in legume crops. Pages 295–318 in *Botrytis: Biology, Pathology and Control* (Elad Y, Williamson B, Tudzynski P and Delen N, eds.). The Netherlands: Kluwer Academic Publishers.
- Haware MP, Narayana Rao J and Pundir RPS.** 1992. Evaluation of wild *Cicer* species for resistance to four chickpea diseases. International Chickpea Newsletter 27:16–18.
- Pande S, Kishore GK, Ramsay G, Williamson B, Senthil G, Shivram LP, Mallikarjuna N, Gaur PM and Rao JN.** 2004. Biology and epidemiology of botrytis grey mould of chickpea. Page 10 in Proceedings of the XIII International Botrytis Symposium, 25–31 October 2004, Antalya, Turkey.
- Pande S, Siddique KHM, Kishore GK, Baaya B, Gaur PM, Gowda CLL, Bretag T and Crouch JH.** 2005. Ascochyta blight of chickpea (*Cicer arietinum* L.): A review of Biology, pathogenicity and disease management. Australian Journal of Agricultural Research 56(4):317–332.
- Singh G, Kaur L and Sharma YR.** 1992. Exploitation of host-plant resistance to manage biotic stresses in chickpea. Page 76 in Program and abstracts, Second International Food Legume Research Conference, 12–16 Apr 1992, Cairo, Egypt.