

**Published in Plant Genetic Resources : Characterization and Utilization (2011) 9(1); 59–69**

<http://dx.doi.org/10.1017/S1479262110000407>

**This is Author version post print for archiving in the official open access repository of the ICRISAT [www.icrisat.org](http://www.icrisat.org)**

**Large genetic variation for heat tolerance in the reference collection of chickpea (*Cicer arietinum* L.) germplasm.**

L. Krishnamurthy<sup>1,\*</sup>, P.M. Gaur<sup>1</sup>, P.S. Basu<sup>2</sup>, S. K. Chaturvedi<sup>2</sup>, S. Tripathi<sup>1</sup>, V. Vadez<sup>1</sup>,  
A. Rathore<sup>1</sup>, R.K. Varshney<sup>1</sup> and C.L.L. Gowda<sup>1</sup>

<sup>1</sup> International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru 502 324, Andhra Pradesh, India.

<sup>2</sup> Indian Institute of Pulses Research (IIPR), Kanpur 208 024, India. \* Corresponding author,  
Email: [l.krishnamurthy@cgiar.org](mailto:l.krishnamurthy@cgiar.org)

**Abstract:** Chickpea is the third most important pulse crop worldwide. Changes in cropping system that necessitate late planting, scope for expansion in rice fallows, and the global warming are pushing chickpeas to relatively warmer growing environment. Such changes demand identification of varieties resilient to warmer temperature. Therefore, the reference collection of chickpea germplasm, defined based on molecular characterization of global composite collection, was screened for high temperature tolerance at two locations in India (Patancheru and Kanpur) by delayed sowing and synchronizing the reproductive phase of the crop with the occurrence of higher temperatures ( $\geq 35^{\circ}\text{C}$ ). A heat tolerance index (HTI) was calculated using a multiple regression approach where grain yield under heat stress is considered as a function of yield potential and time to 50% flowering. There were large and significant variations for HTI, phenology, yield and yield components at both the locations. There were highly significant genotypic effects and equally significant G×E interactions for all the traits studied. A cluster analysis of the HTI of the two locations yielded five cluster groups as

stable tolerant (n=18), tolerant only at Patancheru (34), tolerant only at Kanpur (n=23), moderately tolerant (n=120), and stable sensitive (n=82). The pod number per plant and the harvest index explained  $\geq 60\%$  of the variation in seed yield and  $\geq 49\%$  of HTI at Kanpur and  $\geq 80\%$  of the seed yield and  $\geq 35\%$  of HTI at Patancheru indicating that partitioning as a consequence of poor pod set is the most affected trait under heat stress. A large number of heat tolerant genotypes also happened to be drought tolerant.

**Keywords:** climate change, harvest index, heat tolerance index, high temperature, shoot biomass

## **Introduction**

Chickpea (*Cicer arietinum* L.) is the third most important pulse crop globally, with a production of 9.8 M t from an area of 11.1 M ha (FAO STAT, 2009). It is even more important for India as the country's production accounts for 67% of the global chickpea production and chickpea constitutes about 40% of India's total pulse production. In spite of India being the largest chickpea producing country, a deficit exists in domestic production and demand which is met through imports.

Chickpea is a winter-season crop and often experiences increasing high temperature stress with advancing stages of crop growth. During the past three decades, there has been a significant shift in the growing environment of chickpea in India from the cooler, long-season environments of northern India to the warmer, short-season environments of central and southern India (Gaur et al., 2008; Gowda et al., 2009). Terminal drought and heat stresses are major constraints to chickpea production in warmer short-season environments. Also, the chickpea area under late-sown conditions is increasing, particularly in northern and central India, due to inclusion of chickpea in new cropping systems and intense sequential cropping practices leading to a prolonged exposure of chickpea to high temperature. Heat stress during the reproductive period is a major limitation in this situation too. It is also estimated that about 11.7 million ha of rice area in India, currently remains fallow after late harvest of rice during the winter season in the central and north-eastern India (Subbarao et al. 2001). These lands potentially offer expansion in chickpea cultivation provided genotypes capable of standing heat stress are made available. Finally, heat stress is expected to be an increasingly important constraint

in near future due to climate change and global warming. By 2050, a rise in temperature by at least 2<sup>0</sup>C, particularly the night temperatures, is being predicted with higher levels of warming in northern parts of India. It can be envisaged that the increases in temperature will have more adverse effects on cool-season crops (e.g. chickpea) than the rainy-season crops (Kumar, 2006). So, there is an urgent need to search the gene bank for diverse sources of heat tolerance. However, no such systematic search had been taken up in chickpea except for a limited effort with 25 diverse genotypes leading to the identification of two genotypes, ICCV 88512 and ICCV 88513, to have heat tolerance at reproductive stage (Dua, 2001).

Flowering and podding in chickpea is known to be very sensitive to changes in external environment, and exposure to heat stress at this stage is known to lead to reduction in seed yield (Summerfield et al., 1984). Drastic reductions in chickpea seed yields were observed when plants at flowering and pod development stages were exposed to high (35<sup>0</sup>C) temperatures (Summerfield et al., 1984, Wang et al., 2006). Heat stress is known to adversely affect pollen viability, fertilization and seed development leading to a reduced harvest index. Yet, it is still not clear how heat affects the growth and development of chickpea and whether that can explain part of the differences in seed yield under heat stress. So, a pre-requisite, before undertaking a more thorough physiological analysis of the traits involved in heat stress tolerance, is the identification of heat tolerant genotypes. Also there is an urgent need to develop simple and effective screening techniques for screening germplasm and breeding materials for reproductive stage heat tolerance in chickpea.

Therefore the objectives of this study were to develop a screening method and to screen the reference collection of chickpea germplasm in contrasting chickpea growing locations for high temperature tolerance. The reference collection is a representative subset assembled based on the molecular diversity of the global composite germplasm collection of chickpea (Upadhyaya et al. 2008). Screening of such a diverse germplasm collection has provided contrasting diverse sources of chickpea genotypes for breeding to develop

high temperature-tolerant, climate change-resilient chickpea varieties. In addition, it was also aimed to identify traits that were most closely related to seed yield under heat stress.

## Materials and Methods

### Crop management

Field evaluation of the reference collection of chickpea germplasm devoid of the wild accessions and very long duration accessions (n=280) was conducted during the post-rainy and summer season of 2009-10 in two sowing dates (normal and late sowing) on a Vertisol (fine montmorillonitic isohyperthermic typic pallustert) at ICRISAT, Patancheru (17° 30' N; 78° 16' E; altitude 549 m) in peninsular India and in an Inceptisol (Sandy loam) at the New Research Farm, Indian Institute of Pulses Research, Kanpur in northern India. The soil depth of the field used at ICRISAT was  $\geq 1.2$  m and known to retain about 230 mm of plant available water. The soil depth and maximum retainable water was 1.5 m and 180 mm at Kanpur. At ICRISAT, the field used was solarized using polythene mulch during the preceding summer to sanitize the field, particularly to eradicate wilt causing fungus *Fusarium oxysporum f. sp. ciceri*. After the soil solarization in summer, the field was kept fallow. At Kanpur, the soil was deep ploughed twice and kept fallow after harvest of greengram (mungbean) in the end of September, for another one and half months, before sowing chickpea.

At ICRISAT, the field was prepared into a broad bed and furrows with 1.2 m wide beds flanked by 0.3 m furrows for the normal time sowing, while it was 60 cm ridges and furrows for the late sowing. Surface application and incorporation of 18 kg N ha<sup>-1</sup> and 20 kg P ha<sup>-1</sup> as di-ammonium phosphate was carried out before sowing. The plot size was 4 m × 0.75 m with a 30 × 10 cm spacing for the normal sowing and 2m x 0.6m (one row) with a 60 × 10 cm spacing for the late sowing. The design was a 14 x 20 alpha design (280 accessions) with three replications in normal and two in late sowings. The normal time sown crop was grown under receding soil moisture condition without any irrigation (apart from a post-sowing irrigation) while it was optimally irrigated in late sown condition receiving irrigations on 0, 18, 30, 35, 45, 55, 65 and 75 days after sowing. Seeds were treated with 0.5% Benlate® (E.I. DuPont India Ltd., Gurgaon, India) + Thiram® (Sudhama Chemicals Pvt. Ltd. Gujarat, India) mixture in both the sowings. The normal sown experiment was planted on 31 Oct 2009 in 30 × 10cm spacing and the late sown one on 2 Feb 2010 in 60 × 10cm spacing with

two seeds per hill that was later thinned to one. During both the plantings, the fields were inoculated with Rhizobium strain IC 59 using liquid inoculation method (Brockwell, 1982). A 50 mm irrigation through perforated pipes was applied the next day to ensure complete emergence. Need-based insecticide sprays against pod borer (*Helicoverpa armigera*) were provided and the plots were kept weed free by manual weeding.

At Kanpur, both the normal and late sowings were sown on a flat bed with a plant spacing of 60 x 10 cm and a plot size of 3 x 0.6 m on 13 Nov 2009 and 13 Jan 2010, respectively. The experiments were planted in an 8 x 35 alpha design (280 accessions) with three replications. The seeds were treated with Bavistin (BASF India Ltd, Panoli, Bharuch, Gujarat) containing carbendazim 50% W/P @ 1g per 100 g seeds and was hand planted with more than 50 seeds on a row and later thinned to maintain approximately 10 cm distanced plants. After presowing irrigation, a 50 mm irrigation through surface irrigation was applied on 2 Feb 2010 (80 days after sowing) for the normal sowing but three such irrigations on 2 Feb, 12 Mar and 26 Mar 2010 (19, 37 and 50 days after sowing) were applied for the late-planted crop. Although pod borer (*Helicoverpa armigera*) is not a major pest in Kanpur, Endosulfan, EC 35%, (Excel Crop Care, Limited, Mumbai) (at 2 ml L<sup>-1</sup> of water) was sprayed when 1 to 2 larvae plot<sup>-1</sup> were noticed, more as a prophylactic pest control measure. Pre-emergence weedicide Pendimethalin @ 3ml litre<sup>-1</sup> was applied immediately after sowing the crop. Manual weeding was followed thereafter at regular intervals.

### **Phenology**

By regular observation, the date when 50% or more of the plants in a plot flowered was recorded as 50% flowering time of the plot and when 80% of the pods in a plot were mature was recorded as the time of maturity for each plot.

### **Final harvest**

At physiological maturity, plant aerial parts were harvested from an area of 4 m x 0.75 m (3.0 m<sup>2</sup>) under normal sowing and 4m x 0.6 m (2.4 m<sup>2</sup>) under late sown condition in Patancheru and 3 m x 0.6 m (1.8 m<sup>2</sup>) under both normal and late-sown conditions in Kanpur in each plot, dried to constant weight in hot air dryers at 80°C, and total shoot dry weights were recorded. Grain weights were recorded after threshing. Harvest index (%) was calculated as  $100 \times (\text{seed yield}/\text{total shoot biomass at maturity})$ .

### **Heat tolerance index (HTI) estimation**

Differences in crop duration and yield potential (Saxena et al., 1987) are known to contribute to the seed yield under both drought and salinity stress and the removal of these effects from seed yield under stress provides a reliable measure of stress tolerance *per se* (Vadez et al., 2007). Similar escape mechanism is also expected with heat. Since the temperature increased linearly during the late planting period and all the short-duration genotypes could start flowering and filling seeds even before the temperatures increased to critical levels (Saxena et al., 1987). Previous work related to drought has shown that the residual yield remaining unexplained after removal of effects due to drought escape (early flowering) and yield potential (optimally irrigated yield) of a genotype gave a good indication of the true drought tolerance of that genotype (Bidinger et al., 1987; Saxena et al., 1987; Saxena, 2003; Vadez et al., 2007; Krishnamurthy et al. 2010). These residuals were calculated using the multiple regression approach of Bidinger et al. (1987). This approach considers grain yield under drought stress condition ( $Y_s$ ) as a function of yield potential ( $Y_p$ ), time to 50% flowering (F), and a drought tolerance index (DTI) such that the yield of a genotype can be expressed as follows:

$$Y_{si} = a + bY_p + cF_i + DTI_i + E,$$

where E is random error with zero mean and variance  $\sigma$ . The Drought Tolerance Index (DTI), was calculated as the difference between the actual and estimated yields under stress upon the standard error of the estimated yield ( $\sigma$ ). For this multiple regression, 50% flowering ( $F_i$ ) under stress for every individual plot, and for yield potential ( $Y_p$ ) arithmetic mean across the three replications were considered. Similar approach was

adopted for estimating HTI, as flowering time and yield potential are expected to determine the yields of genotypes that are limited by heat stress.

### **Statistical analysis**

The replication-wise values of HTI along with other traits were used for statistical analysis of each environment using ReML (Harville 1977) considering genotypes as random. Variance components due to genotypes ( $\sigma^2_g$ ) and error ( $\sigma^2_e$ ) and their standard errors were determined. Environment wise best linear unbiased predictors (BLUPs) for the germplasm accessions of the reference collection were calculated for the different environments. The significance of genetic variability among accessions was assessed from the standard error of the estimate of genetic variance  $\sigma^2_g$ , assuming the ratio  $\sigma^2_g/SE(\sigma^2_g)$  to follow normal distribution asymptotically.

While pooling the data over two sites, Bartlett (Snedecor and Cochran, 1983) test indicated heterogeneity in error variances. Appropriate transformation was applied and data was tested for presence of G×E interaction. Upon detection of significant G×E interaction, data from each site was analyzed individually and significance of genotypes and their relative ranks were obtained. Spearman's rank correlation coefficient was calculated to have an idea of difference in genotype ranking over sites. Cluster analysis using Ward's incremental sum of squares method was employed to group the genotypes over sites for HTI. All statistical analyses were carried out using Genstat, Release 10.1 (Payne 2002).



## **Results**

### **Variation in weather**

The late sown crop, subjected to heat stress, was sown on 2 Feb 2010 at Patancheru, and on 13 Jan 2010 at Kanpur. A 20-day early sowing date for Kanpur was chosen as the crop duration in general is longer (by 20 to 30 days) at Kanpur compared to Patancheru; and thus the heat stress imposition is applied at the same phenological stage across locations. The maximum temperature reached the threshold level of 35°C at 27 DAS in Patancheru while at 60 DAS at Kanpur (Fig. 1). Also the minimum temperatures were higher than 17°C after this stage at both locations. At the mean flowering time (52 DAS) in Patancheru the maximum air temperature had reached to 39°C while it was much less (31°C) at mean flowering time (56DAS) at Kanpur.

### **Variation in phenology in the reference collection accessions**

There were large and highly significant differences in flowering time of the accessions in both the sowing times and locations. All the genotypes tend to mature more or less close to each other, irrespective of their differences in flowering time at Kanpur. The overall means for each sowing time had shown that late sowing delayed the days to 50% flowering while the days to maturity was hastened at Patancheru. However, both these stages were reached earlier with late sowing in Kanpur (Table 1). In terms of thermal time (growing degree days, °Cd) taken to reach mean flowering it was 1094 °Cd under normal sowing, while it was 1377 °Cd under late sown condition at Patancheru. Such increase in requirement of thermal time to attain any developmental stage by the higher soil moisture grown crop is well documented (Desclaux and Roumet, 1996; Krishnamurthy et al. 1999). However this requirement was mainly to negate the irrigation-led cooling of the microclimate around the plants which is shown to be about 10°C cooler soil temperature (Reddy et al. 1989). At Kanpur, the late sown crop took 1032 °Cd and the normal sown 1486°Cd. Providing optimum irrigation is known to extend the growth duration substantially in chickpea. The late sown crop at Patancheru received irrigations at 8-12 day intervals during the whole growing period while the normal sowing conditions was grown under residual moisture stress. Similarly, the crop

at Kanpur received only 3 irrigations during the vegetative growth period. There was large range of variation for flowering time under late sown conditions in Patancheru (37 to 73 d) as well as in Kanpur (49 to 67 d) leading to an increased level temperature exposure with the delay in flowering time leading to partial disadvantages of the later genotypes.

### **Influence of flowering time and normal sown yield on late sown yield**

At Patancheru, seed yield under heat stress was negatively associated to the time to flowering ( $r^2 = 0.51^{***}$ ; significant above 0.001 level), while it was positively associated with the normal-sown seed yields, considered here as potential normal yield ( $r^2 = 0.50^{***}$ ). Similar significant negative association with 50% flowering time ( $r^2 = 0.18^{***}$ ) and yield under normal sowing ( $r^2 = 0.09^{**}$ ) was also seen at Kanpur. Therefore, categorization of the accessions in terms of seed yield under heat stress for heat response would partly lead to a categorization for escape from heat and yield potential. Therefore, heat tolerance indices were computed to characterize the heat tolerance *per se* in this study, i.e. the proportion of the genetic variation for seed yield under heat that was not accounted for differences in time to flowering and yield potential.

### **Variation in yield and yield components**

Between the two locations the shoot biomass and yield produced in Kanpur was manifolds less than that at Patancheru. This was due to a combination of effects that did not promote a normal crop growth such as broader spacing practiced, sandy and poor water holding nature of the soil, recently developed marginal land and receding soil moisture conditions during major reproductive growth with only three supplementary irrigations after sowing (Table 2). Under heat stress conditions in Patancheru, the shoot biomass produced was higher than the normal sown crop as the heat stressed crop was optimally irrigated, while the normal sown one was on receding soil moisture condition. However, mean seed yield of all the accessions were reduced to two-thirds. In Kanpur the shoot biomass was reduced by half under heat stress and the seed yield by one-fourth (Table 2). The overall harvest indices were lower under heat stress compared to the normal sown conditions and it was higher in Patancheru in any of the sowing conditions.

There were highly significant variations for the shoot biomass as well as seed yield across the accessions and these variations were about two-fold for the shoot biomass at maturity at both heat stressed as well as normal sown crop and many-fold for seed yield among the accessions again at both the sowing times tested (Table 2). There was a highly significant, large range of variation in harvest index in both the sowing times and locations. At Patancheru, the variance component for the HTI of accessions (0.585, SE 0.070) was highly significant and the means ranged from -2.5 to 2.7. Similarly at Kanpur, the variance component (0.298, SE 0.041) was highly significant for HTI and the means ranged from -0.7 to 1.9. The pooled analysis of data from both the locations had revealed that there were highly significant genotype effects and also equally significant genotype  $\times$  location (G $\times$ E) interactions for all the characteristics that were studied except for one yield component, seeds per pod. In spite of this interaction the rank correlation of the accession means between the location had indicated that 50% flowering ( $r = 0.51^{***}$ ), seed yield  $\text{g m}^2$  ( $r = 0.60^{***}$ ), harvest index ( $r = 0.57^{***}$ ) and HTI ( $r = 0.27^{***}$ ) were closely related except for the shoot biomass production ( $r = 0.06^{\text{NS}}$ ).

### **Contribution of yield components**

Among the yield components, pods  $\text{plant}^{-1}$  was most affected by late sowing at Kanpur. Interestingly, late sowing did not impact pods  $\text{plant}^{-1}$  at Patancheru, potentially as a consequence of irrigation that was specific to the late sown plots (Table 3). The range in pod number  $\text{plant}^{-1}$  was large. Time to 50% flowering (representing earliness), shoot biomass at maturity, harvest index, pods borne on a plant and seed size were related either negatively or positively to the seed yield or the HTI to various degree, depending on the sowing time and the location (data not shown). However, the pod numbers per plant ( $r^2 = 0.81$  at Patancheru and 0.64 at Kanpur) and harvest index ( $r^2 = 0.92$  at Patancheru and 0.63 at Kanpur) were the two parameters that were very closely associated with the seed yield (figures not shown) and as a consequence with the HTI (Fig 2) and other related characteristics. These relationships were very close in Patancheru than at Kanpur.

### **Heat response categorization**

As there was a significant interaction between accessions and years, the HTI of the accessions were grouped into representative groups using the BLUPs for HTI by a hierarchical cluster analysis (using Ward's incremental sum of squares method) and this analysis yielded five clusters that differed significantly. Three accessions (ICC 2482, ICC 2593 and ICC 11903) that were tested in Patancheru but not in Kanpur, as three previously tested checks (ICC 5912, ICC 07110 and ICCV 92944) were included in their place. Thus the common entries across locations, those were included in this clustering exercise, were 277. Based on the extent of cluster group means of the HTI, these were identified as: 1. stable tolerant (with HTI means 0.81 in Patancheru and 1.01 in Kanpur), 2. tolerant only at Patancheru (1.04 and 0.09), 3. tolerant only at Kanpur (-0.15 and 0.71), 4. moderately tolerant (0.10 and -0.18) and 5. stable sensitive (-0.71 and -0.15). The stable tolerant group comprised of 18 accessions (Table 4), while the stable sensitive group comprised of 82 accessions out of the 277 used for clustering. For the sake of brevity, the data of 5 genotypes that were the most sensitive and made a sub-cluster within the sensitive cluster is being presented (Table 4). The tolerant only at Patancheru group was comprised of 34 accessions (Table 5), while the tolerant only at Kanpur group was comprised of 23 accessions (Table 6) and the moderately tolerant group comprised 120 entries, respectively. ICC 14778, a stable drought tolerant entry and ICC 4958, a well known drought tolerant genotype with high root mass (Krishnamurthy et al., 2010) have also ranked as stable heat tolerant entries in this study (Table 4). Ten other entries that ranked as the next order drought tolerant accessions in the previous work also appeared as stable heat tolerant ones. Similarly 13 stable sensitive entries appeared also in a previous drought tolerance assessment and 11 of them were ranked to be moderately tolerant (Data not shown).

## Discussion

This work has established the existence of a large genotypic variation for heat response in the reference collection of chickpea germplasm that represents molecular diversity of global composite collection (Upadhyaya et al. 2008). Delayed sowing for heat tolerance screening in chickpea proposed earlier (Gaur et al. 2007) was found effective in this study. Also, there are reports of successfully using a two months delayed planting than normal in a Mediterranean climate to increase the crop exposure to higher temperature with drier conditions and successfully screening 377 germplasm accessions to identify sources of tolerance (Canci and Toker, 2009). With the current understanding of available variation, ICCV 92944 is recognized to be one of the best available heat tolerant sources based on the earlier empirical selections but as this genotype is early it is also thought to escape the heat stress. However, the yield levels of at least 10 entries listed as stable ones in table 4 did possess arithmetically more yields than that of ICCV 92944 ( $183 \pm 28.6 \text{ g m}^{-2}$ ) at Patancheru while 11 did possess significantly more yields ( $17.0 \pm 5.1 \text{ g m}^{-2}$ ) at Kanpur. Moreover, major proportion of stable heat tolerant accessions or accessions that performed well under Patancheru were also drought tolerant genotypes listed in a recently published study (Krishnamurthy et al., 2010). Also, the initial screenings carried out by Dua (2001) indicate that not only drought tolerant sources (ICCV 92501-2) perform promisingly under high temperature but also some cold tolerant sources (ICCV 88512 and ICCV 88513) also do perform good under heat indicating that the tolerance mechanism can be common for both cold and hot temperatures.

Also these sources are expected to have much wider adaptability as these were selected not simply on the basis of seed yield but by heat tolerance index that is to a large extent free from the advantages of yield potential and flowering time. These genotypes represent ideal materials for further characterization of underlying mechanisms of tolerance involved. For example, ICC 14778 (Table 1), listed in this work as the stable heat tolerant one, was also a top drought tolerant accession (Krishnamurthy et al., 2010) that is known not only to yield high under drought but also to maintain a cooler canopy temperature at peak pod filling phase when many other selected drought tolerant genotypes were

relatively warmer (Kashiwagi et al., 2008). There are possibilities of finding large number of common sources of tolerance for both heat and drought. Therefore, some of the selections made for heat tolerant genotypes can also turn out to be good drought tolerant genotypes, as demonstrated by ICC 4958 and ICC 14778.

It was very clear that the pods produced per plant as indicated in previous works (Wang et al. 2006), and as a consequence the harvest index, are the primary yield components that are affected by increased levels of heat stress. The reductions in shoot biomass and seed size also tend to be the consequences of drought stress as it has happened in Kanpur in this study. Though it is necessary to understand the underlying mechanisms of tolerance, for simple and large scale screenings it may be adequate to select either for harvest index or for pod number when precise shoot biomass estimation becomes difficult.

Chickpea has been reported to be relatively sensitive in terms of membrane stability and photo system II function at high temperatures than other legumes such as groundnut, pigeonpea and soybean (Srinivasan et al., 1996). But within the cool season legumes chickpea was found to have a higher critical temperature for heat tolerance than lentil, pea and faba bean (Malhotra and Saxena, 1993) indicating this crop to be more amenable for adaptation to warmer environments.

## **Conclusions**

Large genotypic variation was available among the reference collection of chickpea germplasm for heat tolerance that underlines the utility of the reference collection for applied breeding programme. These new sources of heat tolerance can be used for physiological and genetic studies and in heat tolerance breeding. Harvest index and pod number per plant are the two key traits that can be used in selections. The heritability of yield under heat stress environment was even better than the normal growing condition offering opportunity for direct selection of yield under optimally irrigated vertisols. The HTI represented a selection index devoid of the yield potential and phenology effects and

this index potentially offers a selection criterion for adaptation to higher temperatures valid across wider agro-ecological zones.

### **Acknowledgements**

This work was funded by the Department of Agriculture and Cooperation, Ministry of Agriculture, Government of India through the Integrated Scheme of Oilseeds, Pulses, Oil palm and Maize (ISOPOM). The technical help of Mr J. Shankaraiah, Research Technician, ICRISAT and Mr Raghubanshi Singh, Miss Rinki and Mr Pankaj Kumar Singh, IIPR, Kanpur in conducting the field experiments is sincerely acknowledged.

## References

- Bidinger FR, Mahalakshmi V and Rao GDP (1987) Assessment of drought resistance in pearl millet [*Pennisetum americanum* (L.) Leeke]. II Estimation of genotype response to stress. Australian Journal of Agricultural Research. 38: 49-59.
- Brockwell J (1982) Inoculation methods for field experimenters and farmers. In: Vincent JM (ed) Nitrogen Fixation in Legumes. Academic Press, New York. pp. 211-221.
- Canci H and Toker C (2009) Evaluation of yield criteria for drought and heat resistance in chickpea. Journal of Agronomy and Crop Science 195: 47-54.
- Desclaux D and Roumet P (1996) Impact of drought stress on the phenology of two soybean (*Glycine max* L. Merr.) cultivars. Field Crops Research 46: 61-70.
- Dua RP (2001) Genotypic variations for low and high temperature tolerance in gram (*Cicer arietinum*). Indian Journal of Agricultural Sciences 71: 561-566.
- FAOSTAT (2009) Food and Agriculture Organization of the United Nations 2002. FAO Production year book. Rome, Italy: FAO. <http://apps.fao.org>.
- Gaur PM, Kumar J, Gowda CLL, Pande S, Siddique KHM, Khan TN, Warkentin TD, Chaturvedi SK, Than AM and Ketema D (2008) Breeding chickpea for early phenology: perspectives, progress and prospects. In: Kharkwal MC (ed) Food Legumes for Nutritional Security and Sustainable Agriculture, Vol. 2, New Delhi, India: Indian Society of genetics and Plant Breeding. pp. 39-48.
- Gaur PM, Srinivasan S, Gowda CLL and Rao BV (2007) Rapid generation advancement in chickpea. Journal of SAT Agricultural Research 3(1). Available online at <http://www.icrisat.org/journal/>
- Gowda CLL, Parthasarathy Rao P, Tripathy S, Gaur PM and Deshmukh RB (2009) Regional shift in chickpea production in India. In: Masood Ali, Shiv Kumar (eds) Milestones in Food Legumes Research. Indian Institute of Pulses Research, Kanpur, 208 024, India. pp. 21-35.
- Harville DA (1977) Maximum Likelihood Approaches to Variance Component Estimation and to Related Problems. Journal of the American Statistical Association 72 (358): 320-338.



- Kashiwagi J, Krishnamurthy L, Upadhyaya HD and Gaur PM (2008) Rapid screening technique for canopy temperature status and its relevance to drought tolerance improvement in chickpea. *Journal of SAT Agricultural Research* 6: pp 4.
- Kumar S (2006) Climate change and crop breeding objectives in the twenty first century. *Current Science* 90: 1053-1054.
- Krishnamurthy L, Johansen C and Sethi SC (1999) Investigation of factors determining genotypic differences in seed yield of nonirrigated and irrigated chickpea using a physiological model of yield determination. *Journal of Agronomy and Crop Science* 183: 9-17.
- Krishnamurthy L, Kashiwagi J, Gaur PM, Upadhyaya HD and Vadez V (2011) Sources of tolerance to terminal drought in the chickpea (*Cicer arietinum* L.) minicore germplasm. *Field Crops Research*. 119: 322-330 (in press). DOI 10.1016/j.fcr.2010.08.002.
- Malhotra RS and Saxena MC (1993) Screening for cold and heat tolerance in cool season food legumes. In: Singh KB and Saxena MC (eds). *Breeding for stress tolerance in cool season food legumes*. John Wiley & Sons, Chichester UK. Pp. 227-244.
- Payne RW (ed) (2002) *The guide to GenStat ® Release 6.1. Part: 2 Statistics*. VSN International Ltd, Oxford, UK.
- Reddy PV, Reddy KB, Rao GNSN (1989) Influence of soil moisture content on pod zone temperatures in groundnut. *International Arachis Newsletter* 6: 9-10.
- Saxena NP (1987) Screening for adaptation to drought: case studies with chickpea and pigeonpea. In: Saxena NP and Johansen C (eds) *Adaptation of chickpea and pigeonpea to abiotic stresses*. Proceedings of consultant's workshop. Patancheru 502 324, Andhra Pradesh, India. International Crops Research Institute for the Semi-Arid Tropics. pp. 63-76.
- Saxena NP (2003) Management of drought in chickpea – A holistic approach. In: Saxena NP (ed.). *Management of agricultural drought. Agronomic and genetic options*. Oxford & IBH Publishing co. Pvt. Ltd., New Delhi. pp. 103-122.
- Snedecor GW and Cochran WG (1989) *Statistical Methods*, Eighth Edition, Iowa State University Press.

- Srinivasan A, Takeda H and Senboku T (1996) Heat tolerance in food legumes as evaluated by cell membrane thermostability and chlorophyll fluorescence techniques. *Euphytica* 88: 35-45.
- Subbarao GV, Kumar Rao JVDK, Kumar J, Johansen C, Deb UK, Ahmed I, Krishna Rao MV, Venkataratnam L, Hebbar KR, Sai MVSR and Harris D (2001) Spatial distribution and quantification of rice-fallows in South Asia – potential for legumes, ICRISAT, Patancheru, India.
- Summerfield RJ, Hadley P, Roberts EH, Minchin FR and Rawsthorne S (1984) Sensitivities of chickpeas (*Cicer arietinum* L.) to hot temperatures during the reproductive period. *Experimental Agriculture* 20: 77-93.
- Upadhyaya HD, Dwivedi SL, Baum M, Varshney RK, Udupa SM, Gowda CLL, Hoisington DA, Singh S (2008) Genetic structure, diversity, and allelic richness in composite collection and reference set in chickpea (*Cicer arietinum* L.). *BMC Plant Biology* 8: 106.
- Vadez V, Krishnamurthy L, Serraj R, Gaur PM, Upadhyaya HD, Hoisington DA, Varshney RK, Turner NC and Siddique KHM (2007) Large variation in salinity tolerance in chickpea is explained by differences in sensitivity at the reproductive stage. *Field Crops Research* 104: 123-129.
- Wang J, Gan YT, Clarke F and McDonald CL (2006) Response of chickpea yield to high temperature stress during reproductive development. *Crop Science* 46: 2171-2178.

**Table 1**

Trial means, range of best linear unbiased predicted means (BLUPs) and analysis of variance of the 280 accessions of the reference collection of chickpea germplasm for days to 50% flowering and days to maturity in the field experiments during 2009-10 both at Patancheru and Kanpur postrainy (normal) and summer (heat stress) seasons.

Location/Sowing time	Trial mean	Range of predicted means	S.Ed	$\sigma^2_g$ (SE)
Days to 50% flowering				
Patancheru				
Heat stress	51.8	37.2 – 73.2	3.64	40.1 (4.13)
Normal	48.4	34.8 – 65.7	2.00	37.8 (3.38)
Kanpur				
Heat stress	55.7	49.3 – 66.8	1.81	8.98 (0.93)
Normal	89.4	81.6 – 102.9	2.79	22.72 (2.34)
Days to maturity				
Patancheru				
Heat stress	88.8	76.0 – 107.4	3.21	47.2 (4.51)
Normal	95.2	78.7 – 114.7	3.18	82.0 (7.41)
Kanpur				
Heat stress	NA	NA	NA	NA
Normal	NA	NA	NA	NA

**Table 2**

Trial means, range of best linear unbiased predicted means (BLUPs) and analysis of variance of the 280 accessions of the reference collection of chickpea germplasm for shoot biomass at maturity, seed yield and harvest index in the field experiments during 2009-10 both at Patancheru and Kanpur postrainy (Normal) and summer (Heat stress) seasons.

Season/Environment	Trial mean	Range of predicted means	S.Ed	$\sigma^2_g$ (SE)
Shoot biomass (g m <sup>-2</sup> )				
Patancheru				
Heat stress	473.3	356.6 – 615.6	65.8	4261 (824)
Normal	412.0	282.2 – 549.9	43.1	3031 (379)
Kanpur				
Heat stress	74.1	38.1 – 120.8	19.4	436.7 (68.7)
Normal	146.4	83.8 – 237.1	33.2	1115 (187)
Seed yield (g m <sup>-2</sup> )				
Patancheru				
Heat stress	97.9	8.0 – 265.4	28.6	4150 (384)
Normal	152	44.2 – 231.4	20.9	1343 (137)
Kanpur				
Heat stress	10.4	3.2 – 34.7	5.1	48.2 (5.7)
Normal	41.1	16.8 – 87.4	12.8	215.2 (29.0)
Harvest index				
Patancheru				
Heat stress	22.0	0.7 – 53.3	4.28	242.9 (21.2)
Normal	37.7	11.3 – 57.0	2.77	133.0 (11.6)
Kanpur				
Heat stress	13.8	5.5 – 30.5	5.15	37.5 (5.06)
Normal	27.3	14.8 – 40.2	4.90	30.8 (4.35)

**Table 3**

Trial means, range of best linear unbiased predicted means (BLUPs) and analysis of variance of the 280 accessions of the reference collection of chickpea germplasm for pod number plant<sup>-1</sup>, seed number pod<sup>-1</sup> and 100 seed weight (g) in the field experiments during 2009-10 both at Patancheru and Kanpur poststrainy (Normal) and summer (Heat stress) seasons.

Season/Environment	Trial mean	Range of predicted means	S.Ed	$\sigma^2_g$ (SE)
Pod number plant <sup>-1</sup>				
Patancheru				
Heat stress	42.5	5.3 – 126.1	13.8	777 (75.1)
Normal	42.6	21.2 – 71.2	7.4	106 (12.3)
Kanpur				
Heat stress	6.7	2.6 – 16.7	3.19	10.83 (1.74)
Normal	24.5	11.9 – 44.9	6.68	66.3 (8.60)
Seed number pod <sup>-1</sup>				
Patancheru				
Heat stress	1.20	0.6 – 1.5	0.120	0.0261 (0.0032)
Normal	1.06	0.77 – 1.37	0.106	0.0180 (0.0023)
Kanpur				
Heat stress	0.95	0.63 – 1.38	0.244	0.038 (0.0084)
Normal	1.07	0.76 – 1.39	0.134	0.024 (0.0033)
100 seed weight (g)				
Patancheru				
Heat stress	14.6	7.4 – 35.7	1.76	31.2 (2.79)
Normal	17.3	9.2 – 44.8	1.31	38.3 (3.32)
Kanpur				
Heat stress	20.0	10.1 – 39.3	6.15	31.1 (4.76)
Normal	17.2	9.8 – 38.2	2.18	37.2 (3.33)

**Table 4**

Time to flowering, shoot and seed yield at maturity and heat tolerance indices of the consistently heat tolerant (stable in both locations) and five out of 82 consistently most heat sensitive cluster group members of chickpea reference collection at Patancheru and Kanpur under heat stress during 2010 summer. Flower, shoot, yield and HTI denotes to days to 50% flowering, shoot biomass g m<sup>2</sup>, seed yield g m<sup>2</sup> and heat tolerance index, respectively and the characters P and K for Patancheru and Kanpur, respectively.

S.No	Accession	P-Flower	P-shoot	P-Yield	P-HTI	K-Flower	K-shoot	K-Yield	K-HTI
Stable heat tolerant									
1	ICC 456	50.3	412.2	145.7	0.64	54.9	81.7	19.9	0.55
2	ICC 637	53.6	514.1	139.2	0.68	54.6	102.8	26.9	1.08
3	ICC 1205	48.6	461.8	190.8	0.79	55.9	114.0	23.0	0.96
4	ICC 3362	47.4	491.3	196.3	1.28	55.7	104.3	28.9	1.34
5	ICC 3761	50.3	565.7	130.7	0.68	51.5	85.9	22.1	0.70
6	ICC 4495	49.1	503.1	199.2	1.23	55.7	73.6	21.4	0.81
7	ICC 4958	42.4	485.2	231.4	0.93	51.5	91.4	20.0	0.81
8	ICC 4991	49.9	460.4	180.6	0.50	53.7	90.6	19.7	0.81
9	ICC 6279	42.0	453.3	220.1	0.82	52.4	81.4	27.0	1.29
10	ICC 6874	49.1	503.9	195.2	1.10	55.4	92.7	21.2	0.99
11	ICC 7441	49.5	464.0	199.1	0.74	53.7	97.4	20.8	0.74
12	ICC 8950	45.7	452.4	177.3	0.81	55.1	96.7	22.7	0.90
13	ICC 11944	48.6	515.3	173.1	0.58	54.3	86.6	24.4	0.91
14	ICC 12155	44.1	433.5	191.1	1.05	51.8	97.3	34.7	1.92
15	ICC 14402	42.0	457.2	184.9	0.59	53.2	112.4	28.4	1.26
16	ICC 14778	47.8	441.8	153.5	0.50	55.4	91.7	21.1	0.89
17	ICC 14815	49.9	506.6	178.8	0.93	54.6	110.5	25.6	1.15
18	ICC 15618	48.6	431.5	187.4	0.59	52.3	112.7	23.8	1.00
	Mean	47.7	475.6	182.7	0.81	54.0	95.8	23.7	1.00
Stable heat sensitive									
1	ICC 4567	50.7	535.2	18.8	-1.80	57.9	86.5	6.5	-0.37
2	ICC 10685	47.4	520.7	8.6	-1.88	55.9	85.4	4.3	-0.27
3	ICC 10755	48.6	474.9	14.8	-1.57	55.4	61.6	5.4	-0.37
4	ICC 16374	37.8	356.6	30.4	-2.53	50.9	63.6	5.0	-0.49
5	IG 7087	49.1	505.8	20.1	-1.65	59.0	61.2	4.8	-0.07
	Mean	46.7	478.6	18.6	-1.89	55.8	71.6	5.2	-0.31

**Table 5.** Time to flowering, shoot and seed yield at maturity and heat tolerance indices of the heat tolerant only at Patancheru cluster group members of chickpea reference collection at Patancheru and Kanpur under heat stress during 2010 summer. Flower, shoot, yield and HTI denotes to days to 50% flowering, shoot biomass g m<sup>2</sup>, seed yield g m<sup>2</sup> and heat tolerance index, respectively and the characters P and K stand for Patancheru and Kanpur, respectively.

S.No	Accession	P-Flower	P-shoot	P-Yield	P-HTI	K-Flower	K-shoot	K-Yield	K-HTI
1	ICC 67	48.2	547.6	202.3	1.07	53.2	86.5	12.8	0.12
2	ICC 283	44.1	482.2	217.7	0.94	55.9	82.0	9.7	-0.09
3	ICC 506	46.1	448.8	183.5	0.77	55.1	75.9	10.8	0.07
4	ICC 708	50.7	513.1	186.6	1.26	55.1	86.8	9.0	-0.44
5	ICC 1164	51.6	470.8	164.8	0.99	60.7	72.2	8.2	-0.33
6	ICC 1356	44.9	464.5	203.1	0.75	52.6	90.8	15.7	0.22
7	ICC 2072	49.5	466.5	153.1	0.80	54.6	79.2	15.4	0.07
8	ICC 2263	52.4	496.1	176.7	0.68	58.2	82.8	10.9	0.11
9	ICC 2629	57.4	460.3	93.7	0.81	59.6	88.2	10.8	0.04
10	ICC 2969	50.3	487.3	219.7	1.57	57.9	120.8	19.2	1.20
11	ICC 3325	44.1	480.2	206.8	0.67	54.3	97.5	16.0	0.17
12	ICC 4657	57.0	457.6	127.3	1.07	56.5	75.9	9.4	-0.12
13	ICC 5434	48.6	404.3	162.8	0.89	55.7	41.4	5.9	-0.05
14	ICC 5613	44.5	441.5	180.6	0.92	54.6	68.1	12.2	0.25
15	ICC 5878	45.3	454.9	199.1	1.24	57.9	54.5	6.7	-0.04
16	ICC 6816	43.2	489.6	201.8	0.65	56.2	71.2	15.4	0.26
17	ICC 8318	40.7	434.6	198.7	0.70	51.3	73.3	14.0	0.12
18	ICC 8522	68.2	420.0	8.0	0.86	51.8	56.1	9.7	-0.05
19	ICC 10018	45.3	447.6	205.7	1.19	56.3	56.6	6.9	-0.18
20	ICC 10393	42.8	442.3	195.6	0.84	54.5	79.2	15.3	0.22
21	ICC 10945	46.6	475.6	191.7	0.96	55.1	79.9	14.4	0.41
22	ICC 11279	50.3	442.2	176.3	2.74	54.3	77.3	10.9	-0.05
23	ICC 12492	54.5	521.2	116.0	0.82	54.8	53.2	9.4	-0.08
24	ICC 12654	51.6	488.8	168.4	0.94	49.8	64.0	12.7	0.11
25	ICC 13124	44.1	497.5	265.4	1.52	52.9	64.7	9.2	-0.25
26	ICC 13892	49.1	416.6	157.1	0.73	52.3	72.0	14.5	0.33
27	ICC 14595	44.9	460.2	213.2	1.17	52.6	72.5	12.3	-0.30
28	ICC 14799	44.5	486.6	195.6	1.06	58.5	101.6	12.9	0.21
29	ICC 15612	49.9	436.4	183.6	0.89	53.7	82.9	13.8	0.20
30	ICC 15614	47.4	464.6	220.9	1.42	54.0	78.8	12.8	0.25
31	ICC 15868	49.9	476.0	164.0	0.96	54.6	89.9	18.1	0.47
32	ICC 16915	44.1	486.2	209.8	1.04	53.2	79.3	16.5	0.40
33	IG 5909	52.0	596.5	147.1	1.08	55.9	62.0	5.1	-0.35
34	IG 6154	73.2	599.7	20.6	1.30	63.2	56.6	5.9	0.21
	Mean	49.3	475.2	174.0	1.04	55.2	75.7	11.8	0.09

**Table 6**

Time to flowering, shoot and seed yield at maturity and heat tolerance indices of the heat tolerant only at Kanpur cluster group members of chickpea reference collection at Patancheru and Kanpur under heat stress during 2010 summer. Flower, shoot, yield and HTI denotes to days to 50% flowering, shoot biomass g m<sup>2</sup>, seed yield g m<sup>2</sup> and heat tolerance index, respectively and the characters P and K stand for Patancheru and Kanpur, respectively.

S.No	Accession	P-Flower	P-shoot	P-Yield	P-HTI	K-Flower	K-shoot	K-Yield	K-HTI
Heat tolerant only at Kanpur									
1	ICC 1083	41.6	405.1	174.0	0.24	54.0	92.0	20.3	0.82
2	ICC 1882	42.8	439.3	187.8	0.44	54.1	100.4	18.8	0.46
3	ICC 2507	47.8	526.1	89.9	-0.42	52.1	84.2	19.1	0.68
4	ICC 2884	50.7	429.2	88.5	-0.04	51.8	82.5	18.3	0.50
5	ICC 3631	47.0	466.4	61.3	-1.00	52.9	100.3	19.5	0.69
6	ICC 4182	49.5	451.2	67.1	-0.50	52.1	90.7	22.7	0.77
7	ICC 4363	42.4	462.5	128.6	-0.07	52.3	75.9	17.6	0.69
8	ICC 4418	49.9	508.5	115.8	-0.12	52.9	75.3	20.9	0.82
9	ICC 4814	47.8	479.1	109.1	-0.51	51.5	105.5	30.6	1.35
10	ICC 5383	45.3	477.0	157.7	0.23	54.6	109.0	21.9	0.39
11	ICC 6293	57.0	444.8	28.2	-0.69	52.4	110.1	19.0	0.74
12	ICC 6537	52.8	480.2	127.3	0.21	58.4	81.1	17.3	0.64
13	ICC 6579	51.1	400.2	114.8	0.00	54.0	79.8	16.6	0.50
14	ICC 9002	49.1	443.2	147.6	0.08	56.2	91.9	20.0	0.79
15	ICC 9895	50.7	465.0	112.4	0.11	55.4	67.9	16.0	0.59
16	ICC 11121	51.1	396.9	104.1	-0.75	54.6	98.3	19.5	0.56
17	ICC 11198	52.4	436.0	122.1	0.36	57.3	109.8	16.5	0.41
18	ICC 12028	55.3	546.7	60.4	-0.08	54.6	82.1	15.7	0.67
19	ICC 13524	54.5	506.4	59.6	-0.72	51.8	71.5	17.8	0.61
20	ICC 14669	42.0	425.8	176.7	-0.17	53.2	102.9	30.6	1.65
21	ICC 14831	45.3	601.4	163.1	0.22	54.6	99.6	22.4	0.92
22	ICC 15510	52.0	457.4	96.1	-0.59	54.3	66.8	16.7	0.69
23	ICC 15606	43.2	499.3	196.5	0.21	56.2	94.9	15.4	0.47
	Mean	48.7	467.3	116.9	-0.15	54.0	90.1	19.7	0.71






Fig 1. Daily maximum and minimum temperatures ( $^{\circ}\text{C}$ ) during the late sown crop growing period both at Patancheru and Kanpur in 2010. The 0 day or the sowing date was 2 Feb 2010 at Patancheru and 13 Jan 2010 at Kanpur.




Fig 2. Relationship of pod number per plant with the HTI and the harvest index (%) with the HTI both at Patancheru and Kanpur.