



Development of composite collection and genotyping of foxtail millet [*Setaria italica* (L.) Beauv.] composite collection



Project PI: HD Upadhyaya, ICRISAT

Project Team: ICRISAT – RK Varshney, CT Hash, DA Hoisington, CLL Gowda, V Gopal Reddy, S Chandra, N Lalitha and A Bharathi



Diversity in foxtail millet germplasm.



Regeneration of foxtail millet germplasm.

Introduction

- Foxtail millet belongs to family *Poaceae* and subfamily *Panicoideae*
- A self-pollinating crop with chromosome number $2n=18$
- Domesticated in highlands of central China, possibly about 4000 BC
- Cultivated in 26 countries, and ranks second in total world production of millets
- Produces six million tons of food mainly in southern Europe and in temperate, subtropical, and tropical Asia (Marathee 1993)
- An important crop in China, India, CIS countries, and Syria
- Possesses high nutrient quality.



Nutritional value of foxtail millet grain (100 g⁻¹)

- Protein content: 10% to 12%
- Lysine content: 2.29% to 2.7%
- Fat content: 4% to 5%
- Energy: 351 kcal
- Thiamin: 0.59 (mg)

Classification of foxtail millet germplasm

Two *Setaria* species are known:

1. *S. glauca* and *S. italica* (Sub sp: *viridis* and *italica*)
2. *S. italica* consists of three races: Moharia, Maxima and Indica; and ten subraces: Aristata, Fusiformis, Glabra, Compacta, Spongiosa, Assamense, Erecta, Nana, Glabra and Profusa.



Status of foxtail millet germplasm at ICRISAT

- The entire foxtail millet germplasm (1535 accessions) characterized for important morpho-agronomic characters following descriptors for *S. italica* and *S. pumila* (IBPGR 1985)
- Collection consists of 1470 landraces, 11 improved cultivars and 54 wild accessions
- A core collection (155 accessions), representing entire diversity, was developed based on geographic origin and quantitative traits.

Genebank/Country	Number of accessions
ICRISAT, India	1535
AICSMIP, UAS, Bangalore, India	1300
National Genebank, China	26222
NIR, Russia	4745
ORSTOM-Montpellier, France	3500
Others	17487
Total	54789

Composite collection

- A composite collection of foxtail millet germplasm (500 accessions) (Table 2) constituted based on geographic origin and diversity for morpho-agronomic traits
- Composite collection consists of accessions representing all three races and ten subraces.

Table 2. Accessions in foxtail millet composite collection.

Trait	Number of accessions
1000 grain weight > 3.7 g	9
Grain weight per plant > 19 g	5
Basal tillers > 29	25
Core collection	155
Dwarf (<= 50 cm)	21
Early flowering accessions (< 40 days)	77
Improved cultivars	10
Short inflorescence length (<=50 mm)	40
Long Inflorescence length (>299 mm)	18
Short inflorescence width accessions (=5 mm)	23
Widest inflorescence accessions (>39 mm)	25
Mono culm types (single tiller per plant)	59
Agronomically elite accessions	33
Total	500

Genotyping

Leaf sample of 20-day old seedling of single representative plant from each of 500 accessions taken and DNA extracted by high-throughput procedure.

Primer optimization

- No SSR markers are available in foxtail millet
- SSR markers from closely related species identified
- Thirty-one unlabelled pearl millet SSR markers selected to genotype 8 diverse foxtail millet accessions (representing five countries)
- Primer optimization with Taguchi method (Taguchi et al. 1986) as described in Cobb and Clarkson (1994)
- Twelve markers showed polymorphism
- Additional SSR markers will be selected from other closely related species such as maize and sorghum.

Future plan

- Genotyping 500 accessions of composite collection using 20 SSR markers
- Identifying a reference collection consisting of most diverse accessions for use in crop improvement program.

References

- Cobb BD and Clarkson JM.** 1994. A simple procedure for optimizing polymerase chain reaction (PCR) using modified Taguchi methods. *Nucleic Acid Research* 22: 3801-3805.
- IBPGR.** 1985. Descriptors for *Setaria italica* and *S. pumila*. Rome, Italy: IBPGR Secretariat.
- Marathee JP.** 1993. Structure and characteristics of the world millet economy. Pages 159–178 in *Advances in small millets* (Riley KW, Gupta SC, Seetharam A and Mushonga JN, eds.). 66 Janapath, New Delhi, India: Oxford and IBH Publ. Co. Pvt.
- Taguchi G.** 1986. Introduction to quality engineering. Asian productivity organization. Dearborn, MI, USA: American Supplies Institute INC.