

1 Introgression of *Botrytis* grey mould 2 resistance genes from *Cicer reticulatum* 3 (*bgmr1_{cr}*) and *C. echinospermum* (*bgmr1_{ce}*) 4 to chickpea (*C. arietinum*)

5 D. Ramgopal¹, R. K. Srivastava², S. Pande², A. Rathore², D. R. Jadhav²,
6 M. Sharma², P. M. Gaur² and N. Mallikarjuna^{2*}

7 ¹Department of Genetics & Plant Breeding, College of Agriculture, ANGR Agricultural
8 University, Rajendranagar, Hyderabad 500 030, India and ²International Crops Research
9 Institute for Semi Arid Tropics (ICRISAT), Patancheru P.O., Andhra Pradesh 502 324, India

10 Received 25 October 2012; Accepted 22 November 2012

11 Abstract

12 *Botrytis* grey mould (BGM), caused by the fungus *Botrytis cinerea* Pers. ex Fr., is an important
13 disease of chickpea causing economic losses across the world in chickpea-growing
14 regions. There are no available resistance sources in cultivated chickpea against this disease.
15 *Cicer echinospermum* and *C. reticulatum*, the only two compatible annual wild species,
16 have been reported to have resistance to BGM. Hence, interspecific populations were devel-
17 oped with susceptible cultivars as female parents and *C. echinospermum* accession IG
18 73074 and *C. reticulatum* accession IG 72937 as the pollen donors to transfer and assess
19 the nature of genetic control for BGM. Screening the progeny indicated that resistance to
20 BGM was controlled by a single additive gene/allele (*bgmr1_{cr}* and *bgmr1_{ce}*), which can be
21 introgressed through a backcross breeding programme.

22 **Keywords:** *Botrytis*-grey mould; chickpea; *Cicer arietinum*; *Cicer echinospermum*;
23 *Cicer reticulatum*; disease resistance; interspecific cross

24 Introduction

25 *Botrytis* grey mould (BGM), a disease caused by
26 the necrotrophic fungus *Botrytis cinerea* Pers. ex Fr.,
27 has been reported from more than 15 countries (Nene
28 *et al.*, 1984). BGM is one of the most devastating diseases
29 of chickpea (*Cicer arietinum* L.) and can result in com-
30 plete yield loss (Davidson *et al.*, 2004). It was first
31 reported in the Jujuy Province of Argentina, causing
32 95% crop loss (Carranza, 1965). It is a serious constraint
33 to chickpea production in many Asian countries includ-
34 ing northern India, Nepal, Bangladesh and Pakistan.
35 The disease is considered to be the major cause for

the decline in the chickpea-growing areas of Nepal and 36
Bangladesh (Pande, 1998). More than 80% yield loss has 37
been observed in chickpea crops grown on the 38
Indo-Gangetic plains of India (Pande, 1998). High levels 39
of resistance have not been found in the cultivated germ- 40
plasm (Singh and Bhan, 1986), which has encouraged the 41
search for resistance sources in the related wild species. 42

Evaluation of germplasm accessions of wild species 43
has revealed that they possess a wealth of genes for 44
biotic and abiotic stresses (Haware *et al.*, 1992, Mallikar- 45
juna, 2003; Gaur *et al.*, 2009). They have resistance to 46
three or more stresses such as *Ascochyta* blight, BGM 47
and *Fusarium* wilt (Robertson *et al.*, 1995). *Cicer reticu-* 48
latum and *C. echinospermum*, two wild relatives from 49
the secondary gene pool of chickpea (Mallikarjuna 50
et al., 2011), have been reported to be resistant to 51
BGM (Singh *et al.*, 1991; Singh *et al.*, 1998; Ramgopal, 52

* Corresponding author. E-mail: N.Mallikarjuna@cgiar.org

2006). In addition, accessions of *Cicer bijugum*, *C. pinnatifidum* and *C. judaicum* from the tertiary gene pool are resistant, but these species are currently inaccessible for chickpea improvement due to incompatibility between these species and cultivated chickpea (Mallikarjuna *et al.*, 2011). Wild relatives in the secondary gene pool of chickpea are amenable to wide crossing and gene transfer (Collard *et al.*, 2003; Mallikarjuna *et al.*, 2011). Nevertheless, until now, none of the wild *Cicer* species have been used in the crossing programme to transfer BGM resistance to cultivated chickpea.

Currently, deployment of host plant resistance has limited potential in BGM management, as high levels of resistance have not been identified in cultivated germplasm and because of the variable nature of *B. cinerea* populations (Davidson *et al.*, 2004). Fungicidal control of BGM is expensive and development of fungicide resistance has been reported frequently in *B. cinerea* populations (Leroux, 2004). This study describes the introgression of BGM resistance from *C. reticulatum* and *C. echinospermum* into chickpea and the pattern of inheritance of resistance.

Materials and methods

Plant material

The experiment was conducted at the International Crops Research Institute for Semi Arid Tropics (ICRISAT), Patancheru, Andhra Pradesh, India. The Genetic Resources Unit of ICRISAT provided *C. reticulatum* (collected from Turkey) and *C. echinospermum* (collected from Turkey) seeds, which were multiplied and used in the crossing programme. The crossing programme was carried out using the BGM-resistant accessions of *C. echinospermum* IG 73 074 (ICC 20 192) and *C. reticulatum* IG 72 937 (ICC 20 170) as male parents. Chickpea cultivars ICC 4954 and ICC 92 318, both susceptible to BGM, were used as female parents to develop interspecific populations. The F₁s developed were selfed to develop F₂ and backcrossed to the cultivated chickpea parent to develop BC₁ populations in the glasshouse.

Screening for BGM

For the identification of BGM resistance, 8–10-d-old test seedlings along with the susceptible parent chickpea cultivars ICC 92 318 and ICC 4954 were inoculated with *B. cinerea* on a potato dextrose agar medium, which was isolated from naturally infected chickpea plants collected from the BGM hot-spot location Pantnagar, India. The isolate was from single spore following

standard mycological procedures. Conidia of *B. cinerea* were cultured on autoclaved marigold (*Tagetes erecta*) flowers. The conidia were harvested into sterile distilled water, adjusted to 3×10^5 conidia/ml using a haemocytometer and used as an inoculum. Chickpea seedlings were sprayed with the inoculum using a hand-operated atomizer. The inoculum was allowed to partially dry for about 30 min. Inoculated plants were maintained at $15 \pm 2^\circ\text{C}$ and above 60% relative humidity with a 12 h photoperiod. Disease scores for BGM on each accession were recorded using a 1–9 rating scale at 20 d after inoculation. The disease rating from scores 1 to 3 was treated as resistant, scores 4 to 5 as moderately resistant and scores 6 to 9 as susceptible (Pande *et al.*, 2006).

Data analysis

A χ^2 goodness-of-fit test was calculated as given by Panse and Sukhatme (1967) and the calculated χ^2 values were compared with table values given by Fisher and Yates (1963), against appropriate degrees of freedom (df).

Results

In this study, two crosses, one derived from *C. reticulatum* (ICC 92 318 \times IG 72 937) and another derived from *C. echinospermum* (ICC 4954 \times IG 73 074), were developed to screen for BGM resistance. The disease symptoms in the susceptible cultivar began with wilting and decaying of the leaves followed by decaying of the aerial parts of the plant by 18–20 d post-inoculation (Fig. 1(b)). In the resistant *C. echinospermum* (Fig. 1(a)) and the resistant interspecific derivatives (Fig. 1(c)), the aerial parts remained green and fresh without any signs of wilting and decaying. Wilting and decaying of the aerial parts of the seedlings was observed in disease-susceptible interspecific derivatives (Fig. 1(d)). Plants that did not show any disease symptoms upon transplantation to suitable pots grew further and set flowers and pods. Susceptible plants continued to decay under disease pressure.

The F₂ population derived from the *C. reticulatum* IG 72 937 cross (ICC 92 318 \times IG 72 937) had a total of 16 plants. BGM disease screening tests showed three plants to be resistant while nine plants were found to be moderately resistant and four were susceptible to the disease. The BGM disease reaction fitted into a 1 (resistant):2: (moderately resistant):1 (susceptible) monogenic segregation ratio with additive gene action (χ^2 test ratio 0.38^{ns}, $P = 0.83$ at 2 df). Similarly, the BC₁ population (derived by crossing the F₁ of the same cross to ICC 92 318) had a total of 20 plants in which eight plants were moderately resistant to the disease and 12 plants were susceptible.




Fig. 1. Botrytis grey mould screening of interspecific derivatives between *C. arietinum* and *C. echinospermum*. (a) Wild species, *C. echinospermum*, showing no BGM disease. (b) Cultivated chickpea cultivar showing susceptibility to the disease. (c) Interspecific derivative resistant to the disease. (d) Interspecific derivative showing susceptible reaction to the disease (a colour version of this figure can be found online at journals.cambridge.org/pgr).

149 The χ^2 test (0.8^{ns}) with a P value of 0.37 at 1 df
 150 followed the ratio 1:1, thereby confirming the monogenic
 151 additive F_2 segregation ratio. The progeny derived from
 152 *C. reticulatum* were advanced further and screened for
 153 BGM, and the lines showed resistance to BGM.

154 The F_2 population derived from the *C. echinospermum*
 155 IG 73074 cross (ICC 4954 \times IG 73074) had a total of
 156 63 plants, of which 12 plants were resistant, 37 moder-
 157 ately resistant and 14 susceptible to the disease. The χ^2
 158 test was carried out to test whether the ratio of 1:2:1
 159 fitted. The χ^2 test was 2.05^{ns} with a P value of 0.36 at
 160 2 df. The BC_1 population (obtained by crossing the F_1
 161 of the same cross to ICC 4954) had a total of 17 plants
 162 with seven moderately resistant plants and 10 susceptible
 163 plants. The χ^2 test (0.53 , n.s.) with a P value of 0.47 at

1 df confirmed the ratio of 1:1 for the backcross. These
 results indicate that BGM resistance inherited from both
C. reticulatum and *C. echinospermum* was a single
 gene showing additivity.

Discussion

This is the first report on the introgression of BGM
 resistance from wild relatives of *Cicer*, namely
C. reticulatum and *C. echinospermum*, into *C. arietinum*.
 In the present study, single and additive modes of
 resistance were observed for BGM. The additive alleles
 from *C. reticulatum* (designated as *bgmr1_{cr}*) and *C. echi-*
nospermum (designated as *bgmr1_{ce}*) were needed in two

164
 165
 166
 167

168

169
 170
 171
 172
 173
 174
 175

doses (*bgmr1_{cr}/bgmr1_{cr}*; *bgmr1_{ce}/bgmr1_{ce}*) to impart complete resistance, while heterozygous (*Bgmr1_{cr}/bgmr1_{cr}*; *Bgmr1_{ce}/bgmr1_{ce}*) individuals were moderately resistant. Disease resistance genes present in the wild species are recessive in many crop plants, as seen in *Cajanus platycarpus*, where resistance to *Phytophthora* blight has been reported to be monogenic and recessive (Mallikarjuna et al., 2005). Collard et al. (2003) reported digenic and recessive modes of resistance to Ascochyta blight in the interspecific derivatives of chickpea from *C. echinospermum* accession ICC 17 159. The results from the present study indicated that neither susceptibility nor resistance was dominant over the other. Previous reports on the transfer of BGM resistance from cultivated chickpea lines showed single dominant gene, and between two resistant cultivated chickpea parents showed duplicate dominant genes (Singh, 1997). The nature of resistance was probably moderate to low as the plants succumbed to the disease when the infection was moderate to severe. Some more examples of *Cicer* species contributions to chickpea improvement are successful introgression of *Phytophthora* root rot resistance from *C. echinospermum* (Knights et al., 2008) and introgression of nematode resistance from *C. reticulatum* and *C. echinospermum* (Gaur et al., 2009). *Cicer pinnatifidum*, *C. judaicum* and *C. bijugum* are known to possess resistance to *Fusarium* wilt, Ascochyta blight, BGM and bruchids (Stevenson and Veitch, 1998; Mallikarjuna et al., 2011).

These results indicate that when desired levels of resistance to biotic constraints are lacking in the cultivated or primary gene pool, there is an option for sources of resistance in the secondary gene pool where the species are cross-compatible and offer genetic variability to tackle many of the biotic constraints (van der Maesen et al., 2007; Mallikarjuna et al., 2011). It was possible to transfer *Helicoverpa armigera* resistance from *C. reticulatum* (Mallikarjuna et al., 2007; Mallikarjuna et al., 2011). Therefore, as demonstrated in the present study, *C. reticulatum* and *C. echinospermum* with their valuable sources of variation for BGM resistance offer genetic variability to broaden the genetic base of cultivated chickpea and introduce useful traits not present in the cultivated gene pool.

Further studies on allelism are needed to ascertain whether the two BGM resistance genes (*bgmr1_{cr}* and *bgmr1_{ce}*) reported in this study are allelic variants of the same gene, or are two different genes.

References

Carranza JM (1965) Wilt of chickpea (*C. arietinum*) caused by *B. cinerea* (in Spanish). *Revista de la Facultad de Agronomia. Universidad Nacional de la Plata* 41: 135–138.

Collard BCY, Pang ECK, Ades PK and Taylor PWJ (2003) Preliminary investigation of QTLs associated with seedling resistance to ascochyta blight from *Cicer echinospermum*, a wild relative of chickpea. *Theoretical Applied Genetics* 107: 719–729.

Davidson JA, Pande S, Bretag TW, Lindbeck KD, Kishore GK. (2004) Biology and management of Botrytis spp. in legume crops. In: Elad Y, Williamson B, Tudzynski P, Delen N (eds) *Botrytis: Biology, Pathology and Control*. The Netherlands: Kluwer Academic Publishers, pp. 295–318.

Fisher RA and Yates F (1963) *Statistical Tables for Biological, Agricultural and Medical Research*, 6th edn. London: Oliver and Boyd.

Gaur PM, Mallikarjuna N, Knights T, Beebe S, Debouck D, Mejia A, Malhotra RS, Imtiaz M, Sarker A, Tripathi S, Gowda CLL (2009) Gene introgression in grain legumes. In: *International Conference on Grain Legumes: Quality Improvement, Value Addition and Trade*. Indian Society of Pulses Research and Development, Indian Institute of Pulses Research, Kanpur, 14–16 February, 2009.

Haware MP, Nene YL, Pundir RPS and Narayana Rao J (1992) Screening the world chickpea resistance to *Fusarium* wilt. *Field Crops Research* 30: 147–154.

Knights EJ, Southwell RJ, Schwinghamer MW and Harden S (2008) Resistance to *Phytophthora medicaginis* Hansen and Maxwell in wild *Cicer* species and its use in breeding root rot resistant chickpea (*Cicer arietinum* L.). *Australian Journal of Agricultural Research* 59: 383–387.

Leroux P (2004) Chemical control of Botrytis and its resistance to chemical fungicides. In: Elad Y, Williamson B, Tudzynski P, Delen N (eds) *Botrytis: Biology, Pathology and Control*. Dordrecht: Kluwer Academic Publishers, pp. 195–222.

Mallikarjuna N (2003) Wide hybridization in important food legumes. In: Jaiwal PK, Singh RP (eds) *Improvement Strategies of Leguminosae Biotechnology*. Dordrecht: Kluwer Academic Publishers, pp. 155–170.

Mallikarjuna N, Jadhav Deepak and Reddy MV (2005) Introgression of phytophthora blight disease resistance from *Cajanus platycarpus* into short duration pigeonpea (*Cajanus cajanus* (L.). Millisp.). *Indian Journal of Genetics and Plant Breeding* 55: 261–263.

Mallikarjuna N, Coyne C, Cho S, Rynearson S, Rajesh PN, Jadhav DR, Muehlbauer FJ (2011) Chickpea. In: Kole C (ed) *Wild Crop Relatives: Genomic and Breeding Resources*. Berlin, Heidelberg: Springer, pp. 63–82. doi: 10.1007/978-3-642-14387-8_4.

Mallikarjuna N, Sharma HC and Upadhyaya HD (2007) Exploitation of wild relatives of pigeonpea and chickpea for resistance to *Helicoverpa armigera*. *E-journal of SAT Agricultural Research under Crop Improvement* 3: 4–7.

Nene YL, Sheila VK, Sharma SB. (1984) *A World List of Chickpea (Cicer arietinum) and Pigeonpea (Cajanus cajan L. Millsp.) Pathogens*. ICRISAT Pulse Pathology Report 32. Patancheru: International Crops Research Institute for Semi Arid Tropics.

Pande S (1998) *Diseases of Chickpea in Nepal and Bangladesh. A Survey of Report. Trip Report January 1998*. Patancheru: International Crops Research Institute for Semi Arid Tropics.

Pande S, Ramgopal D, Kishore GK, Mallikarjuna N, Sharma M, Pathak M and NarayanRao J (2006) Evaluation of wild *Cicer* species for resistance to ascochyta blight and Botrytis gray mold in controlled environment at ICRISAT,

Introgression of Botrytis grey mould resistance genes

- Patancheru. *International Chickpea and Pigeonpea Newsletter* 13: 25–26.
- Panse VG and Sukhatme PV (1967) *Statistical Methods for Agricultural Workers*. New Delhi: Indian Council of Agricultural Research.
- Ramgopal D (2006) Characterization and evaluation of annual wild *Cicer* species and study the transfer of Botrytis gray mold and ascochyta blight resistance from *Cicer echinopspermum* into cultivated species. PhD Thesis, Acharya N.G. Ranga Agricultural University, Hyderabad.
- Robertson LD, Singh KB and Ocampo B (1995) *A Catalogue of Annual Wild Annual Cicer Species*. Aleppo: ICARDA, Syria Publication.
- Singh IS (1997) Genetics of resistance to botrytis gray mould of chickpea. In: *Summary Proceeding of the 3rd Working Group Meeting to Discuss Collaborative Research on Botrytis Gray Mold of Chickpea*, pp. 31–32.
- Singh G and Bhan LK (1986) Physiological races of *Botrytis cenerea* causing gray mold of chickpea. *Plant Disease Research* 1: 69–74.
- Singh G, Kaur L and Sharma YR (1991) Ascochyta blight and gray mold resistance in wild species of *Cicer*. *Crop Improvement* 18: 150–151.
- Singh KB, Ocampo B and Robertson LD (1998) Diversity for abiotic and biotic resistance in the wild annual *Cicer* species. *Genetic Resources and Crop Evolution* 45: 9–17.
- Stevenson PC, Veitch NC and Nigel C (1998) A 2-acylbenzofuran from roots of *Cicer bijugum* associated with *Fusarium* wilt resistance. *Phytochemistry* 48: 947–951.
- van der Maesen LJG, Maxted N, Javadi F, Coles S, Davies AMR (2007) Taxonomy of the genus *Cicer* revisited. In: Yadav SS, Redden RJ, Chen W, Sharma B (eds) *Chickpea Breeding and Management*. Wallington: CABI, pp. 14–47.

Author Queries

JOB NUMBER: 97-12

JOURNAL: PGR

- Q1** A short title has been inserted. Please approve or provide an alternative.
- Q2** Please check edit of the following sentence: 'For the identification...'
- Q3** Please check whether the expansion of RH (relative humidity) is correct.
- Q4** Please clarify as to what the superscript 'ns' refer to.
- Q5** Please check the insertion of location details in references Leroux (2004) and Mallikarjuna (2003).
- Q6** Please provide publisher and location details for Singh (1997).