



Soil Testing as a Tool for On-Farm Fertility Management: Experience from the Semi-arid Zone of India

Kanwar L. Sahrawat^{a*} & Suhas P. Wani^a

^a International Crops Research Institute for the Semi-arid Tropics (ICRISAT), Andhra Pradesh, India

Communications in Soil Science and Plant Analysis

Volume 44, Issue 6, 2013. Pages 1011-1032

<http://dx.doi.org/10.1080/00103624.2012.750339>

This is author version post print archived in the official Institutional Repository of ICRISAT www.icrisat.org

Soil Testing as a Tool for On-Farm Fertility Management: Experience from the Semi-Arid Zone of India

KANWAR L. SAHRAWAT AND SUHAS P. WANI

International Crops Research Institute for the Semi-Arid Tropics (ICRISAT)

Patancheru 502 324, Andhra Pradesh, India

Rainfed agriculture in the dry regions is affected by water shortages. Our earlier research showed that not only the deficiencies of major nutrients, but those of sulfur (S) and micronutrients are holding back the potential of agricultural production systems. The objectives of this article are to discuss the efficacy of soil testing to diagnose nutrient

deficiencies using 28 270 diverse soil samples collected from farmers' fields in the semi-arid tropical (SAT) regions of India, and to confirm the efficacy of the soil test-based balanced nutrient management in enhancing productivity of a range of crops in on-farm farmer participatory trials under rainfed conditions. Results of a large numbers of on-farm trials demonstrated that soil testing is indeed an effective tool for on-farm fertility management, a prerequisite, for sustainably enhancing the productivity in rainfed areas in the SAT regions of India. The need to strengthen the soil testing infrastructure in the country is emphasized.

Keywords Balanced nutrient management, crop productivity and quality, diagnosis of nutrient problems, soil quality, soil test-based recommendation, water shortage, water use efficiency

Address correspondence to K. L. Sahrawat, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru 502 324, Andhra Pradesh, India. Email: k.sahrawat@cgiar.org

Introduction

In the rainfed production systems, the importance of water shortage and associated stress cannot be overemphasized, especially in the semi-arid tropical (SAT) regions (Pathak et al. 2009; Passioura and Angus 2010; Rockström et al. 2010; Sahrawat et al. 2010a; Sharma et al. 2010). However, apart from water shortage soil infertility is also the issue for crop production and productivity enhancement in much of the SAT regions of the world and Indian SAT is no exception (El-Swaify et al. 1985; Black 1993; Zougmoré et al. 2003; Sahrawat et al. 2007, 2010b; Singh 2008; Bationo et al. 2008; Twomlow et al. 2008a; Bekunda et al. 2010).

Apart from the deficiencies of major nutrients, nitrogen (N) and phosphorus (P), the deficiencies of secondary nutrients especially of sulfur (S) and micronutrients have been reported with increasing frequencies from the intensified irrigated production

systems (Kanwar 1972; Pasricha and Fox 1993; Takkar 1996; Scherer 2001, 2009; Fageria et al. 2002; Singh 2008). While in the irrigated systems the deficiencies of various plant nutrients have been diagnosed through soil and plant testing and managed through the fertilization of crops, little attention seems to have been paid to diagnosing the deficiencies of secondary nutrients such as S and micronutrients in dryland rainfed production systems especially in SAT India (Sahrawat et al. 2007; Sahrawat et al. 2010b).

Specifically, little attention has been devoted to survey and determine the fertility status of farmers' fields with an objective to diagnose the nutrient problems in the rainfed production systems, which is a prerequisite for developing an effective nutrient management strategy for enhancing agricultural productivity in these areas. Lack of adequate analytical laboratory support infrastructure in developing countries coupled with the lack of awareness that the mining of secondary and micronutrients in production systems is not helping the cause of upgrading the rainfed agriculture. The information on the soil fertility status is needed not only for enhancing crop productivity through balanced nutrient management, but also to promote judicious use of costly external inputs of nutrients and enhancing the efficiency of scarce water resources in developing countries like India (Sahrawat 2006; Wani 2008).

This apparent paradox of lack of application of adequate amount of nutrients from external inputs (Bationo et al. 2008; Katyál 2003) despite the common knowledge that the soil resource base in the rainfed systems of the SAT regions is relatively fragile and marginal compared to that under the irrigated production systems (El-Swaify et al. 1985; Rego et al. 2003; Sahrawat et al. 2007, 2010b) is rather inexplicable.

In the rainfed systems of India, the management of water shortage has been the primary focus of research and developmental activities in these areas, and soil infertility has largely been ignored (El-Swaify et al. 1985; Wani et al. 2003; Sahrawat et al. 2010a, b) or has not been addressed in an integrated manner along with soil and water conservation practices (Wani et al. 2009; Rockström et al. 2010).

However, even in water-limiting environments there is potential to enhance agricultural productivity through efficient management of soil, water and nutrients in an integrated manner (Wani et al. 2002; Twomlow et al. 2008a; Wani et al. 2009, Sahrawat 2010b). To achieve the potential of productivity in water-limited environments, a concept

of water-limited potential yield seems very appropriate as this forms the basis to reach the attainable yield in these environments through management of various constraints other than just water shortage (Passioura 2006; Singh et al. 2009). For example, in Australia, farmers have adopted the notion of water-limited potential yield as a benchmark for yield and if farmers find that their crops are performing below the benchmark, they look for the reasons and attempt to improve their management accordingly (Passioura and Angus 2010). We emphasize that in the concept of water-limited potential yield in the rainfed systems, natural resource management in general and soil fertility management in particular need to be paid due attention along side water stress management in view of the fragile nature of the soil resource base (Sahrawat 2010a, b; Wani et al. 2009).

Moreover, it is a commonly held belief that at relatively low yields of crops in the rainfed systems of India, the deficiencies of major nutrients, especially those of N and P only are important for the SAT Indian soils (El-Swaify et al., 1985; Rego et al. 2003) and consequently little attention has been devoted to diagnose the extent of deficiencies of the secondary nutrients such as S and micronutrients in various crop production systems on millions of small and marginal farmers' fields (Rego et al. 2005, 2007; Sahrawat et al. 2007, 2010b).

It is recognized and duly emphasized that the productivity of the SAT soils is low due to water shortages. Although, low fertility is also an issue, in practice the deficiencies of major nutrients (N and P) are considered important and the role of secondary and micronutrients in enhancing water use efficiency is neglected. Moreover, the input of major nutrients to dryland production systems is meager compared to that in the irrigated systems (Rego et al. 2005; Wani et al. 2009). Also, due to low productivity of the rainfed crops, it is generally assumed that the off take and mining of micronutrients reserves in soils is much less than in irrigated production systems (Rego et al., 2003).

For sustained increase in dryland productivity, soil and water conservation measures need to be integrated with plant nutrition, and choice of crops and their management (Wani et al., 2003; Passioura 2006; Passioura and Angus 2010; Sahrawat et al. 2010a). The on-going farmer participatory integrated watershed management program of the ICRISAT (International Crops Research Institute for the Semi-Arid Tropics) provided an appropriate opportunity to implement nutrient management strategy along

side with soil and water conservation practices in farmers' fields in the Indian semi-arid tropics. For achieving efficient and judicious use of nutrients through fertilizer inputs, assessing the soil's inherent nutrient status is a prerequisite (Sahrawat, 2006).

The objectives of this paper therefore, are to review, analyze and present recent results on the general fertility status of soils in the rainfed systems with emphasis on the deficiencies of secondary and micro-nutrients, and to confirm the efficacy of the soil test-based nutrient management strategy to increase the productivity of a range of crops in farmer participatory on-farm trials in the rain-fed systems of the SAT regions. Preference is given to the results generated from the on-farm research in the SAT regions of India. First, the results on the fertility status of SAT soils are dealt, followed by the response of various food crops to balanced nutrient management considering the various nutrient deficiencies under the on-farm conditions. Equally importantly, the role of soil testing in the diagnosis of nutrient deficiencies has been demonstrated and hence emphasized.

Materials and Methods

Diagnosis of Nutrient Deficiencies by Soil Testing

Since 1997, the natural resources management group at the ICRISAT center in India along with its partners has been conducting systematic and detailed studies on the diagnosis and management of nutrient deficiencies in the semi-arid regions of Asia with emphasis on the semi-arid regions of India. It started with detailed analysis of farmers' fields in Milli watershed at Lalatora in Madhya Pradesh where analysis of soil samples for micronutrients was deliberately included as a part of the baseline characterization of the site.

First a soil sampling methodology was developed and standardized to collect representative soil samples in a watershed. The methodology is based on the stratified random sampling of the watershed considering the soil types including topography, major crops, and farmers' land holding size (for details see Sahrawat et al. 2008b). For effective soil sampling, farmers' fields were divided into three groups based on the position on the toposequence: top, middle and bottom, depending on the elevation and drainage pattern. We separated different soil types in each category. For soil sampling, we randomly selected 20% farmers in each position on the toposequence, proportion to the farm size,

types of soils and crops grown (see Sahrawat et al. 2008b). The soil sampling program of watersheds in various states was undertaken largely during 2002-2009.

Using stratified random sampling methodology (Sahrawat et al., 2008b); we collected 8 to 10 cores of surface (0-15 cm depth) soils to make one composite sample. The soil samples were air-dried and powdered with wooden hammer to pass through a 2-mm sieve. For organic carbon (C) analysis, the soil samples were ground to pass through a 0.25-mm sieve. Prepared samples were analyzed for various fertility characteristics in the ICRISAT Central Analytical Services Laboratory.

To characterize the fertility status of soils under dryland agriculture in the SAT regions of India, we collected 28 270 soil samples from farmers' fields, spread in the Indian states of Andhra Pradesh, Karnataka, Rajasthan and Madhya Pradesh. The number of farmers cultivating arable land varied along with land holding size, crops and cropping systems.

For soil analysis, pH was measured by a glass electrode using a soil to water ratio of 1:2. Organic C was determined using the Walkley-Black method (Nelson and Sommers 1996). Exchangeable potassium (K) was determined using the ammonium acetate method (Helmke and Sparks 1996). Available S was measured using 0.15% calcium chloride (CaCl_2) as an extractant (Tabatabai 1996; Sahrawat et al. 2009); available P (Olsen-P) was measured using sodium bicarbonate (NaHCO_3) as an extractant (Olsen and Sommers 1982). Available zinc (Zn) was extracted by DTPA reagent (Lindsay and Norvell 1978) and available boron (B) was extracted by hot water (Keren 1996).

Based on the results of soil samples collected from farmers' fields, recommendations were developed at block level for balanced nutrient management. For this, critical limits in the soil for various plant nutrients were used (Table 1) to separate deficient soil samples from the non-deficient ones (Sahrawat 2006; Rego et al. 2007; Sahrawat et al. 2007) for the follow up on-farm crop response studies.

On-farm Crop Responses to Soil Test-Based Fertilization

During 2002-2009 cropping seasons (June-September), we conducted a large number of trials in farmers' fields in the SAT regions of Andhra Pradesh, Karnataka, Madhya

Pradesh and Rajasthan in India with a range of locally important field crops. Each farmer for a crop was treated as a replication. The details of the on-farm trials along with various treatments and crop and nutrient management practices followed were similar to those described in Rego et al. (2007).

In the on-farm trials, there were two treatments: (i) control or farmers' nutrient inputs (termed FI) and (ii) balanced nutrient management treatment or BN consisting of applications of nutrients found deficient based on soil test results. The BN treatment consisted of applications of S + B + Zn or SBZn along with N and P over the FI treatment [FI + SBZn + N + P]. These two treatments were imposed on 2000 or 1000 m² plots side by side on the same piece of land. Farmers' crops and variety, and crop management practices were the same in both the treatments. Before implementation of the treatments in the conduct of field trials, the soil test results on soil samples collected from farmers' fields were shared and discussed with the participating volunteer farmers in their own languages. As mentioned earlier, each farmer's field was considered as one replication.

For applying nutrients as per SBZn treatment, we applied S, B and Zn via a mixture, which consisted of 200 kg gypsum (30 kg S ha⁻¹), 5 kg borax or 2.5 kg Agribore (0.5 kg B ha⁻¹) and 50 kg zinc sulfate (10 kg Zn ha⁻¹) ha⁻¹; the mixture was surface broadcast on the plot before the final land preparation. The SBZn + NP or BN treatment consisted of the same amount of S, B and Zn as in SBZn plus 60 kg N for cereals or 20 kg N ha⁻¹ for legumes; and P was added at 30 kg P₂O₅ ha⁻¹. The treatment SBZn was applied along with P plus 20 kg N ha⁻¹ as basal to all crops and 40 kg N ha⁻¹ was top dressed in the case of cereals. In the case of NP treatment, we applied 20 kg N and 30 kg P₂O₅ ha⁻¹ to all crops as basal and 40 kg N ha⁻¹ as topdressing for cereals.

Results and Discussion

Organic Carbon and Extractable Nutrient Status of Farmers' Fields

The soil test results for pH, organic C and extractable P, K, S, B and Zn of soil samples collected from farmers' fields in the SAT regions of Indian states of Andhra Pradesh (3650 farmers' fields), Karnataka (22867), Madhya Pradesh (341) and Rajasthan (421) showed that the results varied with district in a state and had a wide range in soil

chemical fertility parameters (Table 2). In these results, soil organic C has been used as an index of available N (Sahrawat et al. 2010b).

These first results on the fertility status of farmers' fields at a large scale showed that the samples were generally low in organic C (we have used soil organic C as a proxy for N supplying capacity of a soil), low to medium in Olsen extractable P, medium to high in exchangeable K and generally low in calcium chloride extractable S, hot water extractable B and DTPA extractable Zn (Table 2). The results clearly demonstrate that soils are not only low in organic C and Olsen-P, but also low in secondary nutrient such as S and micronutrients such as B and Zn. The number of farmers' fields sampled from 14 districts of Karnataka state was fairly large and based on the results of these samples; some plausible conclusions can be drawn for the prevalence of plant nutrient problems in the state, which is the second largest state in India with rainfed agriculture after Rajasthan. The mean organic C content in the soil samples was 0.45%, Olsen-P was deficient in 47% of the 22867 farmers' fields sampled, exchangeable K was deficient only in 16% farmers' fields, extractable S in 83% fields, hot water extractable B in 66% fields and DTPA extractable Zn in 61% of the sampled farmers' fields.

In Andhra Pradesh, B deficiency was most prevalent (in 85% of the 3650 fields sampled), followed by S, which was deficient in 79% of the farmers' fields and Zn was deficient in 69% of the farmers' fields. Olsen-P was deficient in 38% of the fields and K only in 12% of the fields (Table 2). In Madhya Pradesh (341 farmers' fields sampled), B deficiency was most prevalent (79% fields), followed by S (74%), Olsen-P (74%) and Zn (66%). While in Rajasthan (421 fields sampled), the deficiency of S was most widespread (in 71% of the fields), followed by B (56%), Olsen-P (45%), Zn (40%), and K (15%) (Table2).

Considering the results of analyses of all soil samples from the four states in the SAT region of India, it can be concluded that the deficiency of S (calcium chloride extractable) was most widespread (on an average 82% of the 28270 farmers' fields sampled were deficient), followed by hot water extractable B (68% of the farmers' fields sampled were deficient) and DTPA extractable Zn (62% of the farmers' fields were found deficient); and the finding is indeed revealing. These results are in accord with

those reported earlier with a limited number of soil samples (Rego et al. 2005; Sahrawat et al. 2007, 2010b).

Another important finding emerging from the soil test results of the soil samples analyzed is that K deficiency has not emerged as a prominent nutrient deficiency as on average only 16% of the farmers' fields out of a total of 28270 farmers' fields sampled were found to be deficient in the rainfed production systems of the SAT regions (Table 2).

These results are significant in showing the widespread nature of the occurrence of the deficiencies of major nutrients such as N and P, but more importantly those of S, B and Zn in the rainfed production systems of the SAT India. The extent of deficiencies of plant nutrients appear as widespread as those reported from the intensified irrigated systems (Pasricha and Fox 1993; Takkar 1996; Scherer 2001; Fageria et al. 2002; Sahrawat et al. 2010b). To our knowledge, no relatively large scale on-farm survey on the nutrient status of farmers' fields in the SAT regions of India has been undertaken, and thus no benchmark results are available to compare the extent of the deficiencies of S and micronutrients in farmers' fields. But these results do clearly demonstrate that in addition to water stress, multiple-nutrient deficiencies have to be managed to unlock the potential of rainfed production systems. The earlier research on fertility management has mostly concentrated on the major nutrients and the deficiencies of N and P have been reported to be widespread in the rainfed systems (El-Swaify et al. 1985; Sahrawat et al. 2001; Rego et al. 2003; Bationo et al. 2008).

Soil organic matter and major plant nutrients (N, P and K) depletion remains a major constraint to long-term agricultural sustainability in much of the rainfed agricultural systems in the SAT regions of Asia and Sub-Saharan Africa. Negative nutrient balances (nutrient added minus nutrient harvested in crop) relative to mostly major plant nutrients have been reported as the nutrient removal exceeds input over a long period of time with concomitant decline in soil organic matter status. Organic matter depletion problem is particularly acute in the rainfed systems where the external inputs of organic matter and nutrients is far lower than the loss or removal (Katyal 2003; Rego et al 2003; Bationo et al. 2008; Bekunda et al. 2010).

Soil Test-Based Nutrient Management: Effects on Crop Productivity and Quality

As mentioned in the introduction, soil fertility management research in the rainfed areas has focused mainly on the management of major nutrients (N, P and K) and even the amounts of these nutrients is generally inadequate (Rego et al. 2007; Bationo et al. 2008; Sahrawat et al. 2010b). Water stress by erratic and low rainfall is the major bottleneck for farmers to apply adequate amounts of nutrients in the rainfed systems. However, recent work by ICRISAT and its partners and other researchers has shown that for realizing the potential of rainfed systems, both water stress and nutrient deficiencies need to be attended simultaneously (Wani et al. 2003; Ncube et al. 2007; Bationo et al. 2008; Sahrawat et al. 2010a).

Rego et al. (2007) conducted a number of on-farm trials under rainfed conditions for three years (2002-2004) during the rainy season (June-October) in three districts of Andhra Pradesh in the SAT region of India to evaluate crop responses to balanced nutrient management based on soil test results using mung bean, maize, groundnut, castor and pigeonpea. There were two treatments (i) control or farmer's nutrient input (FI) and (ii) balanced nutrient (BN) management, which consisted of the applications of SBZn + NP over FI or FI + SBZn + NP. The grain yields of maize, castor, mung bean, groundnut (pod yield) and pigeonpea crops were significantly increased under BN with the applications application of SBZn + NP over the FI treatment in the three seasons (Table 3).

A large number of on-farm trials were also conducted in the semi-arid zone of Karnataka state during five rainy seasons (2005-2009) with maize, finger millet, groundnut and soybean as the test crops. Again, as in the case of trials in Andhra Pradesh, BN treatment significantly increased the grain yields of these crops over the farmer's inputs treatment (Table 4). In another set of trials, conducted during 2005-2007 in the semi-arid zone of Karnataka, the balanced nutrient management, BN significantly increased maize grain yield and dry matter over the farmer's inputs treatment; BN also significantly improved the harvest index of the crop during all the three seasons (Rajashekhara Rao et al. 2010).

The results of on-farm trials conducted in the SAT zone of Madhya Pradesh with soybean in the 2008, 2009 rainy season and chickpea in the 2008-2009 post-rainy seasons

confirmed the superiority of the BN treatment over the FI treatment and significantly increased soybean and chickpea grain yields (Table 5). Similar results were obtained in the on-farm trials conducted during the 2008 rainy season in the semi-arid zone of Rajasthan, India with pearl millet and maize as the test crops; and the grain yields of these crops were significantly increased in the BN treatment as compared to FI (Table 6).

On-farm trials were conducted during the 2006-2007 seasons with a number of vegetable crops in watersheds in three (Dharwad, Haveri and Chitradurga) districts of Karnataka to study their responses to balanced nutrient management as compared to farmer's input treatment. The results showed an impressive yield response to balanced nutrient management as compared to farmer's treatment; and the growing of these vegetable under balanced nutrient management crops was economically viable and remunerative (Srinivasarao et al. 2010).

Balanced plant nutrition is not only important for increasing crop productivity but is also critical for enhancing crop quality including grain and stover/straw quality, which has implications for human (grain as food) and animal (straw used as fodder or feed) nutrition. There is a relationship between soil health and food and feed quality which in turn impacts human and animal health. The importance of mineral nutrition of crops along with improved cultivars of crops and crop management cannot be overemphasized for producing nutritious food (Graham et al. 2007; Parthasarathy Rao et al. 2006; Sahrawat et al. 2008a) and fodder (Kelly et al. 1996; Sahrawat et al. 2008a; Rattan et al. 2009).

For example, in the on-farm experiments conducted to determine the effects of S, B and Zn fertilization on the grain and straw quality of sorghum and maize grown under rainfed conditions in the SAT region of India showed that the balanced mineral nutrition (BN) through combined application of S, B, Zn, N and P as compared to the FI (farmer's inputs) increased N, S and Zn concentrations in the grain and straw of these crops (Sahrawat et al. 2008a). These results stress the importance of balanced mineral nutrition of crops for increased produce quality. For example, the S fertilization of oilseed crops such as soybean, canola and sunflower is not only required for increasing dry matter and seed yield, but is also essential for enhancing oil concentration and quality (Saha et al. 2001; Usha Rani et al. 2009; Brennan et al. 2010).

From this discussion on the results obtained in on-farm trials, it is evident that in the SAT region multiple nutrient deficiencies especially of N, P, S, B and Zn are holding back the potential of rainfed systems and are also responsible for low rainwater use efficiency in rainfed areas in the SAT regions (Singh et al. 2009). Also, soil fertility depletion has been recognized as the major biophysical cause of declining food availability in smallholder farms in Sub-Saharan Africa. It was suggested that any program aimed at reversing the trend in declining agricultural productivity and food quality, and preserving the environmental quality must begin with soil fertility restoration and maintenance. The decline in productivity is related to decline in soil fertility, which in turn is directly related to decline in soil organic matter status and depletion of the plant nutrient reserves in various production systems with little or no investment in recuperating soil fertility in agroecosystems (Sanchez et al. 1997; Bationo et al. 2008; Lal 2008; Bekunda et al. 2010).

Soil fertility maintenance is not only a prerequisite for sustainable increase in crop productivity, but is equally essential for maintaining crop quality in terms of food, fodder and feed quality (Kelly et al. 1996; Sahrawat et al. 2008a), especially iron (Fe) and Zn in the grain (Graham et al. 2007; Sahrawat et al. 2008a; Rattan et al. 2009). The results from on-farm studies also show that the productivity of the rainfed systems can be enhanced through management of various nutrient deficiencies. It is demonstrated from the results of a large numbers of on-farm trials conducted in different parts of India that with soil test-based balanced nutrient management productivity in rainfed areas can be increased by harnessing the potential of rainfed agriculture. Unless the constraints to soil fertility management are alleviated, it would not be possible to achieve the potential productivity of the rainfed systems. Because the area under rainfed production is very large, even a modest sustainable increase in yield would contribute in a big way to global food pool, apart from providing source of income and livelihoods to the rural poor.

For practical utilization of the soil test-based nutrient management, we have been mapping using the GIS based extrapolation methodology, the deficiencies of nutrients especially those of S, B and Zn in various districts in the Karnataka state, India (ICRISAT, unpublished results). Finally, the soil test-based fertilizer application has been made web-based so that the recommendations can be downloaded and made available

nutrient wise to farmers using color codes depicting, the deficiency or sufficiency of a nutrient. Such information can be easily used by small holder farmers at the updated. Typical examples of nutrient mapping for extractable (available) S, B, and Zn, using data from selected districts of Karnataka, are shown in Figure 1. Such maps can be extended and used by farmers in a cluster of villages to plan the application of deficient plant nutrients to production systems.

General Discussion and Conclusions

It is recognized that water shortage related plant stress is the primary constraint to crop production and productivity in the rainfed systems in the SAT and consequently the importance of water shortage has globally been rightly emphasized (Wani et al. 2002, 2003; CAWMA 2007; Pathak et al. 2009). However, apart from water shortage, there is the issue of severe soil infertility problems in the rainfed systems (Rego et al. 2007; Sahrawat et al. 2010b; Bekunda et al. 2010) and managing water stress alone cannot sustainably enhance the productivity of rainfed systems; and hence for achieving sustainable gains in rainfed productivity both water shortage and soil fertility problems need to be simultaneously addressed through effective natural resource management practices (Wani et al. 2009; Sahrawat et al. 2010a).

For the first time, a large number of farmers' fields in the SAT regions of India were sampled and analyzed for organic C and extractable or available nutrients in an effort to diagnose the prevalence of major and micronutrient deficiencies. The results on the analyses of 28, 270 soil samples from the farmers' fields (Table 2) demonstrate that the soils in rainfed areas are indeed infertile and they are not only deficient in major nutrients especially N (soil organic C status used as an index for available N) and P but are low in organic matter reserve. The most revealing results however, were the widespread nature of the deficiencies of S, B and Zn (Rego et al. 2007; Sahrawat et al. 2007, 2010b).

A summary of results of on-farm responses of several field crops to applications of deficient nutrients together with N and P demonstrated that balanced nutrient management has indeed the potential to significantly enhance the productivity of a range of crops, improve grain and straw quality in the SAT regions under rainfed conditions.

It would appear from these results that soil test-based nutrient management approach can be an important entry point activity and also a mechanism to diagnose and manage soil fertility in practical agriculture (Wani 2008). Soil and plant tests have long been used as tools to diagnose and manage soil fertility problems in the intensified irrigated systems and commercial crops including fruit and vegetable crops to maximize productivity (Dahnke and Olson 1990; Mills and Jones 1990; Black 1993; Reuter and Robinson 1997). However, soil testing has not been used to diagnose and manage nutrient problems in farmers' fields in the SAT regions at a scale reported in this paper. The critical limits for P, K, S, B and Zn in the soil (Table 1) seem to provide a fair basis for separating deficient soils from those that are not deficient. Soils below the critical limits of the nutrients evaluated responded to the applications of nutrients; although the overall crop response was regulated by the rainfall received during the cropping season (Rego et al. 2007; Sahrawat et al. 2007, 2010b). Soil test-based nutrient application also allows judicious and efficient use of nutrient inputs at the local and regional levels (Black 1993; Sahrawat et al. 2010b).

For more widespread adoption and use of soil testing for the diagnosis and management of plant nutrient deficiencies in the rainfed systems of the SAT regions, there is need to strengthen the soil testing facilities at the local and regional levels for science-based management and maintenance of soil fertility, a prerequisite for sustainable increase in productivity of the rainfed systems (Sahrawat et al. 2007, 2010b). We do hope that the research reported in this paper would stimulate research for widespread use of soil testing as a means for soil fertility management in farmers' fields.

For enhancing the overall agricultural productivity and crop quality of the rainfed systems, the choice of crops and adapted cultivars along with soil, water and nutrient management practices need to be integrated at the farm level (Wani et al. 2009; Sahrawat et al. 2010b). To achieve this, research and extension support and backstopping along with capacity building of all the stake holders need to converge (Sahrawat et al. 2010 b; Wani 2008). Indeed, ICRISAT and its research partners most appropriately advocate the integration of genetics (crops and its cultivars, social aspects) and natural resource management for technology targeting and greater impact of agricultural research in the semi-arid tropics (Twomlow et al. 2008b). The strategy is based on the use of crop

cultivars that are adapted to the harsh conditions of the SAT regions especially water stress and nutrient deficiencies. The soil, water and nutrient management practices are developed around the adapted cultivars to realize the potential of the cultivars in diverse production systems (Ae et al. 1990; Condon et al. 2004; Hiradate et al. 2007; Passioura 2006; Bationo et al. 2008; Sahrawat 2009; Passioura and Angus 2010).

Acknowledgements

We thank the Andhra Pradesh Rural Livelihood Program of the government of Andhra Pradesh, the Department for International Development (DFID, UK), the World Bank Special Fund for Watershed Development, the government of Karnataka, the Sir Dorabji Tata Trust for the financial support of various projects under which the studies were carried out. We are also thankful to all farmers who participated in these studies.

References

- Ae, N., J. Arihara, K. Okada, T. Yoshihara, and C. Johansen. 1990. Phosphorus uptake by pigeon pea and its role in cropping systems of the Indian subcontinent. *Science* 248:477-480.
- Bationo, A., J. Kihara, B. Vanlauwe, J. Kimetu, B. S. Waswa, and K. L. Sahrawat. 2008. Integrated nutrient management: concepts and experience from Sub-Saharan Africa. In *Integrated nutrient management for sustainable crop production*, eds. M. S. Aulakh and C. A. Grant, 467-521. New York: The Haworth Press, Taylor and Francis Group.
- Bekunda, M., N. Sanginga, and P. L. Woomer. 2010. Restoring soil fertility in Sub-Saharan Africa. *Advances in Agronomy* 108:183-286.
- Black, C. A. 1993. *Soil fertility evaluation and control*, 155-452. Boca Raton, Florida, USA: Lewis Publishers.
- Brennan, R. F., R. W. Bell, C. Raphael, and H. Eslick. 2010. Sources of sulfur for dry matter, seed yield, and oil concentration of canola grown in sulfur-deficient soils of south-western Australia. *Journal of Plant Nutrition* 33:1180-1194.
- CAWMA. 2007. *Water for food, water for life: a comprehensive assessment of water*

- management in agriculture*. London, UK: Earthscan.
- Condon, A. G., R. A. Richards, G. J. Rebetzke, and G. D. Farquhar. 2004. Breeding for high water-use efficiency. *Journal of Experimental Botany* 55:2447-2460.
- Dahnke, W. C., and R. A. Olson. 1990. Soil test calibration, and the recommendation. In *Soil testing and plant analysis*, Third edition, Soil Science Society of America Series 3, ed. R. L. Westerman, 45-71. Madison, Wisconsin, USA: Soil Science Society of America.
- El-Swaify, S. A., P. Pathak, T. J. Rego, and S. Singh. 1985. Soil management for optimized productivity under rainfed conditions in the semi-arid tropics. *Advances in Soil Science* 1:1-64.
- Fageria, N. K., V. C. Baligar, and R. B. Clark. 2002. Micronutrients in crop production. *Advances in Agronomy* 77:185-268.
- Graham, R. D., R. M. Welch, D. A. Saunders, I. Ortiz-Monasterio, H. E. Bouis, M. Bonierbale, S. de Haan, G. Burgos, G. Thiele, R. Liria, C. A. Meisner, S. E. Beebe, M. J. Potts, M. Kadian, P. R. Hobbs, R. K. Gupta, and S. Twomlow. 2007. Nutritious subsistence food systems. *Advances in Agronomy* 92:1-74.
- Helmke, P. A., and D. L. Sparks. 1996. Lithium, sodium, potassium, rubidium, and cesium. In *Methods of soil analysis*, Part 3. Chemical Methods—Soil Science Society of America Book Series no. 5, ed. D. L. Sparks, 551-574. Madison, Wisconsin: SSSA and ASA.
- Hiradate, S., J. F. Ma, and H. Matsumoto. 2007. Strategies of plants to adapt to mineral stresses in problem soils. *Advances in Agronomy* 96:65-132.
- Kanwar, J. S. 1972. Twenty-five years of research in soil, fertilizer and water management in India. *Indian Farming* 22(5):16-25.
- Katyal, J. C. 2003. Soil fertility management—a key to prevent desertification. *Journal of the Indian Society of Soil Science* 51:378-387.
- Kelly, T. G., P. Parthasarathy Rao, R. E. Weltzien, and M. L. Purohit. 1996. Adoption of improved cultivars of pearl millet in arid environment: straw yield and quality consideration in western Rajasthan. *Experimental Agriculture* 32:161-171.

- Keren, R. 1996. Boron. In *Methods of soil analysis*, Part 3. Chemical Methods—Soil Science Society of America Book Series no. 5, ed. D. L. Sparks, 603-626. Madison, Wisconsin: SSSA and ASA.
- Lal, R. 2008. Soils and sustainable agriculture. A review. *Agronomy for Sustainable Development* 28:57-64.
- Lindsay, W. L., and W. A. Norvell. 1978. Development of a DTPA test for zinc, iron, manganese, and copper. *Soil Science Society of America Journal* 42:421-428.
- Mills, H. A., and J. B. Jones, Jr. 1996. *Plant analysis handbook: a practical sampling, preparation, and interpretation guide*. Athens, Georgia, USA: Micro Macro Publishing, Inc.
- Nelson, D.W., and L. E. Sommers. 1996. Total carbon, organic carbon, and organic matter. In *Methods of soil analysis*, Part 3. Chemical methods—Soil Science Society of America Book Series no. 5, ed. D. L. Sparks, 961-1010. Madison, Wisconsin: SSSA and ASA.
- Ncube, B., J. P. Dimes, S. J. Twomlow, W. Mupangwa, and K. E. Giller. 2007. Participatory on-farm trials to test response of maize to small doses of manure and nitrogen in small-holder farming systems in semi-arid Zimbabwe. *Nutrient Cycling in Agroecosystems* 77:53-67.
- Olsen, S. R., and L. E. Sommers. 1982. Phosphorus. In *Methods of soil analysis*, Part 2, Second Edition. Agronomy Monograph 9, ed. A. L. Page, 403-430. Madison, Wisconsin: ASA and SSSA.
- Parthasarathy Rao, P., P. S. BIRTHAL, B. V. S. Reddy, K. N. Rai, and S. Ramesh. 2006. Diagnostics of sorghum and pearl millet grains-based nutrition in India. *International sorghum and millets newsletter* 47:93-96.
- Pasricha, N. S., and R. L. Fox. 1993. Plant nutrient sulfur in the tropics and subtropics. *Advances in Agronomy* 50:209-269.
- Passioura, J. B. 2006. Increasing crop productivity when water is scarce—From breeding to field management. *Agricultural Water Management* 80:176-196.
- Passioura, J. B., and J. F. Angus. 2010. Improving productivity of crops in water-limited environments. *Advances in Agronomy* 106:37-74.
- Pathak, P., K. L. Sahrawat, S. P. Wani, R. C. Sachan, and R. Sudi. 2009. Opportunities

- for water harvesting and supplemental irrigation for improving rainfed agriculture in semi-arid areas. In *Rainfed agriculture: Unlocking the potential*, eds. S. P. Wani, J. Rockström, and T. Oweis, 197-221. Wallingford, UK: CABI International.
- Rajashekhara Rao, B. K., K. L. Sahrawat, S. P. Wani, and G. Pardhasaradhi. 2010. Integrated nutrient management to enhance on-farm productivity of rain fed maize in India. *International Journal of Soil Science* 5:216-225.
- Rattan, R. K., K. P. Patel, K. M. Manjaiah, and S. P. Datta. 2009. Micronutrients in soil, plant, animal and human health. *Journal of the Indian Society of Soil Science* 57:546-558.
- Rego, T. J., V. N. Rao, B. Seeling, G. Pardhasaradhi, and J. V. D. K. Kumar Rao. 2003. Nutrient balances—a guide to improving sorghum and groundnut-based dryland cropping systems in semi-arid tropical India. *Field Crops Research* 81:53-68.
- Rego, T. J., S. P. Wani, K. L. Sahrawat, and G. Pardhasaradhi. 2005. *Macro-benefits from boron, zinc and sulfur application in Indian SAT: a step for gray to green revolution in agriculture*. Global Theme on Agroecosystems Report no. 16. Patancheru 502 324, Andhra Pradesh, India: International Crops Research Institute for the Semi-Arid Tropics.
- Rego, T. J., K. L., Sahrawat, S. P. Wani, and G. Pardhasaradhi. 2007. Widespread deficiencies of sulfur, boron, and zinc in Indian semi-arid tropical soils: on-farm crop responses. *Journal of Plant Nutrition* 30:1569-1583.
- Reuter, D. J., and J. B. Robinson, eds. 1997. *Plant analysis: an interpretation manual*, Second edition. Australia: CSIRO.
- Rockström, J., L. Karlberg, S. P. Wani, J. Barron, N. Hatibu, T. Oweis, A. Bruggeman, J. Farahani, and Z. Qiang. 2010. Managing water in rainfed agriculture—The need for a paradigm shift. *Agricultural Water Management* 97:543-550.
- Saha, J. K., A. B. Singh, A. N. Ganeshamurthy, S. Kundu, and A. K. Biswas. 2001. Sulfur accumulation in Vertisols due to continuous gypsum application for six years and its effect on yield and biochemical constituents of soybean (*Glycine Max L. Merrill*). *Journal of Plant Nutrition and Soil Science* 164:317-320.

- Sahrawat, K. L. 2006. Plant nutrients: sufficiency and requirements. In *Encyclopedia of Soil Science*, Second Edition, ed. R. Lal, 1306-1310. Philadelphia, PA, USA: Taylor and Francis.
- Sahrawat, K. L. 2009. The role of tolerant genotypes and plant nutrients in reducing acid-soil infertility in upland rice ecosystem: an appraisal. *Archives of Agronomy and Soil Science* 55:597-607.
- Sahrawat, K. L., M. K. Abekoe, and S. Diatta. 2001. Application of inorganic phosphorus fertilizer. In *Sustaining Fertility in West Africa*, eds. G. Tian, , F. Ishida, and D. Keatinge, Soil Science Society of America Special Publication no. 58, 225-246. Madison, Wisconsin, USA: Soil Science Society of America and American Society of Agronomy.
- Sahrawat, K. L., S. P. Wani, T. J. Rego, G. Pardhasaradhi and K. V. S. Murthy. 2007. Widespread deficiencies of sulphur, boron and zinc in dryland soils of the Indian semi-arid tropics. *Current Science* 93:1428-1432.
- Sahrawat, K. L., T. J. Rego, S. P. Wani, and G. Pardhasaradhi. 2008a. Sulfur, boron and zinc fertilization effects on grain and straw quality of maize and sorghum grown on farmers' fields in the semi-arid tropical region of India. *Journal of Plant Nutrition* 31:1578-1584.
- Sahrawat, K. L., T. J. Rego, S. P. Wani, and G. Pardhasaradhi. 2008b. Stretching soil sampling to watershed: Evaluation of soil-test parameters in a semi-arid tropical watershed. *Communications in Soil Science and Plant Analysis* 39:2950-2960.
- Sahrawat, K. L., K. V. S. Murthy, and S. P. Wani. 2009. Comparative evaluation of Ca chloride and Ca phosphate for extractable sulfur in soils with a wide range in pH. *Journal of Plant Nutrition and Soil Science* 172:404-407.
- Sahrawat, K. L., S. P. Wani, P. Pathak, and T. J. Rego. 2010a. Managing natural Resources of watersheds in the semi-arid tropics for improved soil and water quality: A review. *Agricultural Water Management* 97:375-381.
- Sahrawat, K. L., S. P. Wani, G. Pardhasaradhi, and K. V. S. Murthy. 2010b. Diagnosis of secondary and micronutrient deficiencies and their management in rainfed Agroecosystems: Case study from Indian semi-arid tropics. *Communications in Soil Science and Plant Analysis* 41:346-360.

- Sanchez, P. A., K. D. Shepherd, M. J. Soul, F. M. Place, R. J. Buresh, A. M. N. Izac, A. U. Mokwunye, F. R. Kwesiga, C. G. Nderitu, and P. L. Woomer. 1997. Soil fertility replenishment in Africa: An investment in natural resource capital. In *Replenishing Soil Fertility in Africa*, eds. R. J. Buresh, P. A. Sanchez, and F. Calhoun, Soil Science Society of America Special Publication no. 51, 1-46. Madison, Wisconsin, USA: Soil Science Society of America.
- Scherer, H. W. 2001. Sulphur in crop production. *European Journal of Agronomy* 14:81-111.
- Scherer, H. W. 2009. Sulfur in soils. *Journal of Plant Nutrition and Soil Science* 172:326-335.
- Sharma, B. R., K. V. Rao, K. P. R. Vittal, Y. S. Ramakrishna, and U. Amarasinghe. 2010. Estimating the potential of rainfed agriculture in India: Prospects for water productivity improvement. *Agricultural Water Management* 97:23-30.
- Singh, A. K. 2008. Soil resource management—key to food and health security. *Journal of the Indian Society of Soil Science* 56:348-357.
- Singh, P., P. Pathak, S. P. Wani, and K. L. Sahrawat. 2009. Integrated watershed management for increasing productivity and water-use efficiency in semi-arid tropical India. *Journal of Crop Improvement* 23:402-429.
- Srinivasarao, Ch., S. P. Wani, K. L. Sahrawat, K. Krishnappa, and B. K. Rajasekhara Rao. 2010. Effect of balanced nutrition on yield and economics of vegetable crops in participatory watersheds in Karnataka. *Indian Journal of Fertilisers* 6:39-42.
- Tabatabai, M. A. 1996. Sulfur. In *Methods of Soil Analysis*, Part 3. Chemical Methods—Soil Science Society of America Book Series no. 5, ed. D. L. Sparks, 921-960. Madison, Wisconsin: SSSA and ASA.
- Takkar, P. N. 1996. Micronutrient research and sustainable agricultural productivity. *Journal of the Indian Society of Soil Science* 44:563-581.
- Twomlow, S., D. Love, and S. Walker. 2008a. The nexus between integrated natural resources management and integrated water resources management in southern Africa. *Physics and Chemistry of the Earth* 33:889-898.
- Twomlow, S., B. Shiferaw, P. Cooper, and J. D. H. Keatinge. 2008b. Integrating genetics and natural resource management for technology targeting and greater impact of

- agricultural research in the semi-arid tropics. *Experimental Agriculture* 44:235-256.
- Usha Rani, K., K. L. Sharma, K. Nagasri, K. Srinivas, T. Vishnu Murthy, G. R. Maruthi Shankar, G. R. Korwar, K. S. Sankar, M. Madhavi, and J. K. Grace. 2009. Response of sunflower to sources and levels of sulfur under rainfed semi-arid tropical conditions. *Communications in Soil Science and Plant Analysis* 40:2926-2944.
- Wani, S. P. 2008. Taking soil science to farmers' doorsteps through community watershed management. *Journal of the Indian Society of Soil Science* 56:367-377.
- Wani, S. P., P. Pathak, H. M. Tam, A. Ramakrishna, P. Singh, and T. K. Sreedevi. 2002. Integrated watershed management for minimizing land degradation and sustaining productivity in Asia. In *Integrated Land Management in Dry Areas*. Proceedings of a Joint UNU-CAS International Workshop, ed. Zafar Adeel, pp. 207-230, 8-13 September 2001, Beijing, China.
- Wani, S.P., P. Pathak, L. S. Jangawad, H. Eswaran, and P. Singh. 2003. Improved management of Vertisols in the semiarid tropics for increased productivity and soil carbon sequestration. *Soil Use and Management* 19:217-222.
- Wani, S. P., T. K. Sreedevi, J. Rockström, and Y. S. Ramakrishna. 2009. Rainfed agriculture—past trends and future prospects. In *Rainfed agriculture: Unlocking the potential*, eds. S. P. Wani, J. Rockström, and T. Oweis, 1-35. Wallingford, UK: CAB International.
- Zougmore, R., Z. Zida, and N. F. Kambou. 2003. Role of nutrient amendments in the success of half-moon soil and water conservation practice in semiarid Burkina Faso. *Soil and Tillage Research* 71:143-149.

Table 1. Critical limits in the soil of plant nutrient elements to separate deficient samples from non-deficient samples. The data gleaned from various literature sources (for details see Rego et al. 2007; Sahrawat et al. 2007)

Plant nutrient	Critical limit (mg kg ⁻¹)
Sodium bicarbonate-extractable P	5
Ammonium Acetate-extractable K	50
Calcium chloride-extractable S	8-10
Hot water-extractable B	0.58
DTPA-extractable Zn	0.75

Table2. Chemical characteristics of soil samples collected from farmers' fields in the SAT regions of India.

Andhra Pradesh

District / (No.of Fields)	Parameter	pH	Organic C (%)	Olsen P (mg kg ⁻¹)	Exch.K (mg kg ⁻¹)	Extractable nutrient elements (mg kg ⁻¹)		
						S	B	Zn
Adilabad (63)	Range	6.4--8.9	0.27--1.33	0.2--48.8	46--549	2.0--142.2	0.10--0.74	0.22--2.90
	Mean	8.2	0.62	6.9	204	12.2	0.34	0.62
	% Deficient*			60	2	76	92	75
Anantapur (593)	Range	5.4--9.6	0.11--1.45	0.6--42.4	14--352	0.2--117.3	0.02--1.40	0.14--5.00
	Mean	7.5	0.30	7.7	73	4.5	0.21	0.59
	% Deficient			33	31	94	98	83
Kadapa (114)	Range	5.3--8.8	0.11--0.79	0.2--25.4	17--387	1.7--41.9	0.04--3.02	0.24--5.20
	Mean	7.4	0.27	3.9	80	6.6	0.39	0.76
	% Deficient			75	43	85	81	67
Khammam (102)	Range	5.1--8.8	0.32--1.50	0.2--57.8	31--856	3.6--71.9	0.12--1.22	0.28--6.80
	Mean	6.8	0.70	8.5	180	10.6	0.39	1.09
	% Deficient			60	2	67	87	45
Kurnool (331)	Range	5.6--9.7	0.09--1.06	0.4--36.4	33--509	1.4--53.8	0.04--2.04	0.08--4.92
	Mean	7.9	0.34	7.6	144	6.3	0.37	0.45
	% Deficient			42	5	85	79	91
Mahbubnagar (1035)	Range	5.3--10.2	0.08--2.18	0.2--247.7	16--1263	1.2--801.0	0.02--4.58	0.12--35.60
	Mean	7.4	0.42	12.6	119	16.2	0.30	1.11
	% Deficient			25	10	60	88	59
Medak (258)	Range	5.0--9.1	0.09--3.00	0.5--75.1	11--978	1.7--431.0	0.08--1.84	0.24--3.26
	Mean	7.7	0.49	8.0	161	12.4	0.57	0.78
	% Deficient			45	11	78	59	57
Nalgonda (441)	Range	5.0--9.2	0.12--1.36	0.2--50.4	21--379	1.4--140.3	0.02--1.48	0.08--16.00
	Mean	7.6	0.42	8.9	120	10.2	0.30	0.82
	% Deficient			31	7	78	90	66
Prakasam (492)	Range	6.4--9.3	0.12--1.30	0.2--41.7	28--697	0.6--19.2	0.02--1.86	0.20--10.8
	Mean	8.4	0.43	5.7	205	4.1	0.45	0.53
	% Deficient			56	1	94	71	88
Ranga Reddi (121)	Range	5.1--8.2	0.15--1.56	0.2--60.0	24--405	1.1--81.6	0.06--1.24	0.30--5.72
	Mean	6.7	0.50	8.9	92	3.7	0.26	1.16
	% Deficient			39	17	98	98	35
Warangal (100)	Range	6.1--9.4	0.08--0.84	0.2--53.4	21--280	1.8--48.9	0.10--1.42	0.26--3.88
	Mean	7.8	0.41	16.0	118	9.4	0.38	0.96
	% Deficient			14	5	77	84	50
Andhra Pradesh State Total (3650)	Range	5.0--10.2	0.08--3.00	0.2--247.7	11--1263	0.2--801	0.02--4.58	0.08--35.6
	Mean	7.6	0.41	9.1	129	9.6	0.34	0.81
	% Deficient			38	12	79	85	69

Karnataka

District / (No. of Fields)	Parameter	pH	Organic C (%)	Olsen P (mg kg ⁻¹)	Exch.K (mg kg ⁻¹)	Extractable nutrient elements (mg kg ⁻¹)		
						S	B	Zn
Bengaluru Rural (2223)	Range Mean % Deficient	5.0-9.5 6.4	0.01--1.31 0.41	0.3--220.8 18.9 16	9--847 93 30	0.9-94.5 5.4 94	0.10--5.12 0.39 68	0.14-235 1.47 34
Bidar (1189)	Range Mean % Deficient	5.6-8.7 7.6	0.19--1.98 0.63	0.6--118.6 8.5 49	18--2297 221 1	1.0-181.3 7.2 84	0.12--2.96 0.56 65	0.16-18 0.94 55
Bijapur (1395)	Range Mean % Deficient	6.7--9.2 8.2	0.00--1.21 0.44	0.1--91.9 3.9 80	24--2613 225 3	0.9-4647.4 38.5 77	0.02--18.22 0.93 46	0.15--10.4 0.58 85
Chamaraja Nagara (818)	Range Mean % Deficient	5.1--9.7 7.8	0.05--1.85 0.43	0.2--77.5 9.6 40	25--738 188 3	0.4-119.4 5.6 90	0.08--3.80 0.63 57	0.14--6.4 0.77 62
Chikkaballapur (2257)	Range Mean % Deficient	5.0--9.9 6.9	0.07--1.42 0.39	0.2--430.8 18.0 37	4-1650 95 34	0.5-470.0 9.1 80	0.06--1.98 0.38 80	0.06--21.5 1.15 52
Chitradurga (1489)	Range Mean % Deficient	5.1--10.1 7.8	0.03--1.36 0.40	0.2--480.0 7.0 54	12--1953 137 15	0.8-291.8 7.3 86	0.04--6.94 0.63 64	0.08--40.5 0.64 80
Davangere (1500)	Range Mean % Deficient	5.0--9.0 7.0	0.04--1.38 0.51	0.0--138.8 13.1 34	11--510 109 13	0.9-945.0 12.7 77	0.06--6.30 0.54 66	0.04--11.2 0.74 74
Dharwad (1129)	Range Mean % Deficient	5.1--9.3 7.4	0.17--1.99 0.65 31	0.2--207.0 9.3 53	36--2344 220 1	1.4-715.0 9.7 79	0.10--12.48 0.82 39	0.24--24.3 0.98 44
Gadag (655)	Range Mean % Deficient	5.0--9.2 8.1	0.04--1.41 0.44	0.0--65.6 5.3 65	27--526 178 2	1.0-223.3 7.4 85	0.08--9.62 0.88 36	0.06--4.9 0.44 90
Gulbarga (2811)	Range Mean % Deficient	5.1--10.0 8.0	0.01--2.50 0.46 65	0.0--97.3 7.1 58	14--1722 244 2	0.4-12647 27.6 79	0.02--24.90 0.64 66	0.10--14.8 0.52 87
Haveri (1532)	Range Mean % Deficient	5.1--10.5 7.7	0.08--3.60 0.51	0.1--143.0 12.4 42	25--3750 133 5	0.3-120.3 7.0 85	0.08--8.44 0.71 46	0.20--34.1 0.81 60
Kolar (2161)	Range Mean % Deficient	5.0--10.2 7.0	0.04--1.50 0.38 81	0.0--182.0 20.3 31	9--1144 87 34	0.7-141.2 7.0 85	0.04--1.82 0.34 87	0.14--14.4 1.31 32
Raichur (1667)	Range Mean % Deficient	5.1--9.7 8.3	0.05--1.48 0.43	0.2--169.6 11.8 47	13-1797 209 4	0.8--2488 46.8 64	0.04--26.24 1.17 37	0.12--15.24 0.66 78
Tumkur (3041)	Range Mean % Deficient	5.0--10.0 6.6	0.04--2.08 0.39	0.1--204.0 5.9 65	11--1470 92 34	0.1-128.4 5.5 92	0.03--3.60 0.33 91	0.14--17.26 0.89 50
Karnataka State Total (22867)	Range Mean % Deficient	5.0--10.5 7.4	0.01--3.6 0.45	0.1--480 11.4 47	4--3750 150 16	0.1--12647 14.4 83	0.02--26.24 0.59 66	0.04-235 0.89 61

Madhya Pradesh

District / (No. of Fields)	Parameter	pH	Organic C (%)	Olsen P (mg kg ⁻¹)	Exch.K (mg kg ⁻¹)	Extractable nutrient elements (mg kg ⁻¹)		
						S	B	Zn
Badwani	Range	7.6--8.4	0.28--0.76	0.5--18.4	73--299	4.0--40.4	0.18--0.70	0.30--1.14
(20)	Mean	8.1	0.51	4.6	146	11.8	0.42	0.58
	% Deficient			70	0	55	80	75
Dewas	Range	7.0--8.7	0.31--1.00	0.2--10.8	46--456	3.9--9.5	0.12--0.56	0.24--0.82
(24)	Mean	8.0	0.60	2.1	137	6.3	0.24	0.45
	% Deficient			96	4	100	100	96
Guna	Range	7.2--8.5	0.47--1.11	0.1--10.2	86--303	2.7--14.3	0.22--2.20	0.24--1.74
(38)	Mean	8.0	0.65	3.2	158	6.3	0.67	0.51
	% Deficient			79	0	87	50	95
Indore	Range	7.8--8.3	0.43--1.08	0.5--42.2	129--716	5.9--134.4	0.46--1.30	0.56--3.00
(23)	Mean	8.1	0.66	10.4	263	29.7	0.82	1.11
	% Deficient			39	0	9	17	22
Jhabua	Range	6.4--7.4	0.58--1.53	0.2--42.2	88--506	2.7--28.2	0.26--0.76	0.66--3.18
(22)	Mean	7.0	0.88	9.7	216	6.3	0.40	1.54
	% Deficient			45	0	95	91	5
Mandla	Range	5.9--7.2	0.45--1.25	1.0--7.2	82--287	2.0--13.2	0.06--0.80	0.48--1.14
(21)	Mean	6.6	0.68	2.8	143	4.8	0.29	0.79
	% Deficient			90	0	90	86	52
Raisen	Range	7.9--8.4	0.42--0.97	0.5--13.4	118--275	2.9--12.8	0.20--0.74	0.30--0.98
(20)	Mean	8.1	0.58	3.1	199	6.2	0.35	0.49
	% Deficient			90	0	90	90	90
Rajgarh	Range	6.7--8.3	0.44--1.41	1.6--19.2	51--434	2.9--50.4	0.30--0.92	0.38--3.82
(30)	Mean	7.9	0.78	5.7	203	12.3	0.49	1.14
	% Deficient			60	0	53	73	27
Sagar	Range	6.7--8.0	0.42--2.19	0.5--68.0	149--333	4.2--23.8	0.18--1.22	0.50--3.10
(32)	Mean	7.4	0.72	7.1	265	10.1	0.36	1.04
	% Deficient			78	0	63	91	34
Sehore	Range	7.3--8.4	0.36--0.69	0.5--17.2	48--256	3.0--20.5	0.28--0.62	0.36--0.92
(19)	Mean	8.1	0.50	4.0	167	8.3	0.39	0.53
	% Deficient			84	5	74	95	95
Shajapur	Range	7.1--8.2	0.46--1.15	1.0--25.8	51--249	5.6--42.0	0.18--0.72	0.46--1.42
(20)	Mean	7.7	0.82	8.7	120	17.2	0.43	0.85
	% Deficient			25	0	25	80	40
Vidisha	Range	7.6--8.6	0.31--0.92	0.5--14.1	96--401	1.8--16.6	0.12--0.74	0.10--1.00
(72)	Mean	8.2	0.56	2.8	203	5.5	0.35	0.34
	% Deficient			92	0	96	93	97
Madhya Pradesh State Total	Range	5.9--8.7	0.28--2.19	0.1--68	46--716	1.8--134.4	0.06--2.2	0.10--3.82
(341)	Mean	7.8	0.65	5.0	190	9.6	0.43	0.72
	% Deficient			74	1	74	79	66

Rajasthan

District / (No.of Fields)	Parameter	pH	Organic C (%)	Olsen P (mg kg ⁻¹)	Exch.K (mg kg ⁻¹)	Extractable nutrient elements (mg kg ⁻¹)		
						S	B	Zn
Alwar (30)	Range	7.9--8.8	0.33--0.66	0.5--44.0	53--515	4.5--17.2	0.20--0.68	0.20--2.00
	Mean	8.5	0.46	14.3	128	9.2	0.45	0.56
	% Deficient			10	0	63	87	83
Banswara (30)	Range	6.3--8.1	0.28--1.05	1.0--35.0	31--418	2.4--22.0	0.10--0.54	0.26--2.60
	Mean	7.2	0.56	7.7	107	9.2	0.23	0.70
	% Deficient			50	17	70	100	80
Bhilwara (30)	Range	7.2--8.9	0.32--1.87	0.8--27.0	33--460	4.0--44.9	0.32--1.30	0.16--2.30
	Mean	8.3	0.74	9.2	111	12.8	0.64	0.92
	% Deficient			40	17	43	47	37
Bundi (36)	Range	6.2--8.7	0.18--1.17	0.9--20.1	23--563	3.3--51.0	0.10--0.98	0.20--1.78
	Mean	7.6	0.60	6.2	87	9.2	0.44	0.65
	% Deficient			53	50	72	72	67
Dungarpur (99)	Range	6.2--8.0	0.48--1.99	1.0--28.2	34--240	4.0--31.3	0.28--1.50	0.88--14.10
	Mean	6.9	1.26	6.6	100	9.0	0.70	2.11
	% Deficient			48	8	72	31	0
Jhalawar (30)	Range	8.0--8.6	0.46--1.15	0.9--22.6	51--1358	1.9--78.0	0.22--1.36	0.40--3.40
	Mean	8.4	0.76	10.2	214	8.3	0.49	0.75
	% Deficient			30	0	87	77	60
Sawai Madhopur (44)	Range	7.8--9.4	0.16--0.70	0.2--11.8	44--438	3.1--26.6	0.20--2.18	0.34--28.60
	Mean	8.5	0.38	4.0	137	6.8	0.64	2.54
	% Deficient			73	7	86	52	41
Tonk (78)	Range	6.8--10.2	0.09--1.11	0.2--28.2	14--243	2.3--29.8	0.08--2.46	0.18--14.00
	Mean	8.1	0.36	5.7	83	7.7	0.62	1.61
	% Deficient			55	32	79	64	58
Udaipur (44)	Range	7.3--9.0	0.25--2.37	2.6--41.0	52--288	3.2--274.0	0.22--1.50	0.70--3.92
	Mean	8.2	0.83	15.2	145	26.7	0.83	1.57
	% Deficient			18	0	48	25	5
Rajasthan State Total (421)	Range	6.2--10.2	0.09--2.37	0.2--44	14--1358	1.9--274	0.08--2.46	0.16--28.6
	Mean	7.8	0.72	8.1	116	10.6	0.6	1.49
	% Deficient			45	15	71	56	40
Grand total (28270)	Range	5.0--10.5	0.01--3.6	0.1--480	4--3750	0.1--12647	0.02--26.24	0.04--235
	Mean	7.4	0.45	10.9	147	13.6	0.55	0.88
	% Deficient			46	16	82	68	62

Source: The results on soil analyses of samples from Andhra Pradesh, Rajasthan and Madhya Pradesh are from Sahrawat et al. (2007). The data on Karnataka soil samples are from unpublished ICRISAT results.

Table 3. Grain yields of crops in response to fertilization according to farmer's inputs (FI) and balanced nutrient management (BN, BN = FI +SBZn+NP) treatments in the semi-arid zone of Andhra Pradesh, India during three (2002 to 2004) rainy seasons

Year	Treatment	Grain yield (kg ha ⁻¹)				
		Maize	Castor	Mung bean	Groundnut (pod)	Pigeonpea
2002	FI	2730 (20) ^a	590 (8)	770 (9)	1180 (19)	536 (43)
	BN	4560	880	1110	1570	873
	LSD(0.05)	419	143	145	92	156
2003	FI	2790 (24)	690 (17)	900 (6)	830 (30)	720 (12)
	BN	4880	1190	1530	1490	1457
	LSD(0.05)	271	186	160	96.8	220
2004	FI	2430 (19)	990 (6)	740 (12)	1320 (40)	1011 (21)
	BN	4230	1370	1160	1830	1564
	LSD(0.05)	417	285	131	122.5	106

Source: The results on maize, castor, mung bean and groundnut crops are from Rego et al. (2007), and the data on pigeonpea crop are from ICRISAT unpublished results.

^aThe value in parenthesis is the number of farmers' fields on which on-farm trials were conducted.

Table 4. Grain yields of crops in response to fertilization according to farmer's inputs (FI) and balanced nutrient management (BN, BN = FI +SBZn+NP) treatments in the semi-arid zone of Karnataka, India during five (2005 to 2009) rainy seasons

Year	Treatment	Grain yield (kg ha ⁻¹)			
		Maize	Finger Millet	Groundnut	Soybean
2005	FI	4000 (6) ^a	2100 (16)	1830 (8)	2030 (6)
	BN(0.05)	6090	3280	1910	3470
	LSD	395	338	91.5	664
2006	FI	4050 (22)	1700 (17)	1080 (17)	1120 (7)
	BN	5400	2170	1450	2650
	LSD(0.05)	240	440	341.4	538
2007	FI	5670 (19)	2000 (27)	1310 (23)	2120 (11)
	BN	8710	2940	2160	3120
	LSD(0.05)	572	230	191.4	262
2008	FI	4400 (27)	1680 (152)	940 (149)	1390 (16)
	BN	6130	2650	1430	1640
	LSD(0.05)	336	125	80.3	249
2009	FP	5460 (90)	1630 (165)	1100 (178)	1770 (36)
	IP	7800	2570	1500	2610
	LSD(0.05)	178	91	49.9	184

Source: Unpublished results from ICRISAT.

^a The value in parenthesis is the number of farmers' fields on which on-farm trials were conducted.

Table 5. Grain yields of soybean (rainy season) and chickpea (post-rainy season) in response to fertilization according to farmer’s inputs (FI) and balanced nutrient management (BN, BN = FI + SBZn + NP) treatments in Madhya Pradesh, India during 2008 and 2008-2009 seasons

Year	Treatment	Grain yield (kg ha ⁻¹)	
		Soybean	Chickpea
2008	FI	1490 (117) ^a	1250 (169)
	BN	1840	1440
	LSD(0.05)	56	29
2009	FI	2120 (140)	
	BN	2680	
	LSD(0.05)	95	

Source: Unpublished results from ICRISAT.

^aThe value in parenthesis is the number of farmers’ fields on which on-farm trials were conducted.

Table 6. Yields of maize and pearl millet in response to fertilization according to farmer’s inputs (FI) and balanced nutrient management (BN, BN = FI +SBZn + NP) treatments in the semi-arid zone of Rajasthan, India during the 2008 rainy season

Year	State	Treatment	Grain yield (kg ha ⁻¹)	
			Maize	Pearl Millet
2008		FI	2730 (17) ^a	2310 (16)
		BN	2980	2510
		LSD(0.05)	55	34.3

Source: ICRISAT unpublished results

^aThe value in parenthesis is the number of farmers’ fields on which on-farm trials were conducted.



