

## **Incidence of Winter and Summer Diapause in *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) in Andhra Pradesh, India**

<sup>1</sup>Deepak R. Jadhav, <sup>2</sup>Nigel J. Armes and <sup>1</sup>Vinod S. Bhatnagar

<sup>1</sup>International Crops Research Institute for the Semi-Arid Tropics (ICRISAT),  
Patancheru 502 324 Andhra Pradesh, India

<sup>2</sup>BASF Corporation, 26 Davis Drive, Research Triangle Park, North, Carolina 27709, USA

**Abstract:** The incidence of winter and summer diapause was studied by large-scale field samplings of larvae and pupae of the cotton bollworm, *Helicoverpa armigera* (Hüb) conducted over a period of 13-years during winter and summer seasons in Andhra Pradesh, India. Induction of winter diapause was associated with cooler prevailing temperatures and shorter day lengths of <12 h in November-January. On the other hand, summer diapause was observed at temperatures >32°C during March-April resulting in greater adult emergence with the onset of the monsoon season during mid-June to early July. In summer, few alternate hosts are available for larvae to sustain by allowing local populations colonizing newly germinated host crops with the onset of the rainy season. In addition, the incidence of winter and summer diapause was higher in male pupae than females in the ratio of 4:1 and 3:1 (♂:♀), respectively. Diapause populations of *H. armigera* were most common in the cyclonic weather prevailed for several weeks during 1977-78 (15.11%) and 1995-96 (17.64%). Under these conditions, an average of <4% and <6% of pupae entered winter and summer diapause, respectively and was associated with severe outbreaks of *H. armigera* in subsequent seasons. Higher populations of *H. armigera* on pigeonpea and chickpea entered winter diapause on pigeonpea and chickpea and summer diapause on irrigated maize, pigeonpea, chickpea and a common weed host, *Datura metel*.

**Keywords:** *Heliothis armigera*, monsoon, population dynamics, summer diapause, temperature, winter diapause

### **INTRODUCTION**

The cotton bollworm, *Helicoverpa armigera* (Hübner) is widely distributed across different continents throughout Africa, southern Europe, the Middle East, India, south-east Asia, Australia, eastern and northern New Zealand and many Pacific Islands (Fitt, 1989), including the USA (Borchert *et al.*, 2003). In Peninsular India, *H. armigera* is a polyphagous pest infesting leguminous, fiber, cereal and horticultural crops (Reed and Pawar, 1982). The annual losses due to cotton and pulses alone by *H. armigera* have been estimated as US \$500 billion (Shanower *et al.*, 1999). Since the first major outbreak of *H. armigera* reported in 1977-78 on chickpea (ICRISAT, 1978) and pigeonpea (Rao *et al.*, 1990) in Andhra Pradesh, India, it has gained importance as a pest in cotton and food crop agroecosystems. Moreover, its pest status in southern India has been exacerbated by the extension of growing season due to increasing irrigation facilities and use of new cultivars allowing for protracted host availability compared to the traditional agricultural practices where fallow periods were common. In addition, resistance to most of the commonly available

insecticide classes (Daly and Murray, 1988; Armes *et al.*, 1996), the ability for long-range migration (Farrow and Daly, 1987) has made control measures in fruituous. Other factors contributing to increased pest status include: ability to survive over a wide diversity of host plants (Zalucki *et al.*, 1986, 1994; Fitt, 1989; Murray, 1991; Hamamura, 1998; Casimero *et al.*, 2000), high fecundity (Reed, 1965) and ability to escape adverse seasons *via* diapause (Roome, 1979; Wilson *et al.*, 1979; Eger *et al.*, 1982).

It completes 7-8 generations annually and adults emerge in two weeks depending on the soil temperature. In Southern India, the rainfall is a key climatic factor that influences the regional population dynamics in any season. Higher rainfall early in the season greatly influences the population buildup. Several reports are available on the ability of *H. armigera* to enter pupal diapause (Reed, 1965; Hardwick, 1965; Wilson *et al.*, 1979; Roome, 1979; Hackett and Gatehouse, 1982; Fitt, 1989; Nibouche, 1998; Qureshi *et al.*, 1999, 2000; Shimuzi and Fujisaki, 2002; Shimuzi *et al.*, 2006; Kurban *et al.*, 2007), whose intensity is governed by latitude (Danilevski, 1965; Beck, 1968, 1980; Wu and Guo, 1997b). The influence




Fig. 1: Sampling of *H. armigera* larval populations conducted in various districts of Andhra Pradesh, India (• Marked denote the districts surveyed)

of temperature and photoperiod on the induction, maintenance and termination of winter diapause in *H. armigera* has been reported from Russia (Kuznetsova, 1972), India (Bhatnagar and Davies, 1978), Botswana (Roome, 1979), Australia (Wilson *et al.*, 1979; Kay, 1982), China (Wu and Guo, 1997a; Ge *et al.*, 2005), Japan (Qureshi *et al.*, 1999), Israel (Zhou *et al.*, 2000), the USA (Borchert *et al.*, 2003) and Greece (Mironidis *et al.*, 2010). Similarly, the induction of diapause resulting from long photoperiods and high temperatures during summer has been reported (Masaki, 1980; Butler *et al.*, 1985; Denlinger, 1986,

2002; Danks, 1987, 2005; DeSouza *et al.*, 1995; Nibouche, 1998). In Andhra Pradesh, winter diapause has been reported as a mechanism allowing the survival of populations when few alternate crops are available for the off-season survival of larvae in summer (Bhatnagar and Davies, 1978). King (1994) reported the occurrence of a low proportion of diapause pupae of <2 and 1% in winter and summer seasons, respectively. Martinat (1987) reported that severe cyclones are associated with the occurrence of heavy to incessant rains during November-December, especially in coastal districts of Andhra Pradesh. Normally, the cyclonic

rains continue for 4-10 days with cool and overcast cloudy conditions, which can have a significant impact on the larvae of *H. armigera*. Very few studies have been conducted on the ecological performance of diapause in *H. armigera* (Hackett and Gatehouse, 1982; Nibouche, 1998; Masaki, 2002) and *H. virescens* (Potter and Watson, 1980; Butler *et al.*, 1985). Very few studies are available on the seasonal incidence of diapause in *H. armigera*. These studies are based on long term field work and investigate to relate the agroclimatic factors influencing the incidence of diapause of *H. armigera* in tropical agroecosystems.

**MATERIALS AND METHODS**

Over a 13 year period from 1974-96 large-scale periodic field samplings of *H. armigera* larvae were made on crops such as pigeon pea (*Cajanus cajan* (L.) Millsp.), chickpea (*Cicer arietinum* L.), maize (*Zea mays* L.), sorghum (*Sorghum bicolor* (L.) Moench), pearl millet (*Pennisetum americanum* (L.) Leeke), tomato (*Lycopersicon esculentum* Mill.), cowpea (*Vigna unguiculata* (L.) Walp), black gram (*Vigna mungo* (L.) Hepper.), green gram (*Vigna radiata* (L.) Wilczek), cotton (*Gossypium* spp.), safflower (*Carthamus tinctorius* L.), linseed (*Linum usitatissimum* L.),

sunflower (*Helianthus annuus* L.), soyabean (*Glycine max* (L.) Merrill), marigold (*Tagetes erecta* L.), groundnut (*Arachis hypogaea* L.), peas (*Pisum sativum* L.), okra (*Abelmoschus esculentus* L.), coriander (*Coriandrum sativum* L.), chilli (*Capsicum annum* L.), onion (*Allium cepa* L.k), green mulch crop (*Sesbania bispinosa* (Jacq.) W.F. Wight) and weeds (*Lagascea mollis* Cavanilles; Hindu datura (*Datura metel* L.) in eight districts of Andhra Pradesh (Adilabad, Anantapur, Karimnagar, Kurnool, Mahbubnagar, Medak, Warangal and Hyderabad), India (Fig. 1 and 2).

Sampling sites were chosen based on crops' flowering because adults of *H. armigera* prefer oviposition on flowers and flowering buds where the resulting larvae feed on the flowers and fruiting structures (Zalucki *et al.*, 1986; Jallow and Zalucki, 1997). Collection of larval samples were individually confined in glass tubes (80x20 mm) supplemented with tender pods and fruiting parts of the same host plants where from they were collected till pupation. Data was recorded on the duration of pupal period, sex and the date of adult emergence. The average critical day length at the sampling sites of the districts is shown in Fig. 3. Light traps were placed in ICRISAT fields to monitor the adult moths with an objective of determining the seasonal population dynamics as well as immigration




Fig. 2: Seasonal abundance of *H. armigera* sampled from different host crops and weeds grown during rainy, winter and summer seasons of Andhra Pradesh, India

Table 1: Mean meteorological parameters recorded in Andhra Pradesh, India (1974-96)

Month	Rainy days (no)	Rainfall (mm)	Relative humidity (%)	Temperature (°C)	
				Min.	Max.
June	10	132.0	79.7	23.6	33.6
July	15	200.1	87.6	22.5	30.2
August	15	196.2	89.1	22.1	29.1
September	12	134.1	89.8	21.1	30.0
October	8	115.0	88.1	19.7	30.1
November	3	13.7	87.8	16.3	28.5
December	1	1.0	87.5	13.1	27.7
January	1	10.8	85.6	14.2	28.6
February	4	5.7	73.6	16.9	31.9
March	2	11.9	64.3	20.0	35.7
April	4	30.2	57.9	23.2	38.0
May	4	26.2	56.2	25.4	39.1




Fig. 3: Average critical day lengths recorded during June-May in Andhra Pradesh, India

influx. In addition, observations were recorded at weekly intervals on the agrometeorological data with monthly averages of maximum and minimum temperatures, humidity, rainfall and rainy days (Table 1). The retention of pigmented eyespots in the pupal stage was considered as a diagnostic feature of diapause (Shumakov and Yakhimovich, 1955; Phillips and Newsom, 1966), which usually coincided at >35 days prior to adult emergence.

**Statistical analysis:** The replicated data was pooled together and means and standard errors were calculated. All data were analyzed by Analysis of Variance (ANOVA) to compare the significance using SPSS (Version 11.5).

## RESULTS

The critical day length triggering higher than normal percentage of pupae at 13 h in June-July, 11.5 h in December and >12 h in March-May (Fig. 3). A smaller proportion of *H. armigera* pupae entered diapause in winter during late November to early-December. Significantly higher moth emergence was observed in 7<sup>th</sup> and 8<sup>th</sup> week followed by 6<sup>th</sup> and 11<sup>th</sup> week. However, no significant differences were observed in adult moth emergence between 6<sup>th</sup> and 8<sup>th</sup> week. No emergence of adult moth was recorded from 19<sup>th</sup> week. A continuous trend of increase in pupal

period was recorded throughout the experimental period. However, significantly greater pupal period was observed in 23<sup>rd</sup> and 24<sup>th</sup> week onwards (Fig. 4). There was a greater significance in adult emergence in 23<sup>rd</sup> and 24<sup>th</sup> week followed by 27<sup>th</sup>, 26<sup>th</sup> and 28<sup>th</sup> week. Higher pupal period was observed in 26<sup>th</sup> week; however, there was no significant difference in pupal period in 24<sup>th</sup>, 25<sup>th</sup> and 27<sup>th</sup> week. Pupal period was lowest in 20<sup>th</sup> week (Fig. 5). In Andhra Pradesh, tomato is cultivated throughout the year; cotton and pigeon pea in June-March; okra in July-March; cowpea, mungbean, pearl millet, soybean, groundnut and sunflower crops in both June-July and Dec/Jan-March; coriander in August-April; sorghum in June-February; and chillies in August-December; safflower and chickpea in winter; and with supplemental irrigation in summer and *D. metel* and *L. mollis* Cav. Grow both in rainy and summer (March-May) seasons (Fig. 2).

Significantly greater no. of adult moths was caught between 45<sup>th</sup> to 52<sup>nd</sup> weeks, followed by 34<sup>th</sup>-39<sup>th</sup> week. Moth catches were observed throughout the experimental period. The larval population of *H. armigera* appeared in two peaks each moderately at 26<sup>th</sup> and 29<sup>th</sup> week in the rainy season and larger peaks at 46<sup>th</sup> and 48<sup>th</sup> week in winter season and a single smaller peak at week 21-in the summer season (Fig. 6). During cyclonic years in 1977-78 and 1995-96, a higher proportion of pupae of 15.11 and 17.64% entered diapause, respectively (Table 2). Among the diapause population, the total adult emergence was appeared in the ratio of 3:2 (648♂:419♀) during January to early-June. The staggered pattern of adult emergence occurred mostly between mid-February and March, very few in April, but only two adults each in May and June, respectively. During summer (March-April), a smaller proportion of pupae (2.00-10.43%) entered pupal diapause followed by an adult emergence ratio of 3:1 (35♂:11♀) between June and mid-July (Table 3).

Pupal recovery and diapausing pupae during winter diapause were significantly higher on pigeonpea than on chickpea (Fig. 7A and B). Significantly higher number




Fig. 4: Pupal period and adult moth emergence from winter diapausing pupae of *H. armigera* in Andhra Pradesh, India (1974-96)




Fig. 5: Pupal period and adult moth emergence from summer diapausing pupae of *H. armigera* in Andhra Pradesh, India (1974-96)




Fig. 6: Mean recovery of larvae and light trap catches of adult moths of *H. armigera* in Andhra Pradesh, India (1974-96)

Table 2: Incidence of winter diapause in the field samplings of *H. armigera* on leguminous crops grown in Andhra Pradesh, India (1974-96)


Year	Period of sampling	Crop	Pupal recovery (no) ±S.E.	Diapause pupae		Adult emergence (no.)		
				No.	(%)	♂	♀	Ratio
1974-75	Nov 15-Dec 3	Chickpea	335±2.51 <sup>f</sup>	5 <sup>c</sup>	1.49	1 <sup>c</sup>	4 <sup>c</sup>	1:4
1975-76	Dec 1-5	Pigeonpea	934±2.40 <sup>cd</sup>	14 <sup>c</sup>	1.49	10 <sup>c</sup>	4 <sup>c</sup>	5:2
1976-77	Nov 29-Dec 6	Pigeonpea	1330±1.53 <sup>c</sup>	2 <sup>c</sup>	0.21	1 <sup>c</sup>	1 <sup>c</sup>	1:1
1977-78*	Dec 1-12	Pigeonpea	3500±7.50 <sup>a</sup>	529 <sup>a</sup>	15.11	319 <sup>a</sup>	210 <sup>a</sup>	3:2
1978-79	Dec 1-10	Pigeonpea	518±3.60 <sup>de</sup>	10 <sup>c</sup>	1.93	6 <sup>c</sup>	4 <sup>c</sup>	3:2
1980-81	Dec 3-13	Pigeonpea	435±1.73 <sup>de</sup>	6 <sup>c</sup>	1.38	4 <sup>c</sup>	2 <sup>c</sup>	2:1
1981-82	Dec 10-16	Pigeonpea	216±4.04 <sup>f</sup>	4 <sup>c</sup>	1.85	2 <sup>c</sup>	2 <sup>c</sup>	1:1
1982-83	Nov 25-Dec 7	Pigeonpea	400±1.73 <sup>de</sup>	10 <sup>c</sup>	2.50	6 <sup>c</sup>	4 <sup>c</sup>	3:2
1984-85	Dec 20-Jan 4	Chickpea	284±1.15 <sup>f</sup>	4 <sup>c</sup>	1.41	3 <sup>c</sup>	1 <sup>c</sup>	3:1
1990-91	Dec 2-13	Pigeonpea	300±1.52 <sup>f</sup>	9 <sup>c</sup>	3.00	6 <sup>c</sup>	3 <sup>c</sup>	2:1
1991-92	Nov 24-Dec 3	Pigeonpea	205±0.57 <sup>f</sup>	7 <sup>c</sup>	3.41	5 <sup>c</sup>	2 <sup>c</sup>	5:2
1992-93	Dec 4-12	Pigeonpea	274±3.05 <sup>f</sup>	6 <sup>c</sup>	2.19	4 <sup>c</sup>	2 <sup>c</sup>	2:1
1995-96*	Dec 24-Jan 7	Pigeonpea	2729±3.46 <sup>b</sup>	461 <sup>b</sup>	17.64	286 <sup>b</sup>	175 <sup>b</sup>	3:2

Values (Mean±SEM) carrying same alphabet (s) within a column are not significantly different by Duncan's multiple range test, (p≤0.05); \*: Cyclonic years

Table 3: Incidence of summer diapause in the field sampling of *H. armigera* on different crops and weeds grown in Andhra Pradesh, India (1974-96)

Year	Period of sampling	Crop/weed	Pupal recovery (no) ±S.E.	Diapause pupae		Adult emergence (no.)		
				No.	(%)	♂	♀	Ratio
1974-75	April 15-18	Maize	30±2.51 <sup>d</sup>	1 <sup>bc</sup>	3.33	1 <sup>c</sup>	0 <sup>a</sup>	1:0
1974-75	Feb 18-20	Chickpea	70±1.73 <sup>c</sup>	2 <sup>bc</sup>	2.86	1 <sup>c</sup>	1 <sup>a</sup>	1:1
1977-78	April 10-13	Maize	45±1.52 <sup>d</sup>	1 <sup>bc</sup>	2.22	0 <sup>c</sup>	1 <sup>a</sup>	0:1
1978-79	April 4-7	Datura	65±1.53 <sup>c</sup>	5 <sup>b</sup>	7.69	4 <sup>b</sup>	1 <sup>a</sup>	4:1
1981-82	April 15-19	Chickpea	40±1.00 <sup>d</sup>	4 <sup>b</sup>	10.00	3 <sup>b</sup>	1 <sup>a</sup>	3:1
1984-85	April 4-9	Datura	80±1.53 <sup>c</sup>	7 <sup>ab</sup>	8.75	5 <sup>ab</sup>	2 <sup>a</sup>	5:2
1990-91	March 17-21	Chickpea	150±2.88 <sup>a</sup>	3 <sup>b</sup>	2.00	2	1 <sup>a</sup>	2:1
1990-91	March 12-16	Pigeonpea	108±0.99 <sup>b</sup>	3 <sup>b</sup>	2.78	3 <sup>b</sup>	0 <sup>a</sup>	3:0
1991-92	March 10-15	Chickpea	115±1.00 <sup>b</sup>	12 <sup>a</sup>	10.43	9 <sup>a</sup>	3 <sup>a</sup>	3:1
1992-93	April 13-17	Maize	50±2.51 <sup>d</sup>	3 <sup>b</sup>	6.00	3 <sup>b</sup>	0 <sup>a</sup>	3:0
1995-96	March 15-19	Chickpea	75±2.64 <sup>cde</sup>	5 <sup>b</sup>	6.67	4 <sup>b</sup>	1 <sup>a</sup>	4:1

Values (Mean±SEM) carrying same alphabet (s) within a column are not significantly different by Duncan's multiple range test, (p≤0.05)






Fig. 7: Pupal recovery and incidence of winter and summer diapause and sex ratio of adult emergence in *H. armigera* on different crops and weed in Andhra Pradesh, India (1974-96)




Fig. 8: Non diapausing pupal period of *H. armigera* collected from various host crops and weeds in Andhra Pradesh, India (1974-96)

of males than females was recorded on pigeonpea, while no significant difference was observed between number of males and females on chickpea (Fig. 7C). Among different crops in summer diapause, significantly higher pupal recovery was observed on chickpea and pigeonpea than on *Datura* and maize (Fig. 7D). *Datura* showed significantly higher pupal diapause during summer than pigeonpea and maize. However, no significant difference was recorded in pupal diapause between chickpea and maize (Fig. 7E). Pigeonpea showed the lowest pupal diapause. Significantly greater difference in sex ratio during summer diapause was observed in *Datura*, followed by chickpea and pigeonpea (Fig. 7F). There was no significant difference in pupal period between non diapausing males and females across the year. However, pupal period was significantly higher in December followed by January as compared to the rest of the months. The nondiapausing adult moth emergence was observed throughout the 12 months (Fig. 8).

## DISCUSSION

The incidence of diapause in *H. armigera* during winter and summer is governed by agroclimatic factors. The multi-year data on the patterns of winter and summer diapause development and emergence correlated with the field activity of adult moth

emergence is important in better understanding the role of local population dynamics in the ecology and genetics of *H. armigera*. Present investigation revealed that shorter day length and cooler temperatures in November-December and higher temperatures in March-April are influencing the pupae to enter diapause. The summer diapause can significantly impact the rainy season crops by early emergence of adult populations resulting in considerable yield losses, while the winter diapause doesn't impact the crop damage to same degree because there is already a resident population multiplying at that time. However, there is a possibility of increased population explosion in winter crops if the timing of the cyclonic weather events in Andhra Pradesh coincides with the fruiting and podding stages in legume and oilseed crops and boll formation in cotton. The continuous emergence of moths from winter diapause pupae is a mechanism to ensure increasing their population by breeding with the emerging local population during late-January to April. Further, drought conditions associated with low local populations in the region have greater significance with diapause populations for the overall population dynamics. It was also observed by field sampling of larvae and pupae as well as monitoring of adult catches in light traps indicated that a majority of the pupae of *H. armigera* did not enter winter diapause.

In general, the incidence of diapause in winter and summer was significantly higher not only in male than female pupae but also in the sex ratio for adult emergence. The pupal period of non-diapause populations remain active throughout the year and migrate to crops that are planted in the neighboring areas, which are favorable for their suitability for population buildup. This was supported by the light trap catches of adult moths during September, December and March. Induction and termination of diapause was influenced in both winter and summer by decreased day length and daily temperatures (Mansingh, 1971; Tauber and Tauber, 1976; Masaki, 1980; Tauber *et al.*, 1986; Danks, 2005). However, at extreme temperatures (>35°C) exceeded by more than 2 months in Australia (Dillon, 1998) and China (Wu and Gu, 1997c). Larval rearing of *H. armigera* induced diapause at 18°C and 27°C with a range in pupal duration of 76-103 and 123-

49 days, respectively in the winter and summer seasons (Nel, 1961; Roome, 1979). In general, the winter diapause was induced in <7.7% population of pupae sampled on maize, pigeonpea, cotton, tobacco and field bean (*Dolichos lablab*), which continued over a period of 30-171 days at 20°C in Ukiriguru (Tanzania) (Reed, 1965). In contrast, no diapause has been reported in *H. armigera* populations in Namulonge (Uganda), with an exception of two pupae collected on sunflower entered diapause over a period of 53 and 84 days (Coaker, 1959) and from tropical and sub-tropical regions at room temperatures during winter day lengths (Hardwick, 1965). In contrast, under similar conditions, the pupal diapause was 8% in Rehovot (Israel; 32°N), 19% in Rotorua (New Zealand; 38°S) and 82% at Nelson (New Zealand; 41°S). Similarly, no diapause has been reported in the larvae of *H. zea* reared at 27°C with a day length of 10 h in Louisiana (USA) (30°N) (Phillips and Newsom, 1966), but 8% of *H. armigera* pupae entered diapause when reared at 25.6°C and 10 h day length in Azerbaidzhan SSR (40°N) and Ukrainian SSR (47°C) (Komarova, 1959). Adult emergence of *H. armigera* took shorter period of 35 days at 30°C but increased its duration by 4 days at 25°C and 111 days at 20°C in Stavropol (45°N) (Kuznetsova, 1972). On the other hand, the pupae took 152 days at 20°C; 80 days at 25°C; 73 days at 30°C in Tadzhikistan (40°N) (Kuznetsova, 1972); and 70 days at 35 and 20°C, respectively in Japan (34°N) (Qureshi *et al.*, 1999).

The critical day length for induction of diapause ranged between 12L: 12D and 14L: 10D during the larval stage in *H. armigera*, but the termination took a longer period without any latitudinal variation at 20°C (Qureshi *et al.*, 2000). In contrast, the diapause in *H. zea* was associated with the latitude from tropical regions and summer diapause at 37°C in Burkina Faso (Nibouche, 1998). While the developmental period of *H. armigera* pupae was prolonged with increased latitude but shortened at low temperatures in central and northern China (Wu and Guo, 1997 b, c, d).

Frequent outbreaks of *H. armigera* have been reported in late-fruiting cultivars, larger fields and extended growing season of cotton, pigeonpea and chickpea crops. Late-flowering/fruiting cultivars usually favored large over-wintering populations of *H. armigera* resulting in predisposed infestation in subsequent crop season. Temperature and photoperiod although considered as key stimuli for the induction and termination of diapause in *H. armigera*, increased latitude also played a significant role. Diapausing pupae have the ability to survive during winter in southern Japan (Qureshi *et al.*, 1999, 2000; Izumi *et al.*, 2005). In Western Tanganyika, a greater proportion of diapause pupae (45%) have been reported on maize and pigeon pea, followed by cotton (2.5%) and tobacco (2.3%) during April-July which period coincided with a decrease in day length (Reed, 1965). In south-eastern Australia, a significant proportion of *H. armigera*

overwinter as pupae during summer and with a high recovery on cotton (Duffield, 2004). In Australia, adult emergence during mid-October-mid-November had two weeks later in the southern region compared to northern parts of New South Wales (Duffield and Dillon, 2005). A smaller proportion of pupae entered in winter diapause in Tanzania (Reed, 1965), Rhodesia (Jones, 1937), South Africa (Parsons, 1939; Nel, 1961), Russia (Komarova, 1959) and Botswana (Roome, 1979). In general, the adult emergence was more rapid from the summer diapausing pupae compared to winter ones. Higher incidence of diapause was induced not only by low temperatures but also by photoperiod in *H. zea* (Pullen *et al.*, 1992), *H. punctigera* (Cullen and Browning, 1978), *H. virescens* (Henneberry, 1994) and *H. armigera* (Qureshi *et al.*, 1999). Despite variation in photoperiodic conditions the Okayama population of *H. armigera* did not enter diapause at 30°C (Li and Xie, 1983). Present results also indicated that a small proportion of *H. armigera* pupae entered winter diapause representing a bimodal-shape curve that has coincided with the cyclonic years. Similar trend in bimodal emergence influenced by fluctuating environments has been reported in *H. zea* (Waldbauer, 1978). In addition, long-term studies of summer diapause coincided with maximum temperatures ranged between 32° and 35°C by mid-March associated with an adult emergence in 70-115 days in a bimodal fashion. Similar bimodal emergence of adults has been reported for *Chloridea obsoleta* in USSR (Danilevski, 1965), tobacco budworm (*Protoparce sexta*) (Rabb, 1966) and *H. virescens* in the USA (Butler *et al.*, 1985). Long-duration of adult emergence was also recorded in *H. armigera* in Sudan (Hackett and Gatehouse, 1982) and India (Tripathi and Singh, 1993). Geographic variation in photoperiod and temperature sensitivity to *H. armigera* populations for diapause was 40 and 70% in Japan and China respectively at 25°C and 10L: 14D (Li and Xie, 1983; Wu *et al.*, 1997). Present investigation also showed a similar tendency of entering diapause by a greater proportion of male pupae than females. Similar observations were made in *H. virescens* (Butler *et al.*, 1985) and *H. armigera* (Wu and Guo, 1997a).

Summer diapause in *H. virescens* was associated not only with high temperatures (>32°C) (Butler *et al.*, 1985) but in low rainfall years (Schneider, 2003). In Sudan, the duration of diapause extended to more than 2 months at low temperatures with fewer pupae of *H. armigera* still continued to remain in diapause at high temperatures and commenced development only when the temperatures was lowered (Hackett and Gatehouse, 1982). Present studies also indicated that diapause terminates more rapidly at higher temperatures. These results are in consonance with earlier reports (Mangat and Apple, 1966; Roach and Adkisson, 1970; Kuznetsova, 1972). Higher incidence of diapause in

winter during October-December and summer diapause in April-May was reflected by the recovery of pupae from the crops such as maize, pigeonpea, chickpea and *Datura*. This situation contributed to increased population buildup that led to severe infestation in the succeeding crops.

Further, the large-scale appearance of *H. armigera* larvae is indicative of greater fitness and faster development. In *H. virescens*, the photoperiod however had a negligible effect on the induction of summer diapause (Butler *et al.*, 1985). Early emergence of moths from summer diapausing pupae in June-July was associated with a break in rainfall after a prolonged dry season. The wild and volunteer crop plants also played a significant role early in the season for the population buildup of *H. armigera*. Although the populations are initially low they dispersed widely by maintaining optimal populations until suitable crops become available in the monsoon season where the major agricultural diversity begins. Current studies provide a basis in understanding of *H. armigera* population dynamics for their impact on rainy season crops due to summer diapause in southern India. From these studies, it can be inferred that diapause is an adaptive trait in *H. armigera* in the semi-arid tropics and its occurrence is maintained by natural selection and not by intrusion of genes from elsewhere.

### CONCLUSION

The cotton bollworm, *H. armigera* is a major pest of agricultural crops in the semi-arid regions of Indian sub-continent. The long term studies suggest the pupae undergoes facultative diapause and remains concern with the sudden appearance of larvae in the beginning of the monsoon crops. The long-term studies give the pests ability to survive the favorable and adverse weather conditions and be pestiferous. It is also recommended for the farming community to deep plough the soil where summer diapausing pupae are buried, so they could be killed which may result in lesser population build-up and could prevent out-breaks in cyclonic conditions. In untilled land, increased survival of pupae may allow the buildup of early populations.

### ACKNOWLEDGMENT

Financial support was provided in 1974-88 by the Farming Systems Research Program of ICRISAT Cropping Systems Entomology and 1991-96 by the Natural Resources Institute, UK, through adaptive research of the UK Government's Department of International Development (DfID). Authors are grateful to Mr. K.V.S. Satyanarayana, M. Satyanarayana and G. Venkateswarlu for technical assistance.

### REFERENCES

- Armes, N.J., D.R. Jadhav and K.R. De Souza, 1996. A survey of insecticide resistance in *Helicoverpa armigera* in the Indian subcontinent. Bull. Entomol. Res., 86: 499-514.
- Beck, S.D., 1968. Insect Photoperiodism. 1st Edn., Academic Press, New York, pp: 288.
- Beck, S.D., 1980. Insect Photoperiodism. 2nd Edn., Academic Press, New York, pp: 387.
- Bhatnagar, V.S. and J.C. Davies, 1978. Factors affecting populations of gram pod borer, *Heliothis armigera* (Hübner) (Lepidoptera: Noctuidae) in the period 1974-77 at Patancheru (Andhra Pradesh). Bull. Entomol., 19: 52-64.
- Borchert, D.M., R.D. Magarey and G.A. Fowler, 2003. Pest assessment: Old World Bollworm, *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae). USDA-APHIS-PPQ-CPHST-PERAL/NCSU. NAPPFAST, May 27, pp: 9.
- Butler, J.G.D., L.T. Wilson and T.J. Henneberry, 1985. *Heliothis virescens* (Lepidoptera: Noctuidae): Initiation of summer diapause. J. Econ. Entomol., 78: 320-324.
- Casimero, V., R. Tsukuda, F. Nakasuji and K. Fujisak, 2000. Effect of larval diets on the survival and development of larvae in the cotton bollworm, *Helicoverpa armigera* Hübner (Lepidoptera: Noctuidae). Appl. Entomol. Zool., 35: 69-74.
- Coaker, T.H., 1959. Investigations on *Heliothis armigera* (Hübner) in Uganda. Empire Cotton. Bull. Entomol. Res., pp: 50: 487-506.
- Cullen, J.M. and T.O. Browning, 1978. The influence of photoperiod and temperature on the induction of diapause in pupae of *Heliothis punctigera*. J. Insect Physiol., 24: 595-601.
- Daly, J.C. and D.A.H. Murray, 1988. Evolution of resistance to pyrethroids in *Heliothis armigera* (Hübner) (Lepidoptera: Noctuidae) in Australia. J. Econ. Entomol., 81: 984-988.
- Danilevski, A.S., 1965. Photoperiodism and Seasonal Development of Insects. Oliver and Boyd. Edinburgh, UK.
- Danks, H.V., 1987. Insect Dormancy: An Ecological Perspective. Biological Survey, Ottawa, Canada.
- Danks, H.V., 2005. Key themes in the study of seasonal adaptations in insects I. Patterns of cold hardiness. Appl. Entomol. Zool., 40: 199-211.
- Denlinger, D.L., 1986. Dormancy in tropical insects. Ann. Rev. Entomol., 31: 239-264.
- Denlinger, D.L., 2002. Regulation of diapause. Ann. Rev. Entomol., 47: 93-122.
- DeSouza, K., J. Holt and J. Colvin, 1995. Diapause, migration and pyrethroid-resistance dynamics in the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae). Ecol. Entomol., 20: 333-342.

- Dillon, M., 1998. Predicting autumn diapause induction in *Helicoverpa* using long-term average temperatures. Proceeding of the 9th Austr Cotton Conference, Broad Beach, Queensland, pp: 475-479.
- Duffield, S.J., 2004. Evaluation of the risk of overwintering *Helicoverpa* spp. pupae under irrigated summer crops in south-eastern Australia and the potential for area-wide management. Ann. Appl. Biol., 144: 17-26.
- Duffield, S.J. and M.L. Dillon, 2005. The emergence and control of overwintering *Helicoverpa armigera* pupae in southern New South Wales. Austr. J. Entomol., 44: 316-320.
- Eger, J.E., J.A. Witz, A.W. Hartstack and W.L. Sterling, 1982. Survival of pupae of *Heliothis virescens* and *Heliothis zea* (Lepidoptera: Noctuidae) at low temperatures. Canad. Entomol., 114: 289-301.
- Farrow, R.A. and J.C. Daly, 1987. Long-range movements as an adaptive strategy in the genus *Heliothis* (Lepidoptera: Noctuidae): A review of its occurrence and detection in four pest species. Aust. J. Zool., 35: 1-24.
- Fitt, G.P., 1989. The ecology of *Heliothis* species in relation to agroeco systems. Annu. Rev. Entomol., 34: 17-52.
- Ge, F., F. Chen, M.N. Paralulee and E.N. Yardim, 2005. Quantification of diapausing fourth generation of *Helicoverpa* spp. (Lepidoptera: Noctuidae) during autumn in northern China. Entomol. Exp. Appl., 116: 1-7.
- Hackett, D.S. and A.G. Gatehouse, 1982. Diapause in *Heliothis armigera* (Hübner) and *H. fletcheri* (Hardwick) (Lepidoptera: Noctuidae) in the Sudan Gezira. Bull. Entomol. Res., 72: 409-422.
- Hamamura, T., 1998. Recent occurrence and injury of *Heliothis armigera*-From a questionnaire investigation. Pl. Prot. (Japan), 52: 407-413.
- Hardwick, D.F., 1965. The corn earworm complex. Memoirs Entomol. Soc. Canada, 40: 247.
- Henneberry, T.J., 1994. Effects of temperature on tobacco budworm (Lepidoptera: Noctuidae) pupal diapause, initiation and final stage of movement of stemmatal eyespots and adult emergence. Southwest. Entomol., 19: 329-333.
- ICRISAT (International Crops Research Institute for the Semi-Arid Tropics), 1978. Annual Report, 1977-78. International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru andhra Pradesh 502 324, India.
- Izumi, Y., K. Anniwaer, H. Yoshida, S. Sonoda, K. Fujisaki and H. Tsumuki, 2005. Comparison of cold hardiness and sugar content between diapausing and nondiapausing pupae of the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae). Physiol. Entomol., 40: 36-41.
- Jallow, M.F.A. and M.P. Zalucki, 1997. Within and between population variation in host plant preference and specificity in Australian *Helicoverpa armigera* (Lepidoptera: Noctuidae). Aust. J. Zool., 44: 503-519.
- Jones, E.P., 1937. The overwintering pupa of *Heliothis armigera* Hubn. (*Obsolete* Fabr.). I. Effect of temperature and moisture. Publ. Brit. A. Afr. Co. No., 6A: 19-36.
- Kay, I.R., 1982. The incidence and duration of pupal diapause in *Heliothis armigera* (Hübner) (Lepidoptera: Noctuidae) in southeast Queensland. J. Aust. Entomol. Soc., 21: 263-266.
- King, A.B.S., 1994. *Heliothis/Helicoverpa* (Lepidoptera: Noctuidae). In: Matthews, G.A. and I.P. Tunstall (Eds.), Insect Pests of Cotton. CAB International, Oxon, UK, pp: 39-106.
- Komarova, O.S., 1959. On the conditions determining the diapause of the hibernating pupae of *Chloridea obsoleta* F. (Lepidoptera: Noctuidae). Entomol. Rev. (Moscow), 38: 352-360.
- Kurban, A., H. Yoshida, Y. Izumi, S. Sonoda and H. Tsumuki, 2007. Pupal diapause of *Helicoverpa armigera* (Lepidoptera: Noctuidae): Sensitive stage for thermal induction in the Okayama (western Japan) population. Bull. Entomol. Res., 97: 219-223.
- Kuznetsova, M.S., 1972. The effects of temperature and photoperiodic conditions on the reactivation of diapausing pupae of the cotton bollworm, *Chloridea obsoleta* F. (Lepidoptera: Noctuidae). Entomol. Rev. (Moscow), 51: 311-315.
- Li, C. and B.Y. Xie, 1983. Combined effects of photoperiod and temperature on the diapause of *Heliothis armigera* (Hb.). Insect Knowl., 18: 58-61.
- Mangat, B.S. and J.W. Apple, 1966. Termination of diapause in corn earworm. Proc. North Central Branch, Entomol. Soc. Amer., 21: 123-125.
- Mansingh, A., 1971. Physiological classification of dormancies in insects. Canad. Entomol., 103: 983-1009.
- Martinat, P.J., 1987. The Role of Climatic Variation and Weather in Forest Insect Outbreaks. In: Barbosa, P. and J.C. Schultz (Eds.), Insect Outbreaks. Academic Press, New York, pp: 241-268.
- Masaki, S., 1980. Summer diapause. Annu. Rev. Entomol., 25: 1-25.
- Masaki, S., 2002. Ecophysiological consequences of variability in diapause intensity. Europ. J. Entomol., 99: 143-154.
- Mironidis, G.K., D.C. Stamopoulos and M. Savopoulou-Soultani, 2010. Overwintering survival and spring emergence of *Helicoverpa armigera* (Lepidoptera: Noctuidae) in Northern Greece. Environ. Entomol., 39: 1068-1084.

- Murray, D.A.H., 1991. Investigations into the development and survival of *Heliothis* spp. pupae in South-East Queensland. Ph.D. Thesis, University of Queensland, Brisbane, Australia.
- Nel, J.J.C., 1961. The seasonal history of *Heliothis armigera* (Hübner) on lupins in the South Western Cape Province. Sth. Afr. J. Agric. Sci., 4: 575-588.
- Nibouche, S., 1998. High temperature induced diapause in the cotton bollworm *Helicoverpa armigera*. Entomol. Exptl. Applic., 87: 271-274.
- Parsons, F.S., 1939. Investigations on the cotton bollworm, *Heliothis armigera* (Hübner). (*obsoleta* Fabr.). Part I. The annual march bollworm incidence and related factors. Bull. Entomol. Res., 30: 321-338.
- Phillips, J.R. and L.D. Newsom, 1966. Diapause in *Heliothis zea* and *Heliothis virescens* (Lepidoptera: Noctuidae). Ann. Entomol. Soc. Amer., 59: 154-159.
- Potter, M.F. and T.F. Watson, 1980. Induction of diapause in the tobacco budworm in Arizona. J. Econ. Entomol., 73: 820-823.
- Pullen, S.R., R.W. Meola and J.D. Lopez, 1992. Temperature as a sole factor for diapause induction after pupation in *Helicoverpa zea* (Lepidoptera: Noctuidae). Environ. Entomol., 21: 1404-1409.
- Qureshi, M.H., T. Murai, H. Yoshida, T. Shiraga and H. Tsumuki, 1999. Effects of photoperiod and temperature on the development and diapause induction in the Okayama population of *Helicoverpa armigera* (Hb.) (Lepidoptera: Noctuidae). Appl. Entomol. Zool., 34: 327-331.
- Qureshi, M.H., T. Murai, H. Yoshida and H. Sumuki. 2000. Populational variation in diapause-induction and termination of *Helicoverpa armigera* (Hb.) (Lepidoptera: Noctuidae). Appl. Entomol. Zool., 35: 357-360.
- Rabb, R.L., 1966. Diapause in *Protoparce sexta* (Lepidoptera: Sphingidae). Ann. Entomol. Soc. Amer., 59: 160-165.
- Rao, K.T., N. Venugopal Rao and A. Satyanarayana, 1990. Pigeon pea pod borer, *Helicoverpa armigera* incidence in relation to the rainfall at Lam farm, Guntur andhra Pradesh, India, 1977-78. Intl. Pigeonpea Newsl., 3: 97-98.
- Reed, W., 1965. *Heliothis armigera* (Hb.) (Noctuidae) in Western Tanganyika. I. Biology with special reference to the pupal stage. Bull. Entomol. Res., 56: 117-125.
- Reed, W. and C.S. Pawar, 1982. *Heliothis*: A global problem. Proceeding of the International Workshop on *Heliothis* Management, 15-20 November 1981, ICRIAT Center, Patancheru andhra Pradesh, India Ed. By Reed W, Kumble V, Patancheru andhra Pradesh, India: International Crops Research Institute for the Semi-Arid Tropics. pp: 9-14.
- Roach, S.H. and P.L. Adkisson, 1970. Role of photoperiod and temperature in the induction of pupal diapause in the bollworm, *Heliothis zea*. J. Insect. Physiol., 16: 1591-1597.
- Roome, R.E., 1979. Pupal diapause in *Heliothis armigera* (Hübner) (Lepidoptera: Noctuidae) in Botswana: its regulation by environmental factors. Bull. Entomol. Res., 69: 149-160.
- Schneider, J.C., 2003. Overwintering of *Heliothis virescens* (F.) and *Helicoverpa zea* (Boddie) (Lepidoptera: Noctuidae) in cotton fields of Northeast Mississippi. J. Econ. Entomol., 96: 1433-1447.
- Shanower, T.G., J. Romeis and E.M. Minja, 1999. Insect pests of pigeonpea and their management. Annu. Rev. Entomol., 44: 77-96.
- Shimuzi, K. and K. Fujisaki, 2002. Sexual differences in diapause induction of the cotton bollworm, *Helicoverpa armigera* (Hb.) (Lepidoptera: Noctuidae). Appl. Entomol. Zool., 37: 527-533.
- Shimuzi, K., K. Shimizu and K. Fujisaki, 2006. Timing of diapause induction and overwintering success in the cotton bollworm *Helicoverpa armigera* (Hb.) (Lepidoptera: Noctuidae) under outdoor conditions in temperate Japan. Appl. Entomol. Zool., 41: 151-159.
- Shumakov, E.M. and L.A. Yakhimovich, 1955. Morphological and histological peculiarities of the metamorphosis of the cotton bollworm (*Chloridea obsoleta* F.) in connection with the phenomenon of diapause. Dokl. Acad. Sci. (USSR), 101: 379-382.
- Tauber, M.J. and C.A. Tauber, 1976. Insect seasonality: Diapause maintenance, termination and post-diapause development. Annu. Rev. Entomol., 21: 81-107.
- Tauber, M.J., C.A. Tauber and S. Masaki, 1986. Seasonal Adaptations of Insects - Special Cases. Oxford University Press, New York, pp: 161-191.
- Tripathi, S.R. and R. Singh, 1993. Seasonal bionomics of *Heliothis armigera* (Hübner) (Lepidoptera: Noctuidae) in the terai belt of northeastern Uttar Pradesh. Insect Sci. Applic., 4: 439-444.
- Waldbauer, G.P., 1978. Phenological Adaptation and the Polymodal Emergence Patterns of Insects. In: Dingle, H., (Ed.), Evolution of Insect Migration and Diapause. Springer-Verlag, New York, USA, pp: 127-144.
- Wilson, A.G.L., T. Lewis and R.B. Cunningham, 1979. Overwintering and spring emergence of *Heliothis armigera* (Hübner) in the Namoi Valley, New South Wales. Bull. Entomol. Res., 69: 97-109.
- Wu, K. and Y Guo, 1997a. Inheritance experiment of cotton bollworm populations from Xinjiang Uygur autonomous region and Hunan Province in China. Acta Entomol. Sin., 40: 20-24.

- Wu, K. and Y. Guo, 1997b. Study on the diapause conditions of different geographical populations of cotton bollworm (*Helicoverpa armigera*) in China. China Agricultural Sciencetech Press, Institute of Plant Protection, Chinese Acad. Agric. Sci., Beijing, China, pp: 109-114.
- Wu, K. and Y. Guo, 1997c. The characteristics of diapause termination of different geographical populations in cotton bollworm, *Helicoverpa armigera* (Hübner) in China. Acta Entomol. Sin., 40: 25-29.
- Wu, K. and Y. Guo, 1997d. The influences of soil moisture content on emergence and cold hardiness of different geographical populations of cotton bollworm. Acta. Phytophyl. Sin., 24: 142-146.
- Wu, K., G. Yuyuan, W. Jianfu and S. Fuzai, 1997. On the cold hardiness of cotton bollworm, *Helicoverpa armigera* (Hübner). Acta Ecol. Sin., 17: 298-302.
- Zalucki, M.P., G. Darglish, S. Firempong and P.H. Twine, 1986. The biology and ecology of *Heliothis armigera* (Hübner) and *H. punctigera* (Wallengren) (Lepidoptera: Noctuidae) in Australia: What do we know? Austr. J. Zool., 34: 779-814.
- Zalucki, M.P., A.H. Murray, P.C. Gregg, G.P. Fitt, P.H. Twine and C. Jones, 1994. Ecology of *Helicoverpa armigera* (Hubner) and *H. punctigera* (Wallengren) in the Inland of Australia: Larval sampling and host plant relationships during winter and spring. Austr. J. Zool., 42: 329-346.
- Zhou, X., S.W. Applebaum and M. Coll, 2000. Overwintering and spring migration in the bollworm *Helicoverpa armigera* (Lepidoptera: Noctuidae) in Israel. Environ. Entomol., 29: 1289-1294.