

Marker-assisted improvement of grain protein content and grain weight in Indian bread wheat

Manish K. Vishwakarma · B. Arun ·
V. K. Mishra · P. S. Yadav · H. Kumar ·
Arun K. Joshi

Received: 30 March 2015 / Accepted: 10 November 2015 / Published online: 20 November 2015
© Springer Science+Business Media Dordrecht 2015

Abstract The leading Indian wheat cultivar ‘HUW234’ produces grain with relatively low protein content (GPC) and thousand grain weight (TGW). Therefore, marker-assisted backcrossing was used to improve these two important traits by introducing favorable genes/alleles from cv. ‘Glu133’, which harbors both alleles for both GPC (*Gpc-B1*) and TGW. Foreground selection for GPC was achieved using microsatellite markers *Xucw108* linked with *Gpc-B1* and *Xgwm297* linked with TGW. Background selection applied to support recovery of the recurrent parent genotype was based on 86 genomically distributed microsatellites. A selected BC₂F₁ plant was the progenitor of 15 BC₂F₄ families, in which representation of the cv. HUW234 genome ranged

from 89.5 to 93.0 %. The new derivatives of HUW234 were comparable in yield to the original cultivar, but with both significantly higher GPC and heavier kernels.

Keywords Background selection · Foreground selection · Marker assisted selection · *Triticum aestivum*

Introduction

Wheat provides over 20 % of the calories in the human diet and 25 % of the protein (FAO 2012). After many years of increasing productivity, wheat yields have shown evidence of plateauing since the turn of the century (Lobell et al. 2011) while the human population continues to grow. Although yield is the primary trait under selection in most breeding programs and grain destined for human consumption is processed before consumption, end-use quality is also an important breeding objective. Among quality traits, grain protein content (GPC) and thousand grain weight (TGW) are significant quality traits. GPC is a major determinant of nutritional and market value of wheat grain as well as processing quality (Brevis et al. 2010). High GPC varieties not only address the issue of end-use quality, but also provide a relatively cheap source of protein. This is of high significance for the population dense North

Electronic supplementary material The online version of this article (doi:10.1007/s10681-015-1598-6) contains supplementary material, which is available to authorized users.

M. K. Vishwakarma · B. Arun · V. K. Mishra ·
P. S. Yadav · H. Kumar · A. K. Joshi
Department of Genetics and Plant Breeding, Institute of
Agricultural Sciences, Banaras Hindu University,
Varanasi, India

M. K. Vishwakarma
International Crops Research Institute for the Semi-Arid
Tropics (ICRISAT), Hyderabad, India

A. K. Joshi (✉)
CIMMYT, South Asia Regional Office, Singha Durbar
Plaza Road, Kathmandu, Nepal
e-mail: a.k.joshi@cgiar.org

Eastern Plains Zone (NEPZ) of India with a rice–wheat cropping system and with wheat having double the protein content of rice. TGW is correlated with flour yield as well as actual grain yield (GY) (Ketata et al. 1976; Campbell et al. 1999). Despite major advances in yield achieved by the Green Revolution, progress in increasing GPC has been slow (Gupta 2004). Although it is commonly reported that yield and GPC are negatively correlated (Simmonds 1995), there is little agreement on its biological basis (Cox et al. 1985; Brevis and Dubcovsky 2010).

A number of DNA-based markers for various quality-associated traits have been described in wheat (Gupta et al. 2011; Roder et al. 2008). Accession FA-15-3, which originated in Israel, accumulates high GPC (Avivi 1978) partly due to presence of the gene *Gpc-B1* (Joppa et al. 1997; Olmos et al. 2003). This gene has been tagged by a variety of DNA-based markers (Khan et al. 2000; Singh et al. 2001; Distelfeld et al. 2004, 2006; Olmos et al. 2003; Prasad et al. 2003; Kulwal et al. 2005; Wang et al. 2012) and was positionally cloned by Uauy et al. (2006). When *Gpc-B1* was introduced into low GPC cultivars, it showed little evidence of a yield penalty (Kade et al. 2005; Brevis and Dubcovsky 2010), and similar experiences were reported for Indian cultivar derivatives (Tyagi et al. 2014; Vishwakarma et al. 2014).

As an important component of GY, TGW is always a favored trait in wheat breeding programs (Gegas et al. 2010). QTLs on chromosomes 1A, 1B, 1D, 2B, 2D, 4B, 4D, 7A and 7D have been reported for this trait (Huang et al. 2003, 2004; McCartney et al. 2005; Wang et al. 2009; Mir et al. 2012). A major QTL (*QTgw.ipk-7D*) explaining 84.7 % of the phenotypic variation in TGW (Huang et al. 2003, 2004) was later fine-mapped by Roder et al. (2008). More recently, Mir et al. (2012) identified 11 new QTLs.

Marker assisted selection (MAS) has gained considerable attraction as a breeding tool in crop improvement (Gupta et al. 2010; Vishwakarma et al. 2014). Gupta et al. (2010) estimated that at least 60 genes/quantitative trait loci (QTL) have been used in wheat improvement programs, largely (but not exclusively) in more developed countries. Likewise, there has been successful implementation of MAS in India (Gupta et al. 2011). Here, we describe progress in a MAS program aimed at improving both GPC and grain weight in cultivar (cv.) HUW234.

Materials and methods

Plant materials

The recurrent parent was cv. HUW234 (HUW12/Sparrow//HUW12), and the donor of both *Gpc-B1* and the TGW QTL was line Glu133 (DBW16/GluPro//2*DBW16). The introgression scheme is illustrated in Fig. 1. The hybrid status of two F₁ plants from the cross HUW234 × Glu133 (designated HUW234-09) was confirmed by genotyping with the *Xucw108* marker linked to *Gpc-B1* (Uauy et al. 2006). These plants were backcrossed to HUW234. Foreground selection on the BC₁F₁ plants was applied for *Gpc-B1* using the *Xucw108* marker (Uauy et al. 2006) and background selection in conjunction with phenotypic selection for HUW234 morphology was applied using 86 genome-wide (four to six per chromosome) polymorphic microsatellite markers, selected following an initial screen of the parents using 592 microsatellite primer pairs. The chromosomal locations and primer sequences are given in <http://wheat.pw.usda.gov/GG2/> (Roder et al. 1998; Somers et al. 2004). The procedure was repeated on the BC₂F₁ materials. BC₂F₂ homozygotes at *Gpc-B1* were selected by progeny tests on 10 plant samples because marker *Xucw108* is dominant. Homozygous *Gpc-B1* BC₂F₃ families were screened with *Xgwm297* to identify those carrying the TGW QTL (Mir et al. 2012). BC₂F₄ populations were generated from selected homozygous plants.

DNA isolation, PCR conditions and electrophoresis

DNA was isolated from one-month-old plants using a CTAB method (Saghai-Marooof et al. 1984). Each PCR in a 20 µL volume was formulated as described by Vishwakarma et al. (2014). *Xucw108* and *Xgwm297* amplicons were resolved in 2.5 % agarose gels. PCR products for background selection were separated in 10 % polyacrylamide gels and visualized by silver staining.

Agronomic performance and assessment of end-use quality

Each BC₂F₄ selection, and both parents, were grown in three-row, 3 × 3 m plots with 22.5 cm between rows




Fig. 1 Crossing and MAS selection scheme to improve GPC and TGW in cv. HUW234. The numbers of plants selected at each generation are indicated in parentheses. RP Recurrent

parent, FG foreground selection, BG background selection, RPG recurrent parent genome

during the 2012–2013 main season at Varanasi. Recommended fertilizer treatments (120 kg N, 60 kg P₂O₅ and 40 kg K₂O per ha) were applied and the trial managed as a commercial crop (Vishwakarma et al. 2014). P and K dressings were given at sowing, and the N application was split between 50 % at sowing, and 25 % each at the time of first (21 days after sowing) and second (45 days after sowing) irrigations. Data from ten plants per family were recorded for plant height (cm), number of tillers per plant, DM (number of days to maturity), spike length (cm), number of spikelets per spike, TGW, and yield per plant (YPP) in BC₂F₁. BC₂F₄ grain yields were obtained from 3 × 3 m row plots. BC₂F₃ grain was dried to a moisture content of 12 %, then used to measure GPC (%) and zinc (ppm) and iron (ppm) contents. GPC was obtained using an Infratec 1241 Grain Analyser (Foss, Hilleroed, Denmark), and Zn

and Fe contents were measured using an X-ray fluorescence device (EDXRF X-Supreme 8000 spectrometer) following the procedure of Paltridge et al. (2012).

Recovery of recurrent parental genotypes

The proportion of the recurrent parent genome (RPG) in each backcross derivative was estimated from:

$$\text{RPG \%} = \frac{2(R) + (H)}{2N} \times 100$$

where R represents the number of homozygous recurrent parent marker alleles, H, the number of heterozygous marker loci, and N, the number of background selection marker loci. The level of similarity between the recurrent parent and the BC₂F₄ derivatives was a measure of the effect of

marker selection on the recurrent parent genome. Graphical genotypes were generated with GGT software (Van Berloo 1999).

For determining significant differences in performance between the introgressed families and the HUW234 recurrent parent, multiple comparisons were made of least square means using a significance level of $\alpha = 0.05$ by a two-sided Dunnett test (Dunnett 1955) in SPSS 16.0 (SPSS Inc. 2007) software.

Results

The *Xucw108* amplicon derived from the cv. Glu133 template was 217 bp, whereas there was no amplification from HUW234. Two F₁ HUW234-09 plants backcrossed to HUW234 produced 76 progeny, of which 38 carried the Glu133 *Xucw108* allele. The use of 86 polymorphic background selection markers, along with phenotypic selection for the recurrent parent phenotype, identified families HUW234-09-69 and -36, for which the RPG recoveries were 81.8 and 79.5 %, respectively (Table S1). These plants were used to produce 82 BC₂F₁ progeny, 36 of which had the *Xucw108* allele. The latter plants were subjected to background genotypic selection (44 microsatellite

loci) and phenotyping (Table S2). The BC₂F₁ selection with the highest RPG (HUW234-09-69-8, 91.9 %) generated 64 BC₂F₂ progeny, of which 46 retained the *Xucw108* allele in the hetero- or homozygous state. Progeny testing identified 15 BC₂F₃ families that were homozygous for the *Xucw108* allele. These 15 BC₂F₃ families were used to generate BC₂F₄ progeny that were evaluated for agronomic performance (Table 1). Six of a further 48 microsatellite loci located near *Gpc-B1* on chromosome 6B were polymorphic between the recipient and donor cultivars (Fig. 2). Based on genotyping with these six markers, nine lines with the shortest introgressed segments were selected for further investigation.


All of the selected BC₂F₄ families had at least 13 % protein, compared to 11.1 % for the recurrent parent. The range in GPC was 13–14.8 % (Table S3). Fe contents of the 15 families ranged from 40.3 to 49.3 ppm, compared to 38 ppm for HUW234; and Zn contents ranged from 41.4 to 55.7 ppm, well above the recurrent parent level of 28 ppm (Table S3). All characteristics of the selected families were similar to HUW234 (Table S4; Fig. 3); DM ranged from 110 to 117 days (mean 112 days), close to that of cv. HUW234; and mean spike length ranged from 8.0 to 11.0 cm, with a mean across families somewhat below

Table 1 Comparison of the parents and 15 HUW234*4 × Glu133 introgression lines for agronomic traits using Dunnett tests

	Parent/line	Trait						
		DM	PH	SPL	SPN	TLN	TGW	GY
	HUW234	112	99	43.6	11	7	37.3	44.7
	Glu 133	120	82.7	49.4	11	11	41.2	46.3
	HUW234-09-21	2.00	−21.9*	1.60*	−1.00	3.00*	4.10*	2.00
	HUW234-09-47	−2.00	−9.9*	0.70	−2.00*	3.00*	2.70*	1.00
	HUW234-09-48	−2.00	−0.80	7.50*	−2.00*	2.00*	4.60*	4.00*
	HUW234-09-49	−1.00	−9.7*	1.70*	0.00	4.00*	4.30*	0.70
	HUW234-09-65	−1.00	−5.8*	5.80*	−1.00	5.00*	5.10*	1.00
	HUW234-09-112	3.0*	−1.60	3.70*	−2.00*	5.00*	1.50*	0.20
	HUW234-09-114	5.0*	−3.6*	4.50*	−2.00*	4.00*	−0.50	0.40
	HUW234-09-116	3.0*	−9.7*	4.00*	0.00	4.00*	4.20*	0.50
	HUW234-09-117	−1.00	−20.1*	5.90*	−1.00	5.00*	4.70*	1.50
	HUW234-09-128	−2.00	−11.4*	4.60*	−3.00*	2.00*	−0.20	0.70
	HUW234-09-213	−1.00	−2.8*	2.10*	−3.00*	1.00	3.90*	0.00
	HUW234-09-214	0.00	−8.2*	5.50*	−3.00*	1.00	4.50*	−1.50
	HUW234-09-239	2.00	−6.5*	2.30*	−2.00*	2.00*	1.10	0.10
	HUW234-09-255	−2.00	−1.70	−0.10	−1.00	4.00*	4.30*	−0.80
	HUW234-09-256	0.00	−7.3*	8.50*	−2.00*	4.00*	4.50*	0.50
	SE±	0.821	0.810	0.480	0.400	0.394	0.410	0.990

Positive and negative values are relative to HUW234
DM Days to maturity, *PH* plant height, *SPN* spikelet number, *SPL* spikelet length, *TLN* number of tillers, *TGW* 1000-grain weight, *GY* grain yield qt/ht, *SE* standard error
 * Significantly different at $P = 0.05$ level

Fig. 2 Graphical representations of the selected BC₂F₄ lines with respect to the *Gpc-B1* region on chromosome 6B. Colored segments represent segments from the donor, RP and heterozygotes. (Color figure online)



that of the recurrent parent (9.3 vs 11.0 cm); maximum effective tiller number was superior to HUW234 and TGW ranged between 38.4 and 42.4 g, 2.9 and 13.7 % higher than HUW234. Thirteen selected lines out-yielded HUW234; the best by up to 8.9 %, while two were lower yielding.

The performances of BC₂F₄ families were evaluated by Dunnett test. This test treats one family as a control, and compares all others to it. Dunnett tests were significant ($P < 0.05$) for 72 % of 150 comparisons between introgressed families and recurrent parent HUW234 (Table 2). We found significantly

higher GPC, Fe and Zn contents in almost all BC₂F₄ families compared to HUW234. Dunnett tests were significant ($P < 0.05$) for days to maturity, plant height, spikelet number, spikelet length, number of tillers, TGW and GY for 3, 12, 13, 9, 13, 12 and 1 BC₂F₄ families, respectively.

Discussion

The addition of genotype to conventional phenotypic selection succeeded in recovering over 90 % of the

Fig. 3 BC₂F₄ families selected for the presence of *Gpc-B1* and TGW compared to the recurrent parent



Table 2 Comparison of the parents and 15 HUW234*4 × Glu133 introgression lines for GPC and Fe and Zn contents using Dunnett tests

Parents/lines	Traits		
	GPC (%)	Fe (ppm)	Zn (ppm)
HUW234	11.15	38	28
Glu 133	14	45.2	49
HUW234-09-21	2.55*	4.00*	14.10*
HUW234-09-47	2.75*	7.20*	23.60*
HUW234-09-48	3.65*	8.00*	20.30*
HUW234-09-49	2.65*	8.30*	18.00*
HUW234-09-65	2.25*	5.60*	16.40*
HUW234-09-112	1.85*	2.50*	13.40*
HUW234-09-114	2.55*	9.30*	21.90*
HUW234-09-116	1.85*	3.30*	19.00*
HUW234-09-117	2.65*	3.20*	20.20*
HUW234-09-128	2.05*	2.30*	27.70*
HUW234-09-213	2.35*	11.30*	21.80*
HUW234-09-214	1.95*	10.40*	20.70*
HUW234-09-239	2.75*	7.40*	17.80*
HUW234-09-255	2.05*	3.30*	15.90*
HUW234-09-256	2.25*	7.40*	21.80*
SE±	0.50	0.23	0.41

GPC (%) Grain protein content, Fe iron content, Zn zinc content, SE standard error

* Significantly different at P = 0.05

genome of the recipient parent after two backcrosses (the average recovery using only foreground selection is 87.5 %) and a significant improvement to the GPC was attributed to the introduction of *Gpc-B1*. In addition to significantly increased GPC, both Fe and Zn contents of the grain were higher than for HUW234, and progress was also made in improved TGW and some other agronomic traits. Grain Fe and Zn contents were shown to be positively correlated with GPC (Distelfeld et al. 2006; Uauy et al. 2006), and ascribed to a pleiotropic effect of *Gpc-B1* rather than linkage. The significant progress in TGW appeared to be novel as the usual negative correlation between yield and GPC was broken without affecting days to maturity. The beneficial effects on TGW and other traits could be due to pleiotropic effects of *Gpc-B1* as experienced in previous studies (Kumar et al. 2011; Tyagi et al. 2014; Vishwakarma et al. 2014). Pyramiding of eight QTLs including those of GPC and TGW using MAS was demonstrated by Tyagi et al. (2014) in the then most popular Indian cultivar PBW343. Likewise Kumar et al. (2011) introgressed *Gpc-B1* into ten elite Indian varieties, and Vishwakarma et al. (2014) successfully introgressed *Gpc-B1* into HUW468 by backcrossing. In these studies only HUW468 was a NEPZ cultivar; the others were from the north western plains region.

MAS has made a significant impact on crop improvement programs (Gao et al. 2005; Barloy et al. 2007; Gupta et al. 2010; Kumar et al. 2011; Tyagi et al. 2014). Backcrossing strategies are particularly suited to MAS approaches, with markers providing opportunities for both foreground and background selection. MAS allows an accelerated recovery of the recipient parent genotype, the actual purpose of backcrossing. Foreground selection requires a tightly linked single marker (such as *Xucw108*), or less closely linked flanking markers (such as *Xgwm508* and *Xgwm193* for *Gpc-B1* (see Davies et al. 2006) in order to ensure selection of the target gene and its associated trait. To select superior recombinant lines with minimal donor parent contributions we used six flanking markers (*Xgwm132*, *Xcfd190*, *Xgwm508*, *Xgwm193*, *Xgwm361* and *Xgwm219*) covering the gene *Gpc-B1* (Olmos et al. 2003; Vishwakarma et al. 2014).

The goal of the present work was to obtain an improved version of a well-established cultivar. The *Xucw108* marker used to track the presence of *Gpc-B1* is dominant and could not discriminate between the homozygous and heterozygous conditions. While this is not a problem during backcrossing, it necessitates progeny testing of the first selfing generation (BC₂F₂). Phenotypic selection for agronomic features and genotypic selection based on microsatellites to restore the HUW234 genetic background were also undertaken. To select both GPC and TGW in the same lines, we followed a stepwise screening method, by which we screened for *Gpc-B1* with linked marker *Xucw108* up to BC₂F₃, along with phenotypic selection for agronomic traits with special emphasis on TGW since the donor Glu 133 had 15 % higher TGW than the recurrent parent. Following phenotypic selection, screening for TGW was conducted using the linked marker *Xgwm297* in BC₂F₃ (Mir et al. 2012). A benefit of this strategy was to reduce the number of progenies requiring screening during the selfing generations (Kumar et al. 2011).

Although HUW234 is still grown over a large area in the NEP zone it is disadvantaged by modest GPC and low TGW, traits that were targeted in this work. Background selection was considered to be desirable in order to reconstitute the overall genetic background HUW234 as rapidly and thoroughly as possible. Phenotypic selection as recommended by previous workers (Kumar et al. 2011; Tyagi et al. 2014;

Vishwakarma et al. 2014) was also used. We believe this approach provides a model for breeding for wheat quality in areas such as the NEP zone of India and neighboring countries with limited investments in wheat improvement programs.

Acknowledgments We gratefully acknowledge partial funding from a DBT, New Delhi Network Project and a CSISA - Wheat Breeding project funded by the Bill and Melinda Gates Foundation and USAID for field based support. We sincerely thank Prof. Pushpendra K. Gupta, Ch. Charan Singh University, Meerut for sustained intellectual support. Use of the Indian Agriculture Research Institute off-season facility at Regional Station, Wellington, Tamil Nadu is gratefully acknowledged. We thank Dr. Parveen Chuneja (Punjab Agriculture University, Ludhiana, India) for supplying grain of line Glu133 and Dr. R. R. Mir (S. K. University of Agricultural Sciences and Technology, Jammu, India) for providing TGW marker information.

Compliance with ethical standards

Conflicts of Interest The authors declare that they have no conflicts of interest.

References

- Avivi L (1978) High grain protein content in wild tetraploid wheat Korn. In: Ramanujam S (ed) Proceedings of the 5th wheat genet symposium. Indian Soc Genet & Plant Breed, New Delhi, pp 372–380
- Barloy D, Lemoine J, Abelard P, Tanguy AM, Rivoal R, Jahier J (2007) Marker assisted pyramiding of two cereal cyst nematode resistance genes from *Aegilops variabilis* in wheat. *Mol Breed* 20:31–40
- Brevis JC, Dubcovsky J (2010) Effect of chromosome region including the *Gpc-B1* locus on wheat protein and protein yield. *Crop Sci* 50:93–104
- Brevis JC, Morris CF, Manthey F, Dubcovsky J (2010) Effect of the grain protein content locus *Gpc-B1* on bread and pasta quality. *J Cereal Sci* 51:357–365
- Campbell KG, Bergman CJ, Gualberto DG, Anderson JA, Giroux MJ, Hareland G, Fulcher RG, Sorrells ME, Finney PL (1999) Quantitative trait loci associated with kernel traits in soft × hard wheat cross. *Crop Sci* 39:1184–1195
- Cox MC, Qualset CO, Rains DW (1985) Genetic variation for nitrogen assimilation and translocation in wheat. III. Nitrogen translocation in relation to grain yield and protein. *Crop Sci* 26:737–740
- Davies J, Berzonsky WA, Leach GD (2006) A comparison of marker-assisted and phenotypic selection for high grain protein content in spring wheat. *Euphytica* 152:117–134
- Distelfeld A, Uauy C, Olmos S, Schlatter AR, Dubcovsky J, Fahima T (2004) Microcolinearity between a 2-cM region encompassing the grain protein content locus *GPC-6B1* on wheat chromosome 6B and a 350-kb region on rice chromosome 2. *Funct Integr Genomics* 4:59–66

- Distelfeld A, Uauy C, Fahima T, Dubcovsky J (2006) Physical map of the wheat high-grain protein content gene *Gpc-B1* and development of a high-throughput molecular marker. *New Phytol* 169:753–763
- Dunnett C (1955) A multiple comparison procedure for comparing several treatments with a control. *J Am Stat Assoc* 50:1096–1121
- FAOSTAT (2012) Faostat.fao.org/site/339/default.aspx and www.fao.org/worldfoodsituation/wfs-home/csdb/en
- Gao AL, He HG, Chen QZ (2005) Pyramiding wheat powdery mildew resistance genes Pm2, Pm4a and Pm21 by molecular marker-assisted selection. *Acta Agron Sin* 31:1400–1405
- Gegas VC, Nazari A, Griffiths S, Simmonds J, Fish L, Orford S, Sayers L, Doonan JH, Snape JW (2010) A genetic framework for grain size and shape variation in wheat. *Plant Cell* 22:1046–1056
- Gupta RK (2004) Quality of Indian wheat and infrastructure for analysis. In: Joshi AK, Chand R, Arun B, Singh G (eds) A compendium of the training program (26–30 December, 2003) on wheat improvement in eastern and warmer regions of India: conventional and non-conventional approaches. NATP project, (ICAR), BHU, Varanasi, India
- Gupta PK, Langridge P, Mir RR (2010) Marker-assisted wheat breeding: present status and future possibilities. *Mol Breed* 26:145–161
- Gupta PK, Balyan HS, Mir RR, Kumar J, Kumar A, Kumar S, Jaiswal V, Tyagi S, Kumari S (2011) QTL analysis, association mapping and marker-assisted selection for some quality traits in bread wheat—an overview of the work done at CCS University, Meerut. *J Wheat Res* 3(2):1–11
- Huang XQ, Coster H, Ganal MW, Roder MS (2003) Advanced backcross QTL analysis for the identification of quantitative trait loci alleles from wild relatives of wheat (*Triticum aestivum* L.). *Theor Appl Genet* 106:1379–1389
- Huang XQ, Kempf H, Ganal MW, Roder MS (2004) Advanced backcross QTL analysis in progenies derived from a cross between a German elite winter wheat variety and a synthetic wheat (*Triticum aestivum* L.). *Theor Appl Genet* 109:933–9943
- Joppa LR, Du C, Hart GE, Hareland GA (1997) Mapping gene(s) for grain protein in tetraploid wheat (*Triticum turgidum* L.) using a population of recombinant inbred chromosome lines. *Crop Sci* 37:1586–1589
- Kade M, Barneix AJ, Olmos S, Dubcovsky J (2005) Nitrogen uptake and remobilization in tetraploid ‘Langdon’ durum wheat and a recombinant substitution line with the high grain protein gene *Gpc-B1*. *Plant Breed* 124:343–349
- Ketata H, Edwards LH, Smith EL (1976) Inheritance of eight agronomic characters in a winter wheat cross. *Crop Sci* 16:19–22
- Khan IA, Procnunier JD, Humphreys DG, Tranquilli G, Schlatter AR, Marcucci-Poltri S, Froberg R, Dubcovsky J (2000) Development of PCR-based markers for a high grain protein content gene from *Triticum turgidum* ssp. *dicoccoides* transferred to bread wheat. *Crop Sci* 40:518–524
- Kulwal PL, Kumar N, Kumar A, Gupta RK, Balyan HS, Gupta PK (2005) Gene networks in hexaploid wheat: interacting quantitative trait loci for grain protein content. *Funct Integr Genomics* 5:254–259
- Kumar J, Jaiswal V, Kumar A, Kumar N, Mir RR, Kumar S, Dhariwal R, Tyagi S, Khandelwal M, Prabhu KV, Prasad R, Balyan HS, Gupta PK (2011) Introgression of a major gene for high grain protein content in some Indian bread wheat cultivars. *Field Crop Res* 123:226–233
- Lobell DB, Schlenker W, Costa-Roberts J (2011) Climate trends and global crop production since 1980. *Science* 333:616–620
- McCartney CA, Somers DJ, Humphreys DG, Lukow O, Ames N, Noll J, Cloutier S, McCallum BD (2005) Mapping quantitative trait loci controlling agronomic traits in the spring wheat cross RL4452 × ‘AC Domain’. *Genome* 48:870–883
- Mir RR, Kumar N, Jaiswal V, Girdharwal N, Prasad M, Balyan HS, Gupta PK (2012) Genetic dissection of grain weight in bread wheat through quantitative trait locus interval and association mapping. *Mol Breed* 29:963–972
- Olmos S, Distelfeld A, Chicaiza O, Schlatter AR, Fahima T, Echenique V, Dubcovsky J (2003) Precise mapping of a locus affecting grain protein content in durum wheat. *Theor Appl Genet* 107:1243–1251
- Paltridge NG, Palmer LJ, Milham PJ, Guild GE, Stangoulis JCR (2012) Energy-dispersive X-ray fluorescence analysis of zinc and iron concentration in rice and pearl millet grains. *Plant Soil* 361:251–260
- Prasad M, Kumar N, Kulwal PL, Roder M, Balyan HS, Dhaliwal HS, Gupta PK (2003) QTL analysis for grain protein content using SSR markers and validation studies using NILs in bread wheat. *Theor Appl Genet* 106:659–667
- Roder MS, Korzun V, Wendehake K, Plaschke J, Tixier MH, Leroy P, Ganal MW (1998) A microsatellite map of wheat. *Genetics* 149:2007–2023
- Roder MS, Huang XQ, Börner A (2008) Fine mapping of the region on wheat chromosome 7D controlling grain weight. *Funct Integr Genomics* 8:79–86
- Saghai-Marouf MA, Soliman KM, Jorgensen RA, Allard RW (1984) Ribosomal DNA spacer length polymorphisms in barley: mendelian inheritance, chromosomal location, and population dynamics. *Proc Natl Acad Sci U.S.A.* 81:8014–8018
- Simmonds NW (1995) The relation between yield and protein in cereal grain. *J Sci Food Agric* 67:309–315
- Singh H, Prasad M, Varshney RK, Roy JK, Balyan HS, Dhaliwal HS, Gupta PK (2001) STMS markers for grain protein content and their validation using near-isogenic lines in bread wheat. *Plant Breed* 120:273–278
- SPSS Inc. (2007). SPSS for windows, Version 16.0. Chicago, SPSS Inc.
- Somers DJ, Isaac P, Edwards K (2004) A high-density microsatellite consensus map for bread wheat (*Triticum aestivum* L.). *Theor Appl Genet* 109:1105–1114
- Tyagi S, Mir RR, Chhuneja P, Ramesh B, Balyan HS, Gupta PK (2014) Marker-assisted pyramiding of eight QTLs/genes for seven different traits in common wheat (*Triticum aestivum* L.). *Mol Breed* 34:167–175
- Uauy C, Distelfeld A, Fahima T, Blechl A, Dubcovsky J (2006) A NAC gene regulating senescence improves grain protein, zinc, and iron content in wheat. *Science* 314:1298–1301
- Van Berloo R (1999) GGT: software for display of graphical genotypes. *J Hered* 90:328–329

- Vishwakarma MK, Mishra VK, Gupta PK, Yadav PS, Kumar H, Joshi AK (2014) Introgression of the high grain protein gene Gpc-B1 in an elite wheat variety of Indo-Gangetic Plains through marker assisted backcross breeding. *Curr Plant Biol* 1:60–67
- Wang RX, Hai L, Zhang XY, You GX, Yan CS, Xiao SH (2009) QTL mapping for grain filling rate and yield-related traits in RILs of the Chinese winter wheat population Heshangmai × Yu8679. *Theor Appl Genet* 118:313–325
- Wang L, Cui F, Wang J, Jun L, Ding A, Zhao C, Li X, Feng D, Gao J, Wang H (2012) Conditional QTL mapping of protein content in wheat with respect to grain yield and its components. *J Genet* 91:303–312