

1056

C. S. Busso · C. J. Liu · C. T. Hash · J. R. Witcombe  
K. M. Devos · J. M. J. de Wet · M. D. Gale

## Analysis of recombination rate in female and male gametogenesis in pearl millet (*Pennisetum glaucum*) using RFLP markers

Received: 9 May 1994 / Accepted: 8 September 1994

**Abstract** Sex as a factor affecting recovered recombination in plant gametes was investigated in pearl millet, *Pennisetum glaucum*, by using reciprocal three-way crosses [(A × B) × C] × C × (A × B)]. The two populations were mapped at 42 loci pre-selected to cover the majority of the genome. No differences in recombination distances were observed at the whole-genome level and only a few individual linkage intervals were found to differ, all in favour of increased recombination through the male. Distorted segregations found in the three-way crosses provide evidence of post-gametic selection for particular gene(s) or chromosome regions. The significance of these results for the design of pearl millet breeding programmes and inheritance experiments, as well as for other experimental strategies, is discussed.

**Key words** *Pennisetum glaucum* · RFLP analysis  
Recombination rate · Segregation distortion  
Genetic mechanisms

### Introduction

Recombination at meiosis and the factors that influence recombination rates have been studied extensively. One such factor is the relative recombination rates recovered in male and female gametes. In animals such as *Drosophila* (Baker et al. 1976) and humans (Donis-Keller et al. 1987) the evidence shows that recombination is much higher in the female. However, analysis of recombination frequencies in mouse (Reeves 1990) and silk worm (Maeda 1939)

demonstrated higher values in male gametes. In plants, where a clear effect of sex on recombination could be exploited in breeding programmes, the evidence is also equivocal. A number of studies, e.g., in *Zea mays* (Robertson 1984) and *Arabidopsis thaliana* (Zhuchenko et al. 1989), have shown a tendency towards increased recombination in the male. However, these results do not exclude the possibility that the differences are restricted to particular chromosome regions since all the studies involved a limited number of recombination intervals using available morphological or protein marker loci.

With the development of DNA markers and the construction of extensive genetic maps, analyses of recombination frequencies can be extended to an entire genome. In a comparison of male versus female recombination in two backcross populations of tomato using RFLP markers, De Vicente and Tanksley (1991) reported a significantly higher recombination rate in female meiosis. This result was associated with a skewed allelic segregation, as has been observed in many of the molecular mapping projects in plant crop species, e.g., in barley (Heun et al. 1991) and in pearl millet (Liu et al. 1994). These reports indicated that the skewed segregation could not explain the differences in recombination rates. It is clearly important to establish whether the direction and extent of the sex differences in recombination rates in tomato reflect a general situation in plants. From a practical standpoint both differential recombination in male and female-meiosis and distorted segregation, if predictable, can be exploited to the plant breeders' advantage.

In this paper we report the results of experiments in pearl millet, *Pennisetum glaucum*, designed to examine the effects of sex on recombination and segregation distortion with a view to possible exploitation in breeding strategies.

Communicated by P. L. Pfahler

C. S. Busso · C. J. Liu · K. M. Devos · M. D. Gale (✉)  
Cambridge Laboratory, Colney, Norwich, NR4 7UJ, UK

C. T. Hash · J. M. J. de Wet  
International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru PO 502 324, Andhra Pradesh, India

J. R. Witcombe  
Centre for Arid Zone Studies, University of Wales, Bangor  
Gwynedd LL57 2UW, UK

### Materials and methods

#### Genetic material

Three genotypes, each of which had been inbred for more than nine generations, were used in three-way crosses: 81B (a downy mildew-

resistant selection from gamma-irradiated Tift 23DB). ICMP 451 (derived from LCSN 72-1-2-1-1, a selection made in Upper Volta from the ICRISAT Center Late Composite) and BKM 1163 (an inbred derived from crosses between Tift 23DB and a Ghanaian landrace).

$F_1$  plants of 81B $\times$ ICMP 451 were obtained by protogyny-facilitated hand pollination without employing emasculation. A single typical  $F_1$  plant was then crossed as both male and female with the third parent, BKM 1163. All pollinations were carried out at 35 °C in a glasshouse with natural daylength at the ICRISAT Center, and were repeated on 3 consecutive days to reduce the probability of selfing.

The parents were morphologically distinct: 81B is dwarf and has hairy leaf blades; ICMP 451 is tall, with bristled panicles and smooth leaf blades; and BKM 1163 is dwarf, with smooth leaf blades and amber seed colour.  $F_1$  plants of 81B  $\times$  ICMP 451 were distinguished from selfs by the dominant characteristics of the male parent (smooth leaf blades, bristled panicles, and tall plants).  $F_1$  seed of the three-way cross having BKM 1163 as the male parent, TWC $\delta$ , [(81B  $\times$  ICMP 451)  $\times$  BKM 1163] were distinguished from selfed seed by xenia for amber coloured endosperm colour, whereas selfed and  $F_1$  seed in TWC $\varphi$  [BKM 1163  $\times$  (81B  $\times$  ICMP 451)] were indistinguishable.

From each three-way cross, TWC $\varphi$  and TWC $\delta$ , 112 plants were grown to the five-leaf stage and then harvested directly into liquid nitrogen before freeze drying. This material, together with leaf samples from the parental and  $F_1$  genotypes, was further characterized at the Cambridge Laboratory.

#### RFLP analysis

Sixty-eight low-copy PSM DNA clones were chosen to provide coverage of the current pearl millet map (Liu et al. 1994). Of these 41 were found to be informative and were used for mapping the three-way crosses. Analyses with the restriction enzymes *EcoRI*, *EcoRV*, *DraI* and *HindIII* were carried out as described by Liu et al. (1994). The segregation data were analyzed using Mapmaker v1.9, Whitehead Institute, Mass., USA (Lander et al. 1987). The maps are presented in cM, using the Haldane mapping function.

## Results

The use of three-way crosses allowed off-type progenies, in particular those produced by selfing, to be identified unequivocally (Fig. 1). After elimination of such progenies,

the segregation data for the 112 TWC $\varphi$  plants and the remaining 107 TWC $\delta$  plants were analyzed to produce two genetic maps (Fig. 2). In all cases the marker order and linkage group assignment, of the 42 loci that could be mapped using the 41 probes, was consistent between the reciprocal crosses and the map constructed in an independent  $F_2$  (LGD-1-B-10  $\times$  ICMP 85410) population by Liu et al. (1994).

#### Recombination in male and female gametes


The total map distances estimated in the two populations did not differ significantly, being 234 cM in TWC $\varphi$  and 267 cM in TWC $\delta$ . These values were also similar to the 255.3 cM map distance spanned by the same markers in the (LGD-1-B-10  $\times$  ICMP 85410)  $F_2$  map.

Analysis of the seven linkage groups showed that in only one, the short linkage group 3, was there a difference ( $P < 0.05$ ) between the length of the TWC $\delta$  map (22 cM) and the length of the TWC $\varphi$  map (9 cM). However, observation of the remaining six groups shows clearly that the sex difference does not reflect a general trend. The analysis of the 35 linkage intervals for which reciprocal data were available showed that only three intervals were significantly different at  $P < 0.05$ , and of these only two were significantly different at  $P < 0.01$ . These were *Xpsm223-Xpsm607* in linkage group 1,  $\chi^2_{(1)} = 7.4$ , and *Xpsm451B-Xpsm248* in linkage group 3,  $\chi^2_{(1)} = 7.1$ , with the TWC $\delta$  map having the larger value in each case.

#### Segregation distortion

The three-way cross design allowed transmission of the alternative 81B and ICMP 451 alleles through pollen and egg cells to be scored with precision. Severe distortion from the expected 1:1 ratio was observed, only in TWC $\delta$ , for alleles at loci in linkage groups 1, 4 and 7. The 12 loci sig-

**Fig. 1** Segregation of *Xpsm466* in parents (A, B), the  $F_1$  plant and a sample of the TWC $\delta$  [C  $\times$  (A  $\times$  B)] population digested with *DraI*. Note, the individual progenies are classified by parental genotype (A or B), always recovered with the female C allele, and off-type (homozygous C) in the case of two selfed progeny






Fig. 2 Male and female genetic maps. Note, linkage groups or intervals showing differences between the TWC♂ and TWC♀ map are shown by \* $P < 0.5$  and \*\* $P < 0.01$  in the direction of greater recombination. Loci showing distorted segregation, found only in the TWC♂ population, are shown by # $P < 0.05$  and ## $P < 0.01$

nificantly affected in TWC♂ all showed segregation in favour of the ICMP 451 alleles (Fig. 3) and smaller non-significant biases occurred in TWC♀ along part of linkage group 1 and most of linkage group 4 (Fig. 3).

## Discussion

Analysis of the 81B × ICMP 451 three-way crosses confirms the findings of Liu et al (1994) in that the markers mapped in the expected linear arrangement on the same chromosomes, separated by linkage distances similar to those previously observed. Thus the analyses also confirmed the extremely short genetic map described by Liu et al. using 181 random markers. Although the map has not been capped by mapped telomeres it is unreasonable to expect that, with an average of 26 markers per chromosome, a large portion of the genome lies outside the distal markers. Genetic lengths of between 100 and 200 cM per chromosome appear to be the norm in other cereals, e.g., bar-

ley (Kleinhofs et al. 1993) and wheat (Devos et al. 1993). It is probable therefore that the low level of recombination per chromosome will limit the rate of progress achieved by pearl millet breeders. Thus any means of maximising recombination will be of value. For example, more generations of random mating can be made, before selection commences, in the creation of composite populations. Also, in population improvement by recurrent selection, mass selection and full-sib progeny selection, which allow frequent recombination generations, can be used rather than methods which predominantly use selfing generations. However, an advantage of the genetically short chromosomes is that they provide an excellent test bed for experiments, such as the one reported here, to measure the effects of factors that influence recombination. Moreover, marker-assisted recovery of non-target recurrent parent chromosomes in backcross breeding programmes will be easy, and relatively inexpensive, in this species.

## Differences in male and female recombination

The experiments described here were designed to eliminate other factors that could affect recombination rates in plants. Temperature and photoperiod at meiosis were controlled. Individual genotype effects were eliminated by the use of a single F<sub>1</sub> plant as both male and female in the backcrosses. The effects of non-collinearity, as might be found



found possible in an independent  $F_2$  population, there is no reason to believe that they are in any way exceptional. These equivalent recombination rates suggest that backcrossing programmes can be carried out in the most convenient manner, rather than using the  $F_1$  as female as would be suggested from the tomato results (De Vicente and Tanksley 1991). However, breeders wishing to avoid surprises due to segregation distortion should use the segregating genome as the female in crosses. Other applications, such as the construction of inbred substitution lines by marker-assisted selection for entire linkage groups of a donor inbred, will not be able to take advantage of reduced recombination in one gamete or the other to hold linkage groups together. Fortunately, given the low rates of recombination observed in *P. glaucum*, such aides are probably not necessary.

Our results indicate that segregation will be more predictable among female gametes than among male gametes in pearl millet. If breeders are seeking Mendelian segregation ratios in this species, they should examine test-cross progenies where the  $F_1$  has been used as the female parent (contrary to standard practice where it is more often used as the male parent), rather than  $F_2$  populations, or test-crosses where the  $F_1$  is used as the pollen parent. This has recently been demonstrated to hold true where segregation distortion due to gametophytic selection was found in progenies that are predicted to segregate 3:1 or 1:1 for the yellow seedling character of ICMB 88004 (K. N. Rai, personal communication). Further, in a backcrossing program to introduce one or more genes of interest into an elite background, pearl millet breeders should use the recurrent parent as the male, which can conveniently be done by using stored pollen (Hanna 1990) if this facility is available. The male recurrent parent will permit the use of standard formulae (e.g., Sedcole 1977) to calculate the number of plants required to ensure transmission of the donor gene(s) of interest each generation, facilitating their timely and efficient backcross transfer.

**Acknowledgements** This work was supported by the Overseas Development Administration Plant Sciences Research Programme. We are also grateful to Dr. J. K. M. Brown at the John Innes Centre for the assistance in data analysis and Miss T. Pittaway for her technical assistance. Finally, a special thanks to collaborators in ICRISAT: Mr. M. N. V. Ratnaji Rao, Mr. Mohd. Yousuf and Mr. A. C. Reddy.

## References

- Baker BS, Carpenter ATC, Esposito MS, Esposito RE, Sandler L (1976) The genetic control of meiosis. *Annu Rev Genet* 10:53-134
- De Vicente MC, Tanksley SD (1991) Genome-wide reduction in recombination of backcross progeny derived from male versus female gametes in an interspecific cross of tomato. *Theor Appl Genet* 83:173-178
- Devos KM, Millan T, Gale MD (1993) Comparative RFLP maps of the homoeologous group-2 chromosomes of wheat, rye and barley. *Theor Appl Genet* 85:784-792
- Donis-Keller H, Green P, Helms C, Cartinhour S, Weiffenbach B, Stephens K, Keith TP, Bowden DW, Smith DR, Lander ES, Botstein D, Akots G, Rediker KS, Gravioux T, Brown V, Rising MB, Parker C, Powers JA, Watt DE, Kauffman ER, Bricker A, Phipps P, Muller-Ahle H, Fulton TM, Ng S, Schumm JW, Braman JC, Knowlton RG, Barker DF, Crooks SM, Lincoln SE, Daly MJ, Abrahamson J (1987) A genetic linkage group map of human genome. *Cell* 51:319-337
- Hanna WW (1990) Long-term storage of *Pennisetum glaucum* (L.) R. Br. pollen. *Theor Appl Genet* 79:605-608
- Heun M, Kennedy AE, Anderson JA, Lapitan NLV, Sorrells ME, Tanksley SD (1991) Construction of a restriction fragment length polymorphism map for barley (*Hordeum vulgare*). *Genome* 34:437-447
- Kleinohfs A, Kilian A, Saghai Maroof MA, Biyashev RM, Hayes P, Chen FQ, Lapitan N, Fenwick A, Blake TK, Kanazin V, Ananiev E, Dahleen L, Kudrna D, Bollinger J, Knapp SJ, Liu B, Sorrells M, Heun M, Framckowiak JD, Hoffman D, Skadsen R, Steffenson BJ (1993) A molecular, isozyme and morphological map of the barley (*Hordeum vulgare*) genome. *Theor Appl Genet* 86:705-712
- Lander E, Green P, Abrahamson J, Barlow A, Daley M, Lincoln S, Newburg L (1987) MAPMAKER: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* 1:174-181
- Liu CJ, Witcombe JR, Pittaway TS, Nash M, Busso CS, Gale MD (1994) An RFLP-based genetic map of pearl millet (*Pennisetum glaucum*). *Theor Appl Genet* 89:481-487
- Maeda T (1939) Chiasma studies in the silk worm, *Bombyx mori* L. *Jpn J Genet* 15:118-127
- Reeves RH (1990) Sex, strain, and species differences affect recombination across an evolutionarily conserved segment of mouse chromosome 16. *Genomics* 8:141-148
- Robert T, Sarr A, Pernes J (1989) Sélections sur la phase haploïde chez le Mil [*Pennisetum typhoides* (Burm.) Stapf et Hubb.]: effet de la température. *Genome* 32:946-952
- Robert T, Lespinasse R, Pernes J, Sarr A (1991) Gametophytic competition as influencing gene flow between wild and cultivated forms of pearl millet (*Pennisetum typhoides*). *Genome* 34:195-200
- Robertson DS (1984) Different frequency in the recovery of crossover products from male and female gametes of plants hypoploid for B-A translocations in maize. *Genetics* 107:117-130
- Sarr A, Pernes J (1988) Analyses multivariées de descendance de rétrocroisements et mise en évidence de distorsions de ségrégation de caractères quantitatifs chez le mil [*Pennisetum typhoides* (Burm.) Stapf et Hubb.]. *Genome* 30:411-422
- Sarr A, Sandmeier M, Pernes J (1988) Gametophytic competition in pearl millet, *Pennisetum typhoides* (Stapf et Hubb.). *Genome* 30:924-929
- Sedcole TR (1977) Number of plants necessary to recover a trait. *Crop Sci* 17:667-668
- Zhuchenko AA, Korol AB, Vizir IY, Bocharnikova NI, Zamorzaeva NI (1989) Sex differences in crossover frequency for tomato and thale cress (*Arabidopsis thaliana*). *Sov Genet* 24:1104-1110