

Improving Phosphorus Use through Co-inoculation of Vesicular Arbuscular Mycorrhizal Fungi and Phosphate-Solubilizing Bacteria in Maize in an Acidic Alfisol

V. K. SURI,¹ ANIL K. CHOUDHARY,² GIRISH CHANDER,¹
T. S. VERMA,¹ M. K. GUPTA,³ AND N. DUTT¹

¹Department of Soil Science, Himachal Pradesh Agricultural University, Palampur, Himachal Pradesh, India

²Department of Agronomy, Himachal Pradesh Agricultural University, Palampur, Himachal Pradesh, India

³Department of Microbiology, CSK Himachal Pradesh Agricultural University, Palampur, Himachal Pradesh, India

*Performance of three vesicular arbuscular mycorrhizal (VAM) fungi cultures and a phosphate-solubilizing bacteria (PSB) culture alone or in combination with or without 75% of the recommended P₂O₅ dose based on soil-test crop response model was examined in maize in a phosphorus (P)-deficient acidic Alfisol in a glasshouse pot experiment. Sole application of VAM besides co-inoculation with PSB (*Pseudomonas striata*) and inorganic P stimulated mycorrhizal root colonization. Sole application of PSB, VAM_T (*Glomus intraradices*), and VAM_I (*Glomus mosseae*) as well as co-inoculation of VAM with PSB significantly improved crop productivity besides grain protein content, thus indicating a synergistic interaction between VAM and PSB. Application of VAM_T or VAM_I + PSB + 75% P₂O₅ remained at par with sole application of 100% P₂O₅ dose with regard to productivity, nutrient uptake, and soil fertility status (particularly P), thus indicating economization of fertilizer P to the tune of about 25% without compromising crop productivity and soil fertility in an acidic Alfisol.*

Keywords *Glomus mosseae*, maize, phosphate-solubilizing bacteria, productivity, soil fertility, vesicular arbuscular mycorrhizae

Introduction

Most mountain soils of the northwestern Himalayas are deficient in available phosphorus (P) because of their acidic nature (Anonymous 1997) and have high P-fixing power because of excess iron (Fe) and aluminium (Al) ions (Narsian and Patel 2009), resulting into low P availability for crop production (Sharma, Verma, and Bhumbla 1980). Again, phosphatic fertilizers constitute one of the most expensive inputs in agriculture, which warrants their efficient use. In view of environmental concerns and current developments in sustainability, research efforts are concentrated on development of agrotechniques that use less expensive

Received 11 June 2010; accepted 15 March 2011.

Address correspondence to V. K. Suri, CSK Himachal Pradesh Agricultural University, Palampur, Himachal Pradesh, India. E-mail: surivk13jan@gmail.com

sources of plant nutrients such as biofertilizers or inorganic P coupled with biofertilizers (Narsian and Patel 2009). In this context, use of vesicular arbuscular mycorrhizae (VAM) and phosphate-solubilizing bacteria (PSB) might be useful in reversing the process of P fixation in an acidic Alfisol of the northwestern Himalayas, thereby enhancing P supply to maize. The VAM will largely mobilize available P by increasing the surface area of roots and greater exploration of soil volume by its hyphal network, leading to higher overall nutrient use efficiency (George et al. 1992). The PSB will render insoluble phosphates soluble to some extent. The present study was undertaken with the objective of finding the extent to which the combined application of VAM and PSB might reduce P fertilizer dose in maize.

Materials and Methods

Experimental Details

Glasshouse studies were conducted in maize (Kharif 2003) as a test crop in pots filled with 10 kg of 2-mm sieved air-dried soil. Silty clay loam soil (Typic Hapludalf) used in the study had an initial organic carbon (C) status of 3.5 g kg⁻¹ soil. The available nitrogen (N), P, and potassium (K) contents of experimental soil were 49.1, 2.2, and 124.1 mg kg⁻¹ soil, respectively. The experimental design employed was completely randomized design (CRD) with three replications. An additional replication was included especially for making root VAM colonization studies at flowering/silking. Treatments were absolute control, three VAM fungi cultures [VAM_L (local VAM culture, *Glomus mosseae*, developed by CSK Himachal Pradesh Agricultural University, Palampur), VAM_T (VAM culture, *Glomus intraradices*, developed by Centre for Mycorrhizal Research, the Energy and Resources Institute, New Delhi), and VAM_I (VAM culture, *Glomus mosseae*, developed by Indian Agricultural Research Institute, New Delhi)], and a local PSB culture (*Pseudomonas striata*) alone or in combination with or without 75% of recommended P₂O₅ dose based on targeted yield concept following the soil-test crop response (STCR) precision model as well as one treatment with sole application of 100% of recommended P₂O₅ dose based on the STCR model (Ramamoorthy, Narasimham, and Dinesh 1967). The recommended N and K fertilizers (100%) were added in 75% and 100% doses of recommended P₂O₅ treatments based on the STCR model. Fertilizer N, P, and K were calculated as per targeted yield of the STCR concept (Ramamoorthy, Narasimham, and Dinesh 1967) using following fertilizer adjustment equations:

$$FN = 5.67 T - 0.17 SN$$

$$FP_2O_5 = 4.38 T - 5.26 SP$$

$$FK_2O = 2.29 T - 0.10 SK$$

where FN, FP₂O₅, and FK₂O stand for fertilizer N, P₂O₅, and K₂O in kg ha⁻¹ respectively while T is the targeted yield (40 q ha⁻¹). The SN, SP, and SK are the available N, P, and K statuses of soil respectively in their elemental forms in kg ha⁻¹. Fertilizer calculations were made considering the average weight of 1 ha furrow slice (0–0.15 m) soil to be 2.24 × 10⁶ kg (Suri et al. 2006a). The entire P and K doses were applied basally while N was applied in three equal splits (i.e., at basal, knee–high, and tasseling stages). Nitrogen, P, and K were supplied through urea (46% N), single superphosphate (16% P₂O₅), and muriate of potash (60% K₂O), respectively. Maize seeds were sown in pots

on 10 June 2003. One maize seedling was raised and maintained until maturity. The crop was harvested on 15 September 2003.

VAM Inoculation

Soil mixed VAM cultures with VAM spores and fungal hyphae were used in the study. In all the three VAM cultures [VAM_L (*Glomus mosseae*), VAM_T (*Glomus intraradices*), and VAM_I (*Glomus mosseae*) cultures], the spore count was 100, 500, and 400 per 250 g air-dried soil, respectively. The VAM cultures were used at 12 kg ha⁻¹ (54 mg pot⁻¹). These VAM cultures were used on spore equivalent basis taking VAM_T (*Glomus intraradices*) culture into consideration while using the VAM cultures. Local VAM (*Glomus mosseae*) culture was prepared by the investigating scientists (authors) themselves by raising the target crop (maize) until maturity in pots containing 7 kg sterilized soil + 2 kg FYM and 1 kg mother culture. After harvest, rhizosphere soil of pot as well as root biomass constituted the local VAM (*Glomus mosseae*) culture. Actual inoculation of maize seeds with above VAM cultures was performed by preparing soil slurry of cultures and dipping the seeds into it for half hour followed by shade drying for making seed pallets and then sowing in the pots.

Root Infectivity Studies

After 8–10 weeks of growth at flowering, the plants grown in pots were detached and the rhizosphere soil was carefully removed. The roots were washed thoroughly under running tap water. The cleaned roots were then chopped into small pieces (1 cm) and then subjected to fixation, cleaning, rinsing, and bleaching in potassium hydroxide (KOH) following standard techniques for microscopic observations (Rajapakse and Miller 1992). In case VAM propagules established infection, the infected portion in rhizosphere showed the presence of VAM fructifications. On the basis of root length(s) that were infected and uninfected, the percentage of infection was worked out.

Soil and Plant Chemical Analysis

Soil reaction (pH) was measured in a 1:2.5 soil/water suspension using a glass electrode pH meter (Jackson 1967). Organic C was determined by rapid titration method (Walkley and Black 1934), available N was measured by alkaline permanganate method (Subbiah and Asija 1956), available P was found by Olsen's method (Olsen et al. 1954), and available K was determined by neutral normal ammonium acetate extraction method (AOAC 1970). Plant analysis (grain and stover) was done by the standard procedures of Jackson (1967).

Statistical Analysis

The experimental design was a completely randomized design (CRD), and the statistical analysis was done by the standard procedures suggested by Gomez and Gomez (1984).

Results and Discussion

Root Infectivity and Root Biomass

Maize root system inoculated with various VAM cultures gave greater VAM root infectivity at flowering than their noninoculated counterparts (Table 1). Data further revealed the

Table 1
Effects of co-inoculation of VAM fungi, PSB, and applied P on VAM colonization, root weight, crop productivity, and protein content

Treatments	Root colonization (%)	Root weight (g)	Yield (g/pot)		Grain protein content (%)
			Grain	Stover	
Control	8	3.24	28.07	77.23	8.18
VAM _L alone	12	3.42	31.03	79.87	8.25
VAM _T alone	16	3.56	33.97	83.27	8.37
VAM _I alone	16	3.54	33.23	82.17	8.25
PSB	10	3.52	34.07	84.97	8.31
VAM _L + PSB	16	3.57	33.00	83.07	8.37
VAM _T + PSB	20	3.69	35.10	90.10	8.43
VAM _I + PSB	18	3.60	33.97	88.07	8.37
VAM _L + PSB + 75% P ₂ O ₅ based on STCR model	26	6.79	110.17	269.50	9.25
VAM _T + PSB + 75% P ₂ O ₅ based on STCR model	30	6.98	117.03	292.23	9.37
VAM _I + PSB + 75% P ₂ O ₅ based on STCR model	30	6.88	114.93	285.50	9.31
100% P ₂ O ₅ based on STCR Model	16	7.02	118.00	294.77	9.31
CD (<i>P</i> = 0.05)	—	—	3.19	4.68	0.11

superiority of VAM_T (*Glomus intraradices*) and VAM_I (*Glomus mosseae*) cultures over the local VAM (*Glomus mosseae*) culture with respect to root infectivity. With dual inoculation of VAM cultures and PSB, VAM root colonization was further improved, indicating that there exists a synergistic interaction between VAM and PSB. Gryndler et al. (1990) and Singh and Singh (1993) have also reported similar results. Following the root infectivity pattern, root weight at flowering also increased with individual application of biofertilizers (i.e., VAM and PSB) over the control. Co-inoculation of PSB and VAM culture(s) further affected this parameter favorably. Maximum root weight, however, was recorded with sole application of 100% P₂O₅, which was closely followed by the combined application of VAM cultures and PSB along with 75% of the recommended P₂O₅ dose (Table 1). Thus, greater root biomass coupled with extensive VAM hyphal network in the soil–plant continuum can have great bearing on exploration of soil volume by the crop, leading to higher overall nutrient-use efficiency (George et al. 1992).

Crop Productivity

Like root infectivity, different VAM cultures enhanced the maize grain and stover yield over the control with significantly greater yields only in the cases of VAM_T (*Glomus intraradices*) and VAM_I (*Glomus mosseae*) cultures over the control (Table 1). The individual application of PSB also increased grain and stover yield significantly compared

to the control. Following the co-inoculation of VAM_T (*Glomus intraradices*) + PSB and VAM_I (*Glomus mosseae*) + PSB; yield increases were 21.0%, 18.4%, and 21.4% with respect to grain and 7.8%, 6.4%, and 10.0% with respect to maize stover yield, respectively. Rathore and Singh (1995) have reported similar results. Beneficial effect of VAM inoculation on maize yield is attributed to greater exploration of soil volume by its hyphal network, leading to greater overall nutrient- and water-use efficiencies (George et al. 1992). The VAM are particularly known to induce many favorable changes in the crop rhizosphere by way of exudation/secretion of organic acids / chelating agents (Rovira 1969). These phenomena might have influenced maize yields favorably. The PSB helped in the solubilization of insoluble soil phosphates, thereby encouraging better acquisition of nutrients and strengthening the root system. Application of any of the 3 VAM fungi along with PSB and 75% of recommended P₂O₅ dose resulted in consistent and significant improvement in grain and straw yield of maize with superiority of VAM_T (*Glomus intraradices*) culture, which remained at par with sole application of 100% P₂O₅ dose with regard to grain and stover yield. This indicates that with the dual inoculation of efficient VAM strain and PSB along with 75% P₂O₅ dose, 25% of the fertilizer P dose can be saved without impairing the crop yield targets in maize in an acidic Alfisol of the northwestern Himalayas (Harrier and Watson 2003).

Grain Protein Content

Overall, grain protein content followed a trend similar to that of grain yield (Table 1). There was a significant increase in grain protein content over the control in the case of sole inoculation of VAM_T (*Glomus intraradices*) culture and PSB. It was notable that the combined application of various VAM cultures and PSB was additionally effective in enhancing maize grain protein content, indicating a synergistic interaction between VAM and PSB biofertilizers (Sreenivasa and Krishnaraj 1992). Results, on the whole, suggested that co-inoculation of VAM and PSB coupled with the 75% P₂O₅ dose generated savings of 25% of the fertilizer P in maize.

Nutrient Content and Uptake

Data presented in Table 2 reveal that nutrient (NPK) contents and their uptake in maize grain and stover were enhanced with the sole application of either VAM cultures or PSB inoculation though respective magnitude remained at par with that of the control. However, combined application of VAM cultures and PSB improved the NPK contents and their uptake in maize grain/stover with significant improvement in the case of VAM_T (*Glomus intraradices*) and VAM_I (*Glomus mosseae*) cultures alongwith PSB. Dual inoculation of either VAM_T (*Glomus intraradices*) or VAM_I (*Glomus mosseae*) cultures along with PSB and 75% of the recommended P₂O₅ did not exceed sole application of 100% P₂O₅ in these parameters, but the values tended to be close to each other, which indicates that these biofertilizers have some role in nutrient dynamics in the soil-plant system (Narsian and Patel 2009).

Soil Fertility Status

Co-inoculation of VAM_T (*Glomus intraradices*) or VAM_I (*Glomus mosseae*) cultures along with PSB resulted in significant reductions in available N, P, and K levels in soil after crop harvest in pot experiment (Table 3), mainly because of higher utilization of these

Table 2
Effect of co-inoculation of VAM fungi, PSB, and applied P on nutrient content (NPK) and uptake in maize

Treatments	Nutrient content (%)						Nutrient uptake (mg/pot)					
	N		P		K		N		P		K	
	Grain	Stover	Grain	Stover	Grain	Stover	Grain	Stover	Grain	Stover	Grain	Stover
Control	1.31	0.40	0.13	0.046	0.21	0.40	367.3	308.6	36.3	37.1	58.0	308.5
VAM _L alone	1.32	0.41	0.13	0.050	0.21	0.41	409.6	327.5	41.7	40.0	67.3	327.6
VAM _T alone	1.34	0.42	0.14	0.054	0.23	0.43	455.1	350.1	47.5	45.1	76.7	358.1
VAM _I alone	1.32	0.42	0.14	0.051	0.22	0.42	438.8	345.3	46.7	41.9	73.0	345.1
PSB	1.33	0.42	0.15	0.055	0.23	0.42	452.8	356.9	49.8	47.3	78.3	356.9
VAM _L + PSB	1.34	0.42	0.14	0.056	0.23	0.43	442.2	348.9	46.2	46.5	75.9	357.4
VAM _T + PSB	1.35	0.43	0.16	0.060	0.25	0.44	470.8	390.1	56.5	51.4	89.1	396.2
VAM _I + PSB	1.34	0.43	0.15	0.058	0.24	0.44	455.2	378.7	50.9	51.0	81.5	387.7
VAM _L + PSB + 75% P ₂ O ₅ based on STCR model	1.48	0.51	0.25	0.091	0.33	0.55	1630.6	1373.3	275.4	245.3	363.6	1482.1
VAM _T + PSB + 75% P ₂ O ₅ based on STCR model	1.50	0.53	0.27	0.110	0.35	0.57	1755.6	1548.5	316.1	321.9	409.7	1665.8
VAM _I + PSB + 75% P ₂ O ₅ based on STCR model	1.49	0.53	0.26	0.104	0.34	0.56	1712.5	1513.2	295.0	296.7	390.8	1598.8
100% P ₂ O ₅ based on STCR model	1.49	0.52	0.27	0.118	0.35	0.56	1758.2	1533.0	314.8	347.9	413.2	1641.0
CD (<i>P</i> = 0.05)	0.02	0.02	0.02	0.011	0.03	0.03	49.1	52.7	13.6	13.6	22.7	56.6

Table 3
Effect of co-inoculation of VAM fungi, PSB, and applied P on available nutrient status (mg kg⁻¹ soil) of soil after maize harvest

Treatments	Available nutrient status (mg kg ⁻¹ soil)		
	N	P ₂ O ₅	K ₂ O
Control	34.8	1.7	114.7
VAM _L alone	33.0	1.7	113.4
VAM _T alone	29.9	1.6	111.2
VAM _I alone	31.2	1.6	112.5
PSB	29.9	1.7	111.2
VAM _L + PSB	29.5	1.6	112.5
VAM _T + PSB	27.7	1.6	109.8
VAM _I + PSB	28.6	1.6	110.3
VAM _L + PSB + 75% P ₂ O ₅ based on STCR model	27.7	1.6	104.9
VAM _T + PSB + 75% P ₂ O ₅ based on STCR model	25.9	1.5	102.2
VAM _I + PSB + 75% P ₂ O ₅ based on STCR model	26.8	1.5	103.1
100% P ₂ O ₅ based on STCR model	25.0	1.5	101.3
CD (<i>P</i> = 0.05)	2.2	0.1	1.8
Initial status	49.1	2.2	124.1

nutrients (Table 2) and consequently better crop growth (Table 1). The performance of local VAM culture (*Glomus mosseae*) remained at par with control particularly in terms of soil-available P status. Dual inoculation of biofertilizers further depleted available N, P, and K because of the enhanced exploration of soil reserve nutrients. VAM_T (*Glomus intraradices*) mobilized soil nutrients to the maximum extent followed by VAM_I (*Glomus mosseae*), thereby reducing soil nutrient status significantly over control. This may be ascribed to mobilization of soil N, P, and K from organic and inorganic complexes by the action of VAM fungi due to its mycelial growth or release of organic acids in rhizosphere (Pare, Gregorich, and Nelson 1999). The PSB inoculation also followed the same trend. However, available P in soil under PSB treatment remained at par with the control despite a greater uptake, unlike VAM_T (*Glomus intraradices*) and VAM_I (*Glomus mosseae*) cultures. These facts suggest that PSB inoculation is useful in replenishing the available P status of soil (Suri et al. 2006a). Co-inoculation of PSB with either VAM_T (*Glomus intraradices*) or VAM_I (*Glomus mosseae*) coupled with 75% recommended P₂O₅ dose remained statistically at par with the 100% P₂O₅ dose with respect to soil fertility status. Thus, this study indicates that dual inoculation of PSB and VAM coupled with 75% of the recommended P₂O₅ dose based on targeted yield STCR precision model can bring an economy of about 25% of applied P in maize in a P-deficient acidic Alfisol (Suri et al. 2006).

Conclusions

From this study, it can be inferred that application of VAM fungi alone or with PSB with or without inorganic P fertilization improved the crop productivity, protein content, and nutrient uptake while VAM fungi in combination with PSB as well as 75% of recommended

P₂O₅ dose helped in economizing soil-test and yield-target-oriented P dose using the STCR precision model by about 25% in maize in a P-deficient acidic Alfisol of the northwestern Himalayas. The *Glomus intraradices* culture also showed its superiority over the other two VAM cultures in terms of crop yield, grain quality, and nutrient uptake in maize.

Acknowledgments

The authors are thankful to the Indian Council of Agricultural Research, New Delhi, India, for providing financial assistance for this study under the World Bank-funded National Agricultural Technology Project (NATP). The authors are also thankful to the Centre for Mycorrhizal Research, the Energy and Resources Institute, New Delhi, for supplying VAM cultures and providing valuable technical guidance.

References

- AOAC. 1970. *Official methods of analysis of the Association of Official Agricultural Chemists*. Washington, D.C.: U.S. Government Printing Office.
- Anonymous. 1997. *Soils of Himachal Pradesh for optimizing land use*. Shimla, India: National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur and Department of Agriculture, Government of Himachal Pradesh.
- George, E., K. V. Haussfer, D. Vetterlien, E. Gorgus, and H. Marschner. 1992. Water and nutrient translocations by hyphae of *Glomus mosseae*. *Canadian Journal of Botany* 70:2130–2137.
- Gomez, K. A., and A. A. Gomez. 1984. *Statistical procedures for agricultural research*. New York: Wiley-Interscience.
- Gryndler, M., J. Lestina, V. Moravec, Z. Prikryl, and J. Lipavsky. 1990. Colonization of maize roots by VAM fungi under conditions of long-term fertilization of varying intensity. *Agriculture, Ecosystems and Environment* 29:183–186.
- Harrier, L. A., and C. A. Watson. 2003. The role of arbuscular mycorrhizal fungi in sustainable cropping systems. *Advances in Agronomy* 79:185–225.
- Jackson, M. L. 1967. *Soil chemical analysis*. New Delhi, India: Prentice Hall of India.
- Narsian, V. T., and H. H. Patel. 2009. Relationship of physicochemical properties of rhizosphere soils with native population of mineral phosphate solubilizing fungi. *Indian Journal of Microbiology* 49:60–67.
- Olsen, S. R., C. V. Cole, F. S. Watanabe, and L. A. Dean. 1954. *Estimation of available phosphorus in soils by extraction with sodium bicarbonate* (USDA Circular No. 939). Washington, D.C.: U.S. Government Printing Office.
- Pare, T., E. G. Gregorich, and S. D. Nelson. 1999. Mineralization of nitrogen from crop residues and N recovery by maize inoculated with vesicular arbuscular mycorrhizal fungi. *Plant Science* 218:11–20.
- Rajakpase, S., and J. C. Miller. 1992. Methods for studying vesicular arbuscular mycorrhizae root colonization and related root physical properties. *Methods in Microbiology* 24:301–315.
- Ramamoorthy, B., R. L. Narasimham, and R. S. Dinesh. 1967. Fertilizer recommendations based on fertilizer applications for specific yield targets of Sonara-64. *Indian Farming* 17:443–451.
- Rathore, V. P., and H. P. Singh. 1995. Influence of vesicular arbuscular mycorrhizae fungi and phosphate on maize. *Journal of Indian Society of Soil Science* 43:207–210.
- Rovira, A. D. 1969. Plant root exudates. *Botanical Reviews* 35:35–37.
- Sharma, P. K., S. P. Verma, and D. R. Bhumbla. 1980. Transformation of added phosphorus into inorganic phosphorus fractions in some acid soils of Himachal Pradesh. *Journal of Indian Society of Soil Science* 28:450.
- Singh, H. P., and T. A. Singh. 1993. The interaction of rockphosphate, *Bradyrhizobium*, VAM-corrhizae, and phosphate solubilizing microbes (PSM) on soybean grown in sub-Himalayan Mollisol. *Mycorrhiza* 4:37–43.

- Sreenivasa, M. N., and Krishnaraj, P. U. 1992. Synergistic interaction between VA-mycorrhizal fungi and phosphate bacteria in chilli (*Capsicum annuum*). *Zentralbl fur Microbiologie* 147:126–130.
- Subbiah, B. V., and G. L. Asija. 1956. A rapid procedure for estimation of available nitrogen in soils. *Current Science* 25:259–260.
- Suri, V. K., A. K. Choudhary, G. Chander, and T. S. Verma. 2006. Studies on VAM fungi as a potential biofertilizer in an acid Alfisol of northwestern Himalayas. In *18th WCSS Congress Abstracts*, p. 515. Philadelphia, PA: International Union of Soil Science.
- Suri, V. K., G. Chander, A. K. Choudhary, and T. S. Verma. 2006a. Co-inoculation of VAM and phosphate solubilizing bacteria (PSB) in enhancing phosphorus supply to wheat in Typic Hapludalf. *Crop Research* 31:357–361.
- Walkley, A. J., and C. A. Black. 1934. An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science* 37:29–38.