



Wiley Interdisciplinary Reviews: Energy and Environment

Crop coefficients of *Jatropha* (*Jatropha curcas*) and *Pongamia* (*Pongamia pinnata*) using water balance approach

Kaushal K. Garg, Suhas P. Wani*, A.V.R. Kesava Rao

International Crops Research Institute for the Semi Arid Tropics, Patancheru, Andhra Pradesh, India

DOI: <http://dx.doi.org/10.1002/wene.88>

This is author version post print archived in the official Institutional Repository of ICRISAT www.icrisat.org

Crop coefficients of *Jatropha* (*Jatropha curcas*) and *Pongamia* (*Pongamia pinnata*) using water balance approach

Kaushal K. Garg, Scientist , Suhas P Wani , Principal Scientist, A. V. R. Kesava Rao, Scientist

International Crops Research Institute for the Semi Arid Tropics, Patancheru 502 324, Andhra Pradesh, India

Abstract

Jatropha and Pongamia are potential source of bio-diesel and grow in a wide range of agro climatic zones and soil conditions. Data and knowledge available on water requirement of Jatropha and Pongamia are very scarce. Crop coefficients are important parameters used for assessing water requirement and irrigation scheduling. In present study, crop coefficients of Jatropha and Pongamia were estimated using water balance approach. Temporal data on soil moisture at different depths in block plantations of Jatropha and Pongamia at ICRISAT farm, Patancheru in India were collected at 15 days interval between 2007 and 2010. Measured soil moisture data were analyzed using one dimensional water balance model. Results showed that annual water requirement of Jatropha is 750 mm and of Pongamia is about 950 mm in semi-arid tropics. Crop-coefficients of Jatropha ranged from 0.10 to 0.95 and of Pongamia from 0.30 to 1.10 depending on plant growth stage in different months. ICRISAT received 820 mm of rainfall in a normal year (data between 2001 and 2010) during the monsoon, of which 52 % (430 mm) contributed to evapo-transpiration, 34 % (280 mm) was stored in soil and, 14 % (110 mm) was lost through surface runoff. Stored soil moisture during monsoon season was subsequently utilized by the Jatropha and 270 mm converted into ET during non-monsoonal period. Pongamia utilized stored soil moisture more effectively than Jatropha as it could remove water from deeper soil layers even at high levels of soil moisture suction.

Key words: Bio-diesel, Jatropha, Pongamia, Green water, Semi arid tropics, Water Impact Calculator

INTRODUCTION

Food and energy security are becoming a major issue both for developing and developed countries all over the world. Biofuel such as ethanol and bio-diesel are considered as one of the potential alternatives to reduce fossil fuel dependency^{1,2}, also to reduce greenhouse gas emission and to sequester carbon^{2,3,4}. In recent years, governments of many countries have targeted use of biofuel by blending it with conventional petroleum fuel and are promoted for wide industrial adaptability^{1,5}. In contrary, it is also hypothesized that biofuel production at large scale may threaten the food security by competing with food crops and thus affecting the food supply and prices^{6,7}. On the other hand, production of bio-diesel from plants grown in degraded lands offers opportunities to address the issues of environmental protection and provide options for improving rural livelihood especially in developing countries^{3,8,9}. *Jatropha curcas* (Jatropha) and *Pongamia pinnata* (Pongamia) are identified as potential bio-diesel crops and may be grown in marginal and degraded lands with a wide range of rainfall and ecological conditions^{4,10,11,12}.

Although, the demand for bio-diesel is increasing, available data and knowledge on yield potential and water requirement of *Jatropha* and *Pongamia* are very limited and thus research efforts are required to address such issues¹³. However, few estimates on water requirements of *Jatropha* have been reported in literatures. For example, Rao *et al.*¹⁴ reported that evapotranspiration (ET) from *Jatropha* plantation ranges from 1410 to 1538 mm in optimal conditions and 614-930 mm in field conditions in a semi-arid tropical location in India. They observed moisture extraction from soil layers 150 cm below from surface by five year old plants. Transpiration requirement of *Jatropha* was estimated as 300 mm and 500 mm at 4th and 12th year of growth stage at Owen Sithole College of Agriculture near Empangeni on the KZN north coast and Makhathini, Southern Africa, respectively¹⁵. Gush and Moodley¹⁶ estimated total transpiration of 1983 ℓ (147 mm) for a 4-year-old *J. curcas* tree grown at a relatively wetter location, and 4884 ℓ (362 mm) for a 12-year-old *J. curcas* tree grown at a dry site of South Africa. Transpiration rate was found high during the warm wet summer months and negligible during winter due to the deciduous nature of the *Jatropha* species.

Water requirement of a plant may be estimated by multiplying crop coefficients (K_c) with atmospheric water demand (ET_0). In this method, effect of crop transpiration and evaporation are integrated by a single coefficient, which is known as crop coefficient¹⁷. The K_c - ET_0 approach provides a simple, convenient and reproducible way to estimate water requirement of crops in different climatic conditions^{17,18}. Crop coefficient values have been reported for several crops¹⁷ and tree plantations^{19,20} and have been widely used for irrigation scheduling^{17,21}, simulation studies²² and decision making process¹⁷.

Several methodologies were developed for estimating K_c , for example, weighing lysimeter based techniques²³, agronomic measurements¹⁸ and remote sensing-based techniques²⁴. Irrespective to methodologies, basis for estimating K_c in all such techniques is common *i.e.*, computation of water and energy balance components. In the present study, we used soil moisture data collected at ICRISAT research station for analyzing water balance of *Jatropha* and *Pongamia* crops using one dimensional hydrological model. Specific objectives of the present study are: i) to estimate K_c for *Jatropha* and *Pongamia* crops and ii) to assess water requirements and analyzing field water balance of *Jatropha* and *Pongamia* crops in semi-arid tropics.

MATERIALS AND METHODS

Experimental setup and data monitoring

International Crops Research Institute for the Semi-Arid Tropics (ICRISAT) is located at 17° 53'N, 78° 27'E, and about 545 m above mean sea level in the Medak district of Andhra Pradesh, India. Climate of the study area is tropical monsoon. Long term (1974-2010) average annual rainfall is 898 mm (standard deviation, $\sigma = 225$ mm), of which about 80-85% is received during June to October. Rainfall is highly erratic, both in terms of total amount and distribution over time. Monsoon is preceded by hot summers (minimum air temperature between 16° and 29°C and maximum air temperature between 30° and 43°C in May) and followed by cool winters (minimum air temperature in between 6° and 20°C and maximum air temperature between 23° and 32°C in December). Estimated potential Evapotranspiration (PET) of the study area is 1650 mm annually.

Two micro-watersheds each of 2 ha area were selected for cultivating *Jatropha* and *Pongamia* crops at ICRISAT research station, respectively. Soil of experimental site is classified as Vertisols having maximum depth up to 3 m. Topography of micro-watersheds is relatively flat with average slope between 1.0 and 2.0 %. Soil physical and chemical properties of experimental site e.g. bulk density, texture, organic carbon and soil water retention parameters were measured. Mean values of soil organic carbon was 0.65 % (standard deviation, $\sigma=0.2$) in 0-15 cm depth and 0.50 % ($\sigma=0.10$) in 15-30 cm soil depth. Soil bulk density of upper soil layers is relatively higher (1.4 g cm^{-3} up to 75 cm layer) compared to lower layers (1.2 g cm^{-3} between 120 and 150 cm). Bulk density of surface soil varied between 1.30 and 1.55 with average of 1.35 g cm^{-3} across the watersheds. Subsurface soils are high in clay content and hold more amount (soil moisture availability = 0.25 g/g) of soil moisture compared to shallow soils (soil moisture availability = 0.14 g/g).

Jatropha (3 m × 2 m spacing) and *Pongamia* (5 m × 5 m spacing) seedlings were planted in year 2004. Crops were grown under rainfed conditions and chemical fertilizers were applied every year according to need of plants at its different growth stages. Nitrogen and phosphorus was applied @ 76 kg ha^{-1} and 10 kg ha^{-1} , respectively during 2006 and 2007. During subsequent growth stages, fertilizers dose was increased. Nitrogen and phosphorus was applied @ 92 kg ha^{-1} and 50 kg ha^{-1} , respectively during 2008 and 2009 Source of N and P in above fertilization was urea and single super phosphate, respectively¹⁴. Total 12 neutron probe tubes in each of *Jatropha* and *Pongamia* micro-watersheds were installed for monitoring soil moisture content. Schematic layout of the experimental setup is shown in **Figure 1**

describing spacing and location of neutron probe tubes between the plants rows. Soil moisture content of each 15 cm thick layer was monitored from surface to 225 cm depth using calibrated neutron probe (Troxler 4302) at 15 days interval between 2007 and 2010. In addition, moisture in surface soils (0-15 and 15-30 cm) was also measured by gravimetric method. Daily weather data (rainfall, maximum temperature, minimum temperature, wind speed, solar radiation, relative humidity, and pan evaporation) of experimental site were collected from meteorological station located at 200 m distance from the micro-watersheds. Data on agronomic growth, seed yield and oil content were measured both from the *Jatropha* and *Pongamia* fields.

Description of simulation model

A simple and one dimensional hydrological model (Water Impact Calculator, WIC developed in ICRISAT) was used to analyze field water balance in the present study. WIC requires data on soil (soil depth and water retention), weather (reference evapotranspiration and rainfall), crop growth (biomass and root growth function), topography (land slope, land form conditions), *ex-situ* interventions (water harvesting capacity) and crop management (date of crop sowing and harvesting irrigation method) as an input. WIC analyzes water balance on daily time scale for each one cm soil layer (generate simulation matrix: time vs. soil depth). Rainfall, which is the only source of water for rainfed situation was partitioned into several hydrological components as defined by following mass balance equation:

$$\text{Rainfall} = \text{Surface runoff} + \text{Deep percolation} + \text{Evapotranspiration (Evaporation + Transpiration)} + \text{Change in soil moisture storage} \quad \text{Eq.1}$$

Surface runoff is the flow of excess rainwater after soil becomes saturated and generally occurs either during or after the rainfall event. The SCS curve number technique²⁵ was used to estimate surface runoff in WIC. WIC calculates daily curves number value based on antecedent moisture content of surface soil layer (assumed for 15 cm depth in current analysis) and partition rainfall into surface runoff. Excess amount of infiltrated water after satisfying soil storage capacity is percolated below the crop root zone and is defined as deep percolation. Evapotranspiration comprises two basic hydrological processes: evaporation and transpiration. WIC calculates evaporation and transpiration based on imposed surface boundary conditions and moisture accessibility from surface soil layer and root zone. Water available in top 10 cm soil layer is contributed towards the evaporation demand, whereas, moisture available in root zone is used to meet crop water demand as defined by Allen et al.¹⁷. Crop water requirement (CWR) for a given crop is calculated such as:

$$CWR = K_c \times ET_0 \quad \text{Eq. 2}$$

$$\text{if : } \sum_{i=1}^{\text{rootzone}} AWC > CWR \text{ then } ET = CWR$$

$$\text{otherwise } ET = \sum_{i=1}^{\text{rootzone}} AWC \quad \text{Eq. 3}$$

where AWC is available water and i denotes each cm increment in soil layer reaching up to root zone. It was further assumed that evaporation from soil surface is inversely proportional to vegetative growth/stage. Thus, after achieving full vegetative crop growth ($K_c \geq 1.0$), evaporation from the soil surface is considered almost negligible. Detailed description of WIC, model development, testing and validation procedure are shown by Garg *et al.*²⁶

Estimation of crop coefficients

According to *FAO-56* nomenclature, K_c calculates water requirements under growing conditions having a high level of management and with little or no water stress¹⁷. Appropriate corrections need to be incorporated while estimating crop coefficients in case of water stress situation due to less water uptake. Allen *et al.*¹⁷ described the effects of soil water stress on crop evapotranspiration (ET_c) by multiplying the crop coefficient (K_c) and the water stress coefficient (K_s) with ET_0 such as: $ET_c = K_s \times K_c \times ET_0$

$$\text{Eq. 4}$$

Soils of the experimental watershed are characterized by Vertisols which has medium to high water holding capacity and average depth of 2.5-3.0 m. In current study, K_c is estimated under the fully developed stage (between 2007 and 2011) and is expected to experience with minimal water stress situation. Despite this, whenever available soil moisture depleted below the threshold limit, stress adjustment in ET_c was made as defined by Allen *et al.*¹⁷ and K_c was estimated using Eq. 4. Seventy per cent of total available water (soil moisture) is considered as readily available for crop uptake and further it declined linearly. Water stress coefficient (K_s in Eq.4) is assumed 0.7 for *Jatropha* and *Pongamia* as both are stress tolerant and deep rooted trees^{17,27,28,29}. Crop coefficient of *Jatropha* and *Pongamia* crops were estimated using the inverse optimization technique. Solver is an optimization program built in Microsoft excel (developed for mixed integer nonlinear parameterization) was used for the current study. Based on field measurements, field capacity and permanent wilting point are defined for every 15 cm soil layer up to 300 cm depth. Maximum root zone depth for *Jatropha* and *Pongamia* plant is

considered as 150 cm and 250 cm, respectively.^{27,28,29} Model is set by assigning atmospheric boundary condition (rainfall and PET) on the soil surface as upper boundary and deep percolation from the bottom layer as the lower boundary. Simulation period is started after the four years of plant establishment. Total soil moisture status in root zone was used as the auxiliary variable in the objective function and crop coefficient at different crop growth stages were estimated using inverse optimization technique. WIC performance was assessed by comparing the simulated and measured soil moisture content using the root-mean-squared error (RMSE) of prediction:

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - y_i)^2} \quad \text{Eq. 5}$$

where n is the number of data points of measured (x_i) and simulated (y_i) profile soil moisture.

Estimation of crop water balance

The calibrated model set-up was used to simulate water balance of 10-year time period between 2001 and 2010 capturing large variability of rainfall. Water balance components (ET, runoff and soil moisture) are derived for dry, normal and wet years according to following classification (India Meteorological Department, Pune, India; <http://www.imdpune.gov.in>): Rainfall less than 20% of the long-term average = dry; Rainfall between -20% to +20% of the long-term average = normal; Rainfall greater than 20% of long-term average = wet.

Model further was used to estimate ET under non-limiting water stress situation both for *Jatropha* and *Pongamia* crops by applying auto irrigation setting in WIC between 2001 and 2010. Crop water stress was analyzed by considering difference between actual ET and ET under non-limiting water condition such as:

$$\text{Water Stress Factor (-)} = 1 - \frac{\text{ET actual}}{\text{ET under Non limiting water situation}} \quad \text{Eq. 6}$$

RESULTS AND DISCUSSION

Variation of soil moisture in *Jatropha* and *Pongamia* watersheds

Soil moisture variation in block plantations of *Jatropha* and *Pongamia* are depicted in **Figures 2a** and **2b**, respectively. Figures are divided into four panels and each one shows moisture profile of three consecutive months. In *Jatropha* field, new leaf flushes and biomass growth started by beginning of

April; flowering initiated by May/June; and pod formation and harvesting stage occurred between September and December¹⁴. Previous stored soil moisture into subsurface layers was used by *Jatropha* plants to fulfill water requirements during leaf sprouting. Thus, moisture level depleted significantly between April and June (**Figure 2a**). Large variation in moisture profile is observed between July and September, as this period coincided with peak vegetative growth and occurrence of monsoonal rainfall. Total soil moisture was found highest during September. Moisture in soil profile is depleted slightly during pod formation and harvesting stage during November/December. Variation in profile moisture was found minimal between January and March because crop entered into dormancy phase and leaf started falling down (**Figure 2a**). Soil moisture fluctuation generally was found up to 150 cm indicating maximum effective rooting depth for *Jatropha* crop.²⁷

In *Pongamia* field, low variation in moisture profile was observed during April and June, indicating lesser water uptake (**Figure 2b**). Soil moisture content was increased during monsoon season and depleted subsequently in non-monsoonal period as expected. Data showed that *Pongamia* being a perennial crop could extract water from deeper and drier soil conditions. Several other studies indicated that *Pongamia* has an extensive lateral root system and thick-long tap root²⁸, which could be developed maximum up to 5-10 m length at fully matured stage²⁹. Soil moisture level in *Pongamia* cultivated field is depleted much higher compared to *Jatropha* cultivated fields. For example, soil moisture content for *Jatropha* and *Pongamia* cultivated field in month of May-June was recorded as 0.2-0.3 cm³ cm⁻³ and 0.03-0.15 cm³ cm⁻³, respectively.

Model performance

Soil moisture measured in *Jatropha* and *Pongamia* fields is compared with model simulated moisture level between 2007 and 2011. Simulated soil moisture fluctuations are found comparable with observed numbers at most of the times both for *Jatropha* and *Pongamia* cultivated fields. Root mean square error in estimating profile moisture is 6.0 cm and 5.0 cm, which is equivalent to less than 10 % of total moisture in given soil profiles, thus model performance is considered acceptable.

Crop coefficients and crop water requirements

Crop coefficients estimated for *Jatropha* and *Pongamia* crops are summarized in **Table 1**. K_c for *Jatropha* and *Pongamia* are found in range between 0.10-0.95 and 0.30-1.10, respectively. Maximum K_c was found during August and September for both the crops.

Water balance components of Jatropha and Pongamia fields are summarized in **Table 2**. During normal years, rainfall recorded was 911 mm (820 mm in monsoon and 91 mm in non-monsoon) at experimental site. In these years, ET from Jatropha and Pongamia crop were estimated to be 685 and 830 mm year⁻¹, respectively. Surface runoff generated from both the crop lands is found almost similar, i.e., 130-135 mm (15 % of total rainfall). In Jatropha field, nearly 100 mm rainfall was harvested in terms of green water. In Pongamia fields, 35 mm of net deficit in green water storage was observed by end of the crop season in normal years.

ET from Jatropha and Pongamia field was found larger than the rainfall received during dry years. About, 130-150 mm harvested green water in previous years was utilized by crops during dry years as shown by negative balance (**Table 2**). Surface runoff and deep percolation volume were estimated to be 90-100 mm, i.e., 14% of rainfall received. Results showed that poor rainfall in dry years led to develop water shortage by 21 % (in terms of ET) than the required quantity for Pongamia crop (shown by water stress factor in Table 2). On the other hand, crop water demand and availability for Jatropha crop was found in comparable range.

To show a detail description of mass balance, the monthly water balance of Jatropha and Pongamia field is presented for one of the selected normal year (**Figure 3**). Total 850 mm of rainfall was received during monsoon in year 2003. Thirty four per cent of monsoonal rain was captured as soil moisture, 52 % (440 mm) contributed to ET and 14 % (120 mm) was exported as surface runoff. In non-monsoon period (Nov-May), harvested green water acted as source of water as shown in **Figure 3**. Results showed that water requirements of both the crops were nearly same during monsoon period (460 mm) but Pongamia required additional 200 mm water during non-monsoon season compared to Jatropha (240 mm). June is found the critical period for Jatropha growth and supplemental irrigation could be helpful in dry years.

Conclusions

Jatropha and Pongamia are potential bio-diesel crops and could be grown in wide range of agro-climatic and soil conditions. Water requirement of these two crops are estimated in the current study. Key findings of this study are:

1. Water requirements of Jatropha and Pongamia are estimated to be 750 and 950 mm in semi-arid tropics, respectively. Water balance showed that at ICRISAT out of 820 mm rainfall received

during monsoon period, 52 % was contributed to ET, 34 % stored as soil moisture and 14 % exported outside from watershed as surface runoff. Consumptive water use by Jatropha and Pongamia are found to be nearly same in monsoon, i.e., 460-500 mm. Water requirement (ET) of Jatropha during non-monsoon period is found as 265 mm. Pongamia required 200 mm additional water than Jatropha during non-monsoonal period.

2. Crop coefficients of Jatropha and Pongamia ranged from 0.10-0.95 and 0.30-1.1 depending on growth stage, respectively. Pongamia is found to utilize green water from deeper and drier soil layers compared to Jatropha crop.
3. Water availability for Pongamia crops was found adequate in wet years but slightly less (10% shortage) in normal year. Poor rainfall in dry years led to develop water shortage by 20 % than the required quantity in Pongamia field, whereas Jatropha experienced negligible water stress in all the years. *In-situ* moisture availability played an important role in supplying water both for Jatropha and Pongamia crops especially in non-monsoon period.

Crop coefficient estimated in current study could be useful to assess technical feasibility to grow Jatropha and Pongamia crops in different agro-ecological regions and soil conditions.

References

1. Government of India. National Policy on Biofuels. Ministry of New & Renewable Energy 2009 New Delhi.
2. Achten WMJ, Maes WH, Aerts R, Verchot L, Trabucco A, Mathijs E, Singh VP, Muys B. Jatropha: From global hype to local opportunity. *Journal of Arid Environments* 2010, 74:164–165.
3. Wani SP, Sreedevi TK, Marimuthu S, Rao AVRK, Vineela C. Harnessing the potential of Jatropha and Pongamia plantations for improving livelihoods and rehabilitating degraded lands. 6th International Biofuels Conference 2009, March 4-5, New Delhi, India.
4. Wani SP, Chander G, Sahrawat KL, Rao CS, Raghvendra G, Susanna P, Pavani M. Carbon sequestration and land rehabilitation through Jatropha curcas (L.) plantation in degraded lands. *Agriculture, Ecosystems and Environment* 2012, 161: 112– 120.
5. EC. Communication from the European Commission on An EU Strategy for Biofuels, COM 2006, 34 final. European Commission,
6. Doornbosch R, Steenblik R. Round Table on Sustainable Development Biofuels: Is the cure worse than the disease? Organization for Economic Cooperation and Development (OECD) 2007, 11-12 September.
7. Naylor R, Steinfield H, Falcon W, Galloway J, Smil V, Bradford E, Alder J, Mooney H. Losing the Links between Livestock and Land. *Science* 2007, 310: 1621-1622.
8. Wani SP, Sreedevi TK. Pongamia's journey from forest to microenterprise for improving livelihood, International Crop Research Institute for the Semi-Arid Tropics. Research Report 2005 Global Theme of Agroecosystems, Patancheru, India.

9. Sreedevi TK, Wani SP, Srinivasa Rao CH, Chaliganti R, Reddy RL. *Jatropha* and *Pongamia* rain-fed plantations on wastelands in India for improved livelihoods and protecting environment. 6th International Biofuels Conference 2009, March 4-5 New Delhi, India.
10. Wani SP, Osman M, D'Silva E, Sreedevi TK. Improved livelihoods and environmental protection through biodiesel plantations in Asia. *Asian Biotechnology and Development Review* 2006 8(2): 11-29.
11. Achten WMJ, Verchot L, Franken YJ, Mathijs E, Singh VP, Aerts R. Review: *Jatropha* bio-diesel production and use. *Biomass & Bioenergy* 2008, 32: 1063–1084.
12. Garg KK, Karlberg L, Wani SP, Berndes G. *Jatropha* production on wastelands in India: opportunities and trade-offs for soil and water management at the watershed scale. *Biofuels, Bioproducts & Biorefining* 2011, 5: 410–430.
13. Raju KV. Biofuels in South Asia: An overview, editorial introduction. *Asian Biotechnology and Development Review* 2006: 8(2): 1-9.
14. Rao AVRK, Wani SP, Singh P, Srinivas K, Srinivasa Rao Ch. Water requirement and use by *Jatropha curcas* in a semi-arid tropical location. *Biomass & Bioenergy* 2012, 39: 175-181.
15. Holl MA, Gush MB, Hallows J, Versfeld DB. *Jatropha curcas* in South Africa: An Assessment of its Water Use and Bio-Physical Potential. Report submitted to the Water Research Commission on the project "Investigation into the Impacts of Large-Scale Planting of *Jatropha curcas* on Water Resources, through Process-Based Research and Modelling". WRC Report No 1497/1/07, NOVEMBER 2007.
16. Gush MB and Moodley M. Water use assessment of *Jatropha curcas*. In: Holl M, Gush MB, Hallows J and Versfeld DB (eds.) *Jatropha curcas* in South Africa: An Assessment of its Water Use and Bio-Physical Potential. WRC Report No. 1497/1/07 (Chapter 4). Water Research Commission, Pretoria, RSA
17. Allen RG, Pereira LS, Raes D, Smith M. Crop evapotranspiration: Guidelines for computing crop water requirements, Irrigation and Drainage Paper 56. United Nations FAO, Rome, 300 p. <http://www.fao.org/docrep/X0490E/X0490E00.htm> 1998
18. Allen RG, Pereira LS. Estimating crop coefficients from fraction of ground cover and height. *Irrigation Science* 2009, 28: 17–34.
19. Teixeira AH, de C, Bastiaanssen GM, Moura MSB, Soares JM, Ahmad MD, Bos MG. Energy and water balance measurements for water productivity analysis in irrigated mango trees, Northeast Brazil. *Agricultural and Forest Meteorology* 2008, 148: 1524–1537.
20. Paco TA, Ferreira MI, Rosa RD, Paredes P, Rodrigues GC, Conceicao N, Pacheco CA, Pereira LS. The dual crop coefficient approach using a density factor to simulate the evapotranspiration of a peach orchard: IMDualKc model versus eddy covariance measurements. *Irrigation Science* 2012, 30: 115–126.
21. Chen F, Robinson PJ. Estimating reference crop evapotranspiration with ET gauges. *Journal of Irrigation and Drainage Engineering* 2009, 135 (3): 335-342.
22. Cancela JJ, Cuesta TS, Neira XX, Pereira LS. Modelling for improved irrigation water management in a temperate region of Northern Spain. *Biosystems Engineering* 2006, 94(1): 151–163.

23. Evett SR, Mazahrih NT, Jitan MA, Sawalha MH, Colaizzi PD, Ayars JE. A weighing lysimeter for crop water use determination in the Jordan Valley, Jordan, *Transactions of the ASABE* 2009, 52(1):155-169.
24. Singh RK, Irmak A. Estimation of Crop Coefficients Using Satellite Remote Sensing. *Journal of Irrigation and Drainage Engineering* 2009, 135: 597-608.
25. USDA-SCS. 1985. National Engineering Handbook, Section 4 - Hydrology. Washington, D.C.: USDA-SCS.
26. Garg KK, Wani SP, Patil MD. Simple and farmer-friendly decision support system for enhancing water use efficiency in agriculture: tool development, testing and validation. *Submitted to "Agricultural water management"* 2013.
27. Krishnamurthy L, Zaman-Allah M, Marimuthu S, Wani SP, Rao AVRK. Root growth in Jatropha and its implications for drought Adaptation. *Biomass & Bioenergy* 2012, 39: 247-252.
28. Scott P T, Pregelj L, Chen N, Hadler J S, Djordjevic M A, Gresshoff P M. Pongamia pinnata: An Untapped Resource for the Biofuels Industry of the Future Bioenerg. Res 2008 1:2–11
29. Rangan L. *Pongamia* - A multipurpose versatile legume *Research Journal of Biotechnology* 2013. 8(1): 1-3.

List of Tables

Table 1: Crop coefficient for Jatropha and Pongamia tree crops on monthly time scale

Month	K _c of Jatropha	K _c of Pongamia
April	0.20	0.30
May	0.40	0.40
June	0.60	0.40
July	0.85	0.70
August	0.90	0.90
September	0.95	1.10
October	0.80	1.10
November	0.70	1.10
December	0.50	0.75
January	0.30	0.65
February	0.10	0.60
March	0.10	0.35

Table 2: Water balance of Jatropha and Pongamia cultivated fields at ICRISAT for 10 years simulation period

Crop cultivated	Year	No of year	Water balance components						Water stress factor (-)
			Rainfall (mm)			Surface runoff + Deep percolation (mm)	ET (mm)	Change in SMC (mm)	
			Jun-Oct	Oct-May	Annual rainfall				
Jatropha	Dry	2	612	40	652	90	690	-128	0.01
Jatropha	Normal	6	820	91	911	130	700	80	0.01
Jatropha	Wet	2	1179	71	1250	322	808	118	0.00
Pongamia	Dry	2	612	40	652	95	710	-153	0.21
Pongamia	Normal	6	820	91	911	115	830	-35	0.10
Pongamia	Wet	2	1179	71	1250	249	953	48	0.00

The calibrated model set-up is used to simulate water balance for a 10-year time period between 2001 and 2010. Water balance results are presented for dry, normal and wet years according to following classification (Indian Meteorological Department, Pune, India; <http://www.imdpune.gov.in>): Rainfall less than 20% of the long-term average = dry; Rainfall between -20% to +20% of the long-term average = normal; Rainfall greater than 20% of long-term average = wet;

List of Figures




Figure 1: Layout of neutron probe access tubes for soil moisture measurement in Jatropha and Pongamia field at ICRISAT, Patancheru.




Figure 2a:




Figure 2b:

Figure 2a Depth wise soil moisture content during 12 months in Jatropha experimental field

Figure 2b Depth wise soil moisture content during 12 months in Pongamia experimental field




Figure 3 Monthly simulated water balance of Jatropha and Pongamia fields at ICRISAT for a selected normal year (2003-2004); the upper part of the graphs show the source of the water (rainfall and change in soil moisture content), and the lower part presents various sink components: ET, surface runoff, deep percolation, change in soil moisture content