

1336 CP522

Genotypic diversity in pearl millet (*Pennisetum glaucum*) for nitrogen, phosphorus and potassium use efficiencies*

S.P. WANI, M.A. ZAMBRE and K.K. LEE

International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru P.O. 502 324, A.P., India

Key words: genetic diversity, grain yield, harvest index, nutrient uptake, pearl millet, *Pennisetum glaucum* (L.) R.Br., translocation index

Abstract

Twelve genotypes of pearl millet (*Pennisetum glaucum* (L.) R.Br. comprising of hybrids, composites, varieties, and landraces were evaluated for N, P, and K uptake, efficiency of grain production per unit of N, P, and K absorbed and their efficiency of transfer from vegetative parts to the grain. The genotypes were grown on Allisols in two fields at the ICRISAT Centre with two N and P levels in a rainy season (20 kg N + 9 kg P ha⁻¹ and 80 kg N + 18 kg P ha⁻¹).

Genotypes varied significantly for total dry matter, grain yield, and for uptake, use efficiency and translocation indices of N, P, and K. Genotype × fertility interactions were not observed for all parameters studied, except for grain yield, indicating that genotypes could be evaluated for N, P, and K use efficiency and translocation indices at different soil fertility levels. Hybrid MBH 110 showed the highest use efficiency for N, P, and K with a maximum harvest index, and it was followed by two other hybrids amongst the genotypes tested. Genotypes (composites, and a landrace) which showed lower use efficiency and translocation indices for N, P, and K also had lower grain yields than the hybrids tested. Positive relationships were found between harvest index and phosphorus use efficiency, and N, and P translocation indices. Thus, a future challenge lies in selecting lines with high N, P, and K use efficiency and translocation indices.

Introduction

Pearl millet is an important rainfed cereal crop grown on marginal soils in the semi-arid tropics. Most of it is grown with little or without fertilizer applied. Nitrogen is usually the nutrient limiting crop production and its poor recovery by crops, when applied as fertilizer, is of worldwide concern. With steadily increasing prices of fertilizers, it becomes important to produce maximum pearl millet grain yields per unit of fertilizer applied. In recent years, crop improvement research specifically directed towards increasing

the efficiency of mineral nutrition of plants has received increased attention (Devine, 1982; Gabelman and Loughman, 1987; Sarić and Loughman, 1983). The differential response of pearl millet genotypes to applied N (ICRISAT, 1988; Kanwar *et al.*, 1973; Murty, 1967) suggests that differences in nutrient uptake and translocation to the grains exist in pearl millet genotypes. Nutritional differences in genotypes have rarely been related to final economic yield. Increased and efficient production of pearl millet may be characterised as functions of increased uptake and accumulation of nutrients by the plant, increased production of dry matter per unit of nutrient assimilated, and increased translocation of nutrients from vegetative parts to the grain.

* Submitted as CP # 522 by the International Crops Research Institute for the Semi-Arid Tropics (ICRISAT).

Genotypes differences with respect to N, P, K and other nutrients for the abovementioned traits have been reported in sorghum (Seetharama *et al.*, 1987). This paper deals with differences among a set of 12 pearl millet genotypes grown at two levels of applied N and P fertility. The genotypes were compared for the extent of variation in the abovementioned traits and for the relationships between grain yield and some of these traits.

Materials and methods

Experimental details

Twelve pearl millet genotypes selected for this study consisted of three hybrids (BJ 104, MBH 110, and ICMH 451) released for commercial cultivation in India, five composites (MC-C8, D2 C6, ERC-Co, EC-C6 and IVC-C7) made at ICRISAT, one composite (RCB 2) released for cultivation in Rajasthan, India, a synthetic (Gam 73) from Senegal, and Local landraces from India (Rajasthan Locals 1 and 2). The experiment was conducted in two fields (locations) (Table 1) during the rainy season on Alfisols at ICRISAT Center, Patancheru, India (17°36'N, 78°16'E, 545 m altitude).

Both experiments were conducted in a split-plot design with two N fertility levels as main plots and pearl millet genotypes as sub-plots.

Table 1. Details of Alfisols at ICRISAT Center, 1987 rainy season, and of field experiments

Properties	Location I	Location II
Soil pH	6.60	7.00
EC (M mhos cm ⁻¹)	0.19	0.24
Organic carbon (g kg ⁻¹)	3.40	3.30
Available P (mg kg ⁻¹)	14.7	14.3
NO ₃ -N + NH ₄ ⁺ -N (mg kg ⁻¹)	19	20.0
Total N (mg kg ⁻¹)	410	410
Gross plot area (m ²)	40.5	24
Harvest area (m ²)	36	13.5
Date of sowing	17 June '87	16 June '87
Date of irrigation	-	22 July and 4 August '87

Each treatment was replicated four times. For the 20 kg N ha⁻¹ treatment 20 kg N ha⁻¹ as urea and 9 kg P ha⁻¹ as single super-phosphate were applied as basal dressings. For the 80 kg N ha⁻¹ treatment a basal dose of 40 kg N ha⁻¹ and 18 kg P ha⁻¹ was applied. The remaining N was applied as urea after thinning (20 DAS). The crop was machine sown on ridges spaced 0.75 m apart, and plant-to-plant spacing of 0.1 m was maintained by thinning the plants 12 DAS. Weeding and inter-row cultivations were carried out as and when required.

At harvest, the above-ground plant parts were harvested. The panicles were separated and threshed. Fresh stover yield was recorded and 10-kg subsamples of stover were collected and chopped. The subsampled stover biomass and the grains were dried at 70°C for 72 h and their dry mass recorded. N and P in ground grains and in stover were determined with the use of a Technicon Autoanalyser, and K was determined using atomic absorption spectrophotometry (Jackson, 1973).

Characteristics used to compare genotypes

The traits used to characterise the genotypes were i) total dry matter production (grain plus stover), ii) harvest index [HI: (grain mass/total dry matter mass × 100)], iii) N, P and K contents of grain and stover (dry mass multiplied by concentrations in the respective parts), iv) total N, P, and K in plants (contents in grains + contents in stover), v) N, P, and K use efficiency (NUE, PUE and KUE: grain mass produced per unit of N, P, and K in total dry matter), vi) N, P, and K translocation indices (NTI, PTI and KTI: N, P, and K contents in grains divided by N, P, and K contents in total dry matter × 100).

Results

Grain and total dry matter yield, and total N, P, and K uptake

Mean grain and total dry matter yields of pearl millet across the genotypes and locations increased significantly to 3.10 t ha⁻¹ and 7.90 t ha⁻¹




Fig 1 a Relationship between total dry matter yield (kg ha^{-1}) and grain yield (kg ha^{-1}) of pearl millet genotypes. $y = 84 + 0.380 \times (R^2 = 0.67^{**}, \text{df} = 190)$ b Relationship between grain yield (kg ha^{-1}) and harvest index (HI %) of pearl millet genotypes $y = 472 + 60.10 \times (R^2 = 0.28^{**}, \text{df} = 190)$ c Relationship between phosphorus use efficiency (PUE) and harvest index (HI) of pearl millet genotypes. $y = 8.64 + 0.162 \times (R^2 = 0.48^{**}, \text{df} = 190)$ d. Relationship between harvest index (HI) and nitrogen translocation index (NTI) of pearl millet genotypes. $y = 1.41 + 0.589 \times (R^2 = 0.44^*, \text{df} = 190)$ e Relationship between harvest index (HI) and phosphorus translocation index (PTI) of pearl millet genotypes $y = 7.86 + 0.478 \times (R^2 = 0.38^{**}, \text{df} = 190)$.

Table 2. Grain and total dry matter yield, nutrient uptake, concentrations in grains and plant dry matter, nutrients use efficiency and translocation index of pearl millet genotypes^a

Genotypes	Genotypes											CV %		
	BJ 104	MBH 110	MC-C8	D2-C6	ERC-CO	ERC-C6	IVC-C7	Gam 73	ICMH 451	Local 1	RCB2		Local 2	SE ±
Grain yield (t ha ⁻¹)	2.8	3.3	3.1	2.9	2.6	2.8	3.1	2.4	3.6	2.5	2.4	2.5	0.097 ^{a,b}	14
Total dry matter yield (t ha ⁻¹)	6.6	6.8	8.1	7.0	7.6	6.9	8.2	6.6	8.5	7.2	6.9	6.4	0.176 ^a	13
Nutrient uptake (kg ha ⁻¹)														
Nitrogen	69	74	72	70	64	67	74	70	81	70	68	67	2.8 ^a	21
Phosphorus	13.5	15.0	16.3	16.0	14.9	15.2	16.6	14.3	17.6	14.0	13.9	13.3	0.47 ^a	16
Potash	124	107	138	130	115	119	127	125	151	130	112	105	6.3 ^a	29
Harvest index	43	48	38	43	34	41	38	36	42	34	35	40	1.08 ^a	11
Nitrogen concentration (mg N g ⁻¹)														
Grain	15.8	16.2	14.1	14.3	15.2	14.7	14.9	15.4	14.7	17.0	16.8	17.0	0.26 ^a	12
Stover	6.1	5.3	5.3	6.4	4.6	5.9	5.3	7.6	5.3	5.5	5.9	5.9	0.20 ^a	20
Phosphorus concentration (mg P g ⁻¹)														
Grain	3.3	3.4	3.4	3.4	3.6	3.4	3.5	3.4	3.5	3.7	3.7	3.5	0.04 ^a	5
Stover	1.1	1.1	1.1	1.5	1.1	1.4	1.1	1.5	1.0	1.0	1.1	1.2	0.04 ^a	23
Potash concentration (mg K g ⁻¹)														
Grain	4.9	4.7	4.6	5.1	4.5	4.6	4.7	5.0	5.0	5.0	4.8	4.7	0.12 ^a	10
Stover	28.7	25.6	24.4	27	20.7	25.0	21.7	25.9	25.9	23.5	22.2	23.2	1.01 ^a	20
Nutrient use efficiency (kg grain produced kg ⁻¹ nutrient uptake)														
Nitrogen	44	48	47	47	44	46	45	37	48	39	39	40	1.02 ^a	15
Phosphorus	209	219	191	185	174	186	187	168	205	178	174	191	3.64 ^a	11
Potash	32	40	29	30	29	33	32	25	31	25	28	35	2.37 ^a	34
Nutrient translocation index (%)														
Nitrogen	66	74	63	63	63	65	63	55	67	62	62	66	1.12 ^a	8
Phosphorus	69	74	65	62	62	64	65	57	72	66	64	66	1.21 ^a	9
Potash	16	19	14	17	14	15	16	12	15	12	12	15	1.16 ^a	35

^a Each value is the average of two locations, two N levels and four replications.

^b ** P < 0.01.

^c * P < 0.05.

respectively, with 80 kg N ha⁻¹ applied as compared to 2.55 t ha⁻¹ and 6.50 t ha⁻¹ respectively, with 20 kg N ha⁻¹ applied. In all genotypes, except D2C6' and IVC-C7, application of 80 kg N ha⁻¹ increased the grain yields significantly over those obtained with 20 kg N ha⁻¹ treatment. However, for total dry matter, the genotype × fertility level interactions were not significant. Even though grain and dry matter yields varied between locations, there were no genotype × location interaction effects for these parameters. Mean grain and total dry matter yields of pearl millet genotypes across the fertility levels and locations varied significantly. The landraces from India (Rajasthan Locals 1 and 2) produced dry matter yields similar to those of improved varieties and hybrids (Table 2). A maximum mean grain yield of 3.6 t ha⁻¹ was observed for hybrid ICMH 451, followed by another hybrid MBH 110. Total N, P, and K uptake varied significantly (Table 2). However, total N, P, and K uptake by landraces and some of the improved genotypes were similar. There was a positive relationship (df = 190) between grain mass and total dry matter ($R^2 = 0.67^{**}$, Fig. 1a), grain mass and HI ($R^2 = 0.28^{**}$, Fig. 1b), grain mass and total N ($R^2 = -0.53^{**}$), P ($R^2 = 0.71^{**}$) and K ($R^2 = 0.31^{**}$), uptake respectively.

Harvest index and nutrient translocation indices

Application of 80 kg N ha⁻¹ or field location (or fertility treatment) had no effects on HI of pearl millet genotypes. The HI varied significantly amongst genotypes. The improved hybrids and the composites, except ERC-Co (an ergot-resistant composite), had higher HI than the local landraces (Table 2). Harvest index was positively correlated with PUE ($R^2 = 0.48^{**}$, df = 190, Fig. 1c).

NTI of pearl millet genotypes did not change with 80 kg N ha⁻¹ applied in comparison with 20 kg N ha⁻¹. However, NTI varied significantly among locations, but without genotype × location interactions. Mean PTI of pearl millet across the locations increased significantly to 68.2% when N was increased from 20 to 80 kg ha⁻¹, whereas in that situation mean KTI decreased from 16.5% to 13%. Nutrient translo-

cation indices varied significantly amongst genotypes. In general, the genotypes with higher HI showed higher nutrients translocation indices (Table 2). Higher proportions (>60%) of plant N and P were translocated to the grains, whereas <80% of total plant K remained in the stover. More variability for NTI and PTI than for KTI was observed (Table 2). There was a positive relationship (df = 190) between HI and NTI ($R^2 = 0.44^{**}$, Fig. 1d) and also with PTI ($R^2 = 0.38^{**}$, Fig. 1e).

Nutrient (N, P, K) concentrations in grain and stover

Mean N concentrations in grain and stover across the genotypes and locations increased significantly from 14 to 17 and from 5.1 to 6.4 mg g⁻¹, respectively, with 80 kg N ha⁻¹ instead of 20 kg N ha⁻¹ applied. Phosphorus concentration in grains was not changed, whereas P concentration declined significantly from 1.3 mg g⁻¹ to 1.07 mg g⁻¹ dry matter when N application was increased from 20 to 80 kg ha⁻¹. Potassium concentration in grains decreased from 4.93 to 4.66 mg g⁻¹ when N applied was increased from 20 to 80 kg ha⁻¹, whereas K concentration in stover was not changed. Locations also had a significant effect on nutrient concentrations in grains and stover.

Mean N, P, and K concentrations in grain and stover varied significantly with genotypes. The landraces had higher concentrations in the grains than the improved genotypes. However, N, P, and K concentrations in stover of local landraces were similar to those in stover of improved genotypes, except for BJ 104 and D2C6 (Table 2). There was no genotype × fertility interaction for N concentration in grain and stover, but a significant interaction was observed between genotypes and fertility levels for P and K concentrations in grains and stover. In a few genotypes P and K concentrations in grain and stover increased when N applied was increased from 20 to 80 kg ha⁻¹, whereas in others it decreased. Grain N concentration was positively correlated (df = 190) with a total N uptake ($R^2 = 0.71^{**}$), and also with total K uptake ($R^2 = 0.50^{**}$) while it was negatively correlated with NUE ($R^2 = 0.83^{**}$), and KUE ($R^2 = 0.34^{**}$).

